



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2017 – 10:43 AM EDT

PDB ID : 3HYL
Title : Crystal Structure of Transketolase from Bacillus anthracis
Authors : Maltseva, N.; Kim, Y.; Kwon, K.; Joachimiak, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : unknown
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	690	 2% 75% 19% . .
1	B	690	 4% 67% 25% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	667	-	-	-	X
3	FMT	A	669	-	-	X	-
3	FMT	A	673	-	-	X	X
3	FMT	B	667	-	-	-	X
3	FMT	B	668	-	-	X	-
3	FMT	B	674	-	-	X	X
4	PEG	A	674	-	-	-	X
4	PEG	A	677	-	-	X	X
4	PEG	A	678	-	-	-	X
4	PEG	B	669	-	-	-	X
7	SO4	B	671	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 10868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transketolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	663	Total	C	N	O	S	Se	0	2	0
			5087	3198	862	1005	1	21			
1	B	663	Total	C	N	O	S	Se	0	3	0
			5096	3205	863	1006	1	21			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MSE	-	EXPRESSION TAG	UNP C3P4P9
A	-22	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-21	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-20	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-19	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-18	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-17	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-16	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-15	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-14	GLY	-	EXPRESSION TAG	UNP C3P4P9
A	-13	VAL	-	EXPRESSION TAG	UNP C3P4P9
A	-12	ASP	-	EXPRESSION TAG	UNP C3P4P9
A	-11	LEU	-	EXPRESSION TAG	UNP C3P4P9
A	-10	GLY	-	EXPRESSION TAG	UNP C3P4P9
A	-9	THR	-	EXPRESSION TAG	UNP C3P4P9
A	-8	GLU	-	EXPRESSION TAG	UNP C3P4P9
A	-7	ASN	-	EXPRESSION TAG	UNP C3P4P9
A	-6	LEU	-	EXPRESSION TAG	UNP C3P4P9
A	-5	TYR	-	EXPRESSION TAG	UNP C3P4P9
A	-4	PHE	-	EXPRESSION TAG	UNP C3P4P9
A	-3	GLN	-	EXPRESSION TAG	UNP C3P4P9
A	-2	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-1	ASN	-	EXPRESSION TAG	UNP C3P4P9
A	0	ALA	-	EXPRESSION TAG	UNP C3P4P9
B	-23	MSE	-	EXPRESSION TAG	UNP C3P4P9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-21	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-20	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-19	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-18	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-17	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-16	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-15	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-14	GLY	-	EXPRESSION TAG	UNP C3P4P9
B	-13	VAL	-	EXPRESSION TAG	UNP C3P4P9
B	-12	ASP	-	EXPRESSION TAG	UNP C3P4P9
B	-11	LEU	-	EXPRESSION TAG	UNP C3P4P9
B	-10	GLY	-	EXPRESSION TAG	UNP C3P4P9
B	-9	THR	-	EXPRESSION TAG	UNP C3P4P9
B	-8	GLU	-	EXPRESSION TAG	UNP C3P4P9
B	-7	ASN	-	EXPRESSION TAG	UNP C3P4P9
B	-6	LEU	-	EXPRESSION TAG	UNP C3P4P9
B	-5	TYR	-	EXPRESSION TAG	UNP C3P4P9
B	-4	PHE	-	EXPRESSION TAG	UNP C3P4P9
B	-3	GLN	-	EXPRESSION TAG	UNP C3P4P9
B	-2	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-1	ASN	-	EXPRESSION TAG	UNP C3P4P9
B	0	ALA	-	EXPRESSION TAG	UNP C3P4P9

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0


- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total O S 5 4 1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	343	Total O 343 343	0	0
8	B	261	Total O 261 261	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.80 Å 70.97 Å 145.79 Å 90.00° 117.35° 90.00°	Depositor
Resolution (Å)	36.35 – 2.16 36.35 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.4 (36.35-2.16) 99.4 (36.35-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.36 (at 2.16 Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.176 , 0.230 0.172 , 0.228	Depositor DCC
R_{free} test set	3417 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10868	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, CL, FMT, SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/5174	0.77	4/6988 (0.1%)
1	B	0.64	0/5187	0.76	6/7005 (0.1%)
All	All	0.67	0/10361	0.76	10/13993 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	ARG	NE-CZ-NH2	-10.79	114.90	120.30
1	A	93	ARG	NE-CZ-NH1	10.20	125.40	120.30
1	B	93	ARG	NE-CZ-NH2	-7.87	116.37	120.30
1	A	665	MSE	CG-SE-CE	-6.16	85.36	98.90
1	B	31	MSE	CG-SE-CE	-5.65	86.46	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5087	0	4921	134	0
1	B	5096	0	4930	198	0
2	A	12	0	16	2	0
3	A	15	0	5	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	18	0	6	7	0
4	A	21	0	30	7	0
4	B	7	0	10	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
7	B	5	0	0	0	0
8	A	343	0	0	5	0
8	B	261	0	0	12	0
All	All	10868	0	9918	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:ASP:HB2	8:B:866:HOH:O	1.26	1.30
1:B:616:MSE:SE	8:B:741:HOH:O	2.12	1.16
1:B:465:ILE:HD11	1:B:616:MSE:HE2	1.26	1.12
1:B:616:MSE:HE3	1:B:653:PHE:CD2	1.85	1.12
1:A:130:MSE:SE	8:A:927:HOH:O	2.21	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	663/690 (96%)	637 (96%)	25 (4%)	1 (0%)	51	50
1	B	664/690 (96%)	634 (96%)	30 (4%)	0	100	100
All	All	1327/1380 (96%)	1271 (96%)	55 (4%)	1 (0%)	55	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	528/527 (100%)	492 (93%)	36 (7%)	18	12
1	B	529/527 (100%)	483 (91%)	46 (9%)	12	6
All	All	1057/1054 (100%)	975 (92%)	82 (8%)	15	9

5 of 82 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	49	HIS
1	B	249	ILE
1	B	551	GLU
1	B	55	THR
1	B	196	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	512	ASN
1	B	78	HIS
1	B	479	GLN
1	A	498	ASN
1	B	488	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 21 ligands modelled in this entry, 3 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	667	-	5,5,5	0.36	0	5,5,5	0.54	0
2	GOL	A	668	-	5,5,5	0.36	0	5,5,5	0.49	0
3	FMT	A	669	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	670	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	671	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	672	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	673	-	0,2,2	0.00	-	0,1,1	0.00	-
4	PEG	A	674	-	6,6,6	0.53	0	5,5,5	1.61	1 (20%)
4	PEG	A	677	-	6,6,6	0.52	0	5,5,5	1.42	1 (20%)
4	PEG	A	678	-	6,6,6	0.52	0	5,5,5	1.82	2 (40%)
3	FMT	B	667	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	668	5	0,2,2	0.00	-	0,1,1	0.00	-
4	PEG	B	669	-	6,6,6	0.60	0	5,5,5	2.25	3 (60%)
7	SO4	B	671	-	4,4,4	0.18	0	6,6,6	0.19	0
3	FMT	B	672	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	673	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	674	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	675	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	667	-	-	0/4/4/4	0/0/0/0
2	GOL	A	668	-	-	0/4/4/4	0/0/0/0
3	FMT	A	669	-	-	0/0/0/0	0/0/0/0
3	FMT	A	670	-	-	0/0/0/0	0/0/0/0
3	FMT	A	671	-	-	0/0/0/0	0/0/0/0
3	FMT	A	672	-	-	0/0/0/0	0/0/0/0
3	FMT	A	673	-	-	0/0/0/0	0/0/0/0
4	PEG	A	674	-	-	0/4/4/4	0/0/0/0
4	PEG	A	677	-	-	0/4/4/4	0/0/0/0
4	PEG	A	678	-	-	0/4/4/4	0/0/0/0
3	FMT	B	667	-	-	0/0/0/0	0/0/0/0
3	FMT	B	668	5	-	0/0/0/0	0/0/0/0
4	PEG	B	669	-	-	0/4/4/4	0/0/0/0
7	SO4	B	671	-	-	0/0/0/0	0/0/0/0
3	FMT	B	672	-	-	0/0/0/0	0/0/0/0
3	FMT	B	673	-	-	0/0/0/0	0/0/0/0
3	FMT	B	674	-	-	0/0/0/0	0/0/0/0
3	FMT	B	675	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	674	PEG	O2-C2-C1	2.05	119.62	110.15
4	A	678	PEG	O2-C3-C4	2.08	119.77	110.15
4	A	678	PEG	O2-C2-C1	2.15	120.07	110.15
4	B	669	PEG	O2-C2-C1	2.16	120.14	110.15
4	A	677	PEG	O2-C3-C4	2.32	120.84	110.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	667	GOL	2	0
3	A	669	FMT	3	0
3	A	672	FMT	1	0
3	A	673	FMT	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	677	PEG	4	0
4	A	678	PEG	3	0
3	B	668	FMT	3	0
3	B	672	FMT	1	0
3	B	674	FMT	2	0
3	B	675	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	642/690 (93%)	-0.12	16 (2%) 58 65	17, 42, 77, 120	0
1	B	642/690 (93%)	0.22	31 (4%) 31 39	19, 53, 91, 126	0
All	All	1284/1380 (93%)	0.05	47 (3%) 42 48	17, 47, 87, 126	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	TYR	5.1
1	B	142	ASP	5.0
1	A	142	ASP	4.3
1	B	269	VAL	3.9
1	A	144	TYR	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	PEG	A	674	7/7	0.53	0.35	13.24	78,80,80,81	0
4	PEG	A	677	7/7	0.90	0.36	12.05	68,71,73,74	0
4	PEG	B	669	7/7	0.77	0.26	11.96	50,58,66,67	0
4	PEG	A	678	7/7	0.36	0.25	9.89	76,78,82,83	0
3	FMT	A	673	3/3	0.79	0.19	7.36	45,45,47,49	0
3	FMT	B	674	3/3	0.76	0.17	2.60	46,46,47,50	0
2	GOL	A	667	6/6	0.85	0.19	2.39	43,53,58,58	0
3	FMT	B	667	3/3	0.92	0.19	2.26	41,41,47,49	0
7	SO4	B	671	5/5	0.84	0.37	2.12	121,121,122,124	0
3	FMT	A	669	3/3	0.90	0.16	1.40	48,48,51,53	0
3	FMT	B	673	3/3	0.84	0.21	0.76	65,65,70,71	0
3	FMT	B	672	3/3	0.70	0.17	0.29	60,60,65,66	0
2	GOL	A	668	6/6	0.86	0.15	0.28	64,68,72,74	0
3	FMT	A	671	3/3	0.92	0.11	-0.41	53,53,57,59	0
3	FMT	B	668	3/3	0.96	0.12	-0.44	43,43,45,47	0
6	CL	A	676	1/1	0.97	0.09	-0.77	63,63,63,63	0
3	FMT	A	670	3/3	0.95	0.09	-0.78	44,44,45,47	0
5	MG	B	670	1/1	0.94	0.09	-0.83	42,42,42,42	0
3	FMT	B	675	3/3	0.87	0.12	-	65,65,66,67	0
3	FMT	A	672	3/3	0.89	0.16	-	54,54,58,61	0
5	MG	A	675	1/1	0.94	0.10	-	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.