



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 04:09 am GMT

PDB ID : 3HYV
Title : 3-D X-Ray structure of the sulfide:quinone oxidoreductase from the hyperthermophilic bacterium Aquifex aeolicus
Authors : Marcia, M.; Ermler, U.; Peng, G.H.; Michel, H.
Deposited on : 2009-06-23
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	430	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
1	B	430	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
1	C	430	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>
1	D	430	<div> <div>9%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>.</div> </div> </div>
1	E	430	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
1	F	430	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LMT	A	600	-	-	-	X
3	LMT	B	600	-	-	-	X
3	LMT	C	600	-	-	-	X
3	LMT	D	600	-	-	-	X
3	LMT	E	600	-	-	-	X
3	LMT	F	600	-	-	-	X
4	H2S	A	700	-	-	-	X
4	H2S	B	700	-	-	-	X
4	H2S	C	700	-	-	-	X
4	H2S	D	700	-	-	-	X
4	H2S	E	700	-	-	-	X
4	H2S	F	700	-	-	-	X
5	PS9	A	800	-	-	X	X
5	PS9	B	800	-	-	X	X
5	PS9	C	800	-	-	-	X
5	PS9	D	802	-	-	-	X
5	PS9	E	800	-	-	-	X
5	PS9	F	800	-	-	-	X
7	SO4	C	432	-	-	X	-

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 21480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfide-quinone reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	8	0
			3370	2182	556	609	23			
1	B	429	Total	C	N	O	S	0	7	0
			3364	2179	556	606	23			
1	C	429	Total	C	N	O	S	0	6	0
			3358	2174	555	606	23			
1	D	429	Total	C	N	O	S	0	6	0
			3357	2175	555	604	23			
1	E	429	Total	C	N	O	S	0	4	0
			3346	2166	555	602	23			
1	F	429	Total	C	N	O	S	0	4	0
			3346	2166	555	602	23			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: $C_{24}H_{46}O_{11}$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			35	24	11		
3	B	1	Total	C	O	0	0
			35	24	11		
3	C	1	Total	C	O	0	0
			35	24	11		
3	D	1	Total	C	O	0	0
			35	24	11		
3	E	1	Total	C	O	0	0
			35	24	11		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	C	O	0	0
			35	24	11		

- Molecule 4 is HYDROSULFURIC ACID (three-letter code: H2S) (formula: H₂S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	S	0	0
			1	1		
4	B	1	Total	S	0	0
			1	1		
4	C	1	Total	S	0	0
			1	1		
4	D	1	Total	S	0	0
			1	1		
4	E	1	Total	S	0	0
			1	1		
4	F	1	Total	S	0	0
			1	1		

- Molecule 5 is OCTATHIOCANANE (three-letter code: PS9) (formula: S₈).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total S 8 8	0	0
5	B	1	Total S 6 6	0	0
5	C	1	Total S 6 6	0	0
5	D	1	Total S 1 1	0	0
5	D	1	Total S 5 5	0	0
5	E	1	Total S 8 8	0	0
5	F	1	Total S 8 8	0	0

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	C	1	Total	O	S	0	0
			5	4	1		
7	C	1	Total	O	S	0	0
			5	4	1		
7	C	1	Total	O	S	0	0
			5	4	1		
7	C	1	Total	O	S	0	0
			5	4	1		
7	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	D	1	Total	O	S	0	0
			5	4	1		
7	D	1	Total	O	S	0	0
			5	4	1		
7	E	1	Total	O	S	0	0
			5	4	1		
7	E	1	Total	O	S	0	0
			5	4	1		
7	E	1	Total	O	S	0	0
			5	4	1		
7	E	1	Total	O	S	0	0
			5	4	1		
7	F	1	Total	O	S	0	0
			5	4	1		
7	F	1	Total	O	S	0	0
			5	4	1		
7	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	97	Total	O	0	0
			97	97		
8	B	107	Total	O	0	0
			107	107		
8	C	105	Total	O	0	0
			105	105		
8	D	57	Total	O	0	0
			57	57		
8	E	92	Total	O	0	0
			92	92		
8	F	113	Total	O	0	0
			113	113		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sulfide-quinone reductase




- Molecule 1: Sulfide-quinone reductase




- Molecule 1: Sulfide-quinone reductase





- Molecule 1: Sulfide-quinone reductase







• Molecule 1: Sulfide-quinone reductase



• Molecule 1: Sulfide-quinone reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	112.85Å 154.92Å 178.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.04 – 2.30 20.04 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.9 (20.04-2.30) 97.9 (20.04-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.235 0.214 , 0.256	Depositor DCC
R_{free} test set	6820 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	21480	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSS, LMT, H2S, PS9, SO4, MES, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3477	0.57	0/4720
1	B	0.50	0/3468	0.58	0/4707
1	C	0.50	0/3459	0.58	0/4696
1	D	0.47	0/3458	0.54	0/4695
1	E	0.48	0/3441	0.57	0/4672
1	F	0.50	0/3441	0.57	0/4672
All	All	0.49	0/20744	0.57	0/28162

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3370	0	3373	25	0
1	B	3364	0	3372	26	0
1	C	3358	0	3359	33	0
1	D	3357	0	3364	31	0
1	E	3346	0	3347	18	0
1	F	3346	0	3347	27	0
2	A	53	0	30	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	53	0	29	0	0
2	C	53	0	29	0	0
2	D	53	0	29	1	0
2	E	53	0	29	1	0
2	F	53	0	29	1	0
3	A	35	0	46	0	0
3	B	35	0	46	0	0
3	C	35	0	46	2	0
3	D	35	0	46	0	0
3	E	35	0	46	8	0
3	F	35	0	46	9	0
4	A	1	0	0	1	0
4	B	1	0	0	1	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	1	0
5	A	8	0	0	2	0
5	B	6	0	0	2	0
5	C	6	0	0	1	0
5	D	6	0	0	1	0
5	E	8	0	0	0	0
5	F	8	0	0	1	0
6	A	12	0	12	0	0
6	B	12	0	12	0	0
6	C	12	0	12	1	0
6	D	12	0	12	0	0
6	E	12	0	12	0	0
6	F	12	0	12	0	0
7	A	20	0	0	0	0
7	B	25	0	0	1	0
7	C	20	0	0	2	0
7	D	15	0	0	1	0
7	E	25	0	0	0	0
7	F	15	0	0	1	0
8	A	97	0	0	2	0
8	B	107	0	0	3	0
8	C	105	0	0	4	0
8	D	57	0	0	0	0
8	E	92	0	0	1	0
8	F	113	0	0	1	0
All	All	21480	0	20685	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:347[B]:CYS:SG	5:D:802:PS9:S9	2.39	1.19
1:C:420:LYS:H	1:C:420:LYS:HD3	1.28	0.98
3:E:600:LMT:H1B	3:F:600:LMT:O2B	1.64	0.97
1:D:327:ASN:HD21	1:D:339:TYR:H	1.17	0.92
3:F:600:LMT:O2B	3:F:600:LMT:H4'	1.72	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	433/430 (101%)	427 (99%)	6 (1%)	0	100	100
1	B	432/430 (100%)	426 (99%)	6 (1%)	0	100	100
1	C	431/430 (100%)	423 (98%)	8 (2%)	0	100	100
1	D	431/430 (100%)	421 (98%)	9 (2%)	1 (0%)	51	63
1	E	429/430 (100%)	417 (97%)	12 (3%)	0	100	100
1	F	429/430 (100%)	422 (98%)	7 (2%)	0	100	100
All	All	2585/2580 (100%)	2536 (98%)	48 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	334	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/356 (102%)	350 (97%)	12 (3%)	43	59
1	B	361/356 (101%)	350 (97%)	11 (3%)	46	63
1	C	360/356 (101%)	343 (95%)	17 (5%)	30	41
1	D	360/356 (101%)	349 (97%)	11 (3%)	45	61
1	E	358/356 (101%)	345 (96%)	13 (4%)	40	55
1	F	358/356 (101%)	348 (97%)	10 (3%)	49	65
All	All	2159/2136 (101%)	2085 (97%)	74 (3%)	42	57

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	338	LYS
1	D	42	THR
1	F	226	ILE
1	C	343	LEU
1	C	390	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	283	ASN
1	D	214	ASN
1	F	214	ASN
1	C	327	ASN
1	C	379	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSS	A	156[A]	-	5,5,7	1.15	0	2,5,8	1.74	1 (50%)
1	CSS	A	156[B]	5	5,6,7	0.83	0	3,6,8	1.46	1 (33%)
1	CSS	B	156[A]	-	5,5,7	1.26	0	2,5,8	1.54	1 (50%)
1	CSS	B	156[B]	5	5,6,7	1.06	0	3,6,8	2.17	2 (66%)
1	CSS	C	156[A]	-	5,5,7	1.40	0	2,5,8	1.60	0
1	CSS	C	156[B]	5	5,6,7	1.22	0	3,6,8	1.10	0
1	CSS	D	156[A]	-	5,5,7	1.33	1 (20%)	2,5,8	2.55	2 (100%)
1	CSS	D	156[B]	5	5,6,7	0.91	0	3,6,8	1.44	1 (33%)
1	CSS	E	156[A]	-	5,5,7	1.31	1 (20%)	2,5,8	1.60	1 (50%)
1	CSS	E	156[B]	5	5,6,7	0.94	0	3,6,8	1.36	1 (33%)
1	CSS	F	156[A]	-	5,5,7	1.22	0	2,5,8	1.54	1 (50%)
1	CSS	F	156[B]	5	5,6,7	0.91	0	3,6,8	1.75	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	A	156[B]	5	-	0/1/5/7	0/0/0/0
1	CSS	B	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	B	156[B]	5	-	0/1/5/7	0/0/0/0
1	CSS	C	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	C	156[B]	5	-	0/1/5/7	0/0/0/0
1	CSS	D	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	D	156[B]	5	-	0/1/5/7	0/0/0/0
1	CSS	E	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	E	156[B]	5	-	0/1/5/7	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	F	156[A]	-	-	0/1/4/7	0/0/0/0
1	CSS	F	156[B]	5	-	0/1/5/7	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	156[A]	CSS	CB-SG	-2.54	1.76	1.81
1	E	156[A]	CSS	CB-SG	-2.09	1.77	1.81

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	156[A]	CSS	CA-CB-SG	-2.84	108.02	114.42
1	A	156[A]	CSS	O-C-CA	-2.45	118.26	125.02
1	A	156[B]	CSS	O-C-CA	-2.45	118.26	125.02
1	D	156[B]	CSS	O-C-CA	-2.21	118.93	125.02
1	D	156[A]	CSS	O-C-CA	-2.21	118.93	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	156[A]	CSS	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 55 ligands modelled in this entry, 7 are modelled with single atom - leaving 48 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	SO4	A	431	-	4,4,4	0.16	0	6,6,6	0.11	0
7	SO4	A	432	-	4,4,4	0.14	0	6,6,6	0.16	0
7	SO4	A	433	-	4,4,4	0.13	0	6,6,6	0.24	0
7	SO4	A	434	-	4,4,4	0.15	0	6,6,6	0.09	0
2	FAD	A	441	4	51,58,58	1.00	1 (1%)	54,89,89	1.79	6 (11%)
3	LMT	A	600	-	36,36,36	0.53	0	47,47,47	0.98	2 (4%)
5	PS9	A	800	1	8,8,8	0.76	0	8,8,8	1.29	1 (12%)
6	MES	A	900	-	12,12,12	2.13	1 (8%)	14,16,16	2.69	7 (50%)
7	SO4	B	431	-	4,4,4	0.12	0	6,6,6	0.17	0
7	SO4	B	432	-	4,4,4	0.18	0	6,6,6	0.17	0
7	SO4	B	433	-	4,4,4	0.21	0	6,6,6	0.32	0
7	SO4	B	434	-	4,4,4	0.14	0	6,6,6	0.11	0
7	SO4	B	435	-	4,4,4	0.14	0	6,6,6	0.13	0
2	FAD	B	441	4	51,58,58	1.00	2 (3%)	54,89,89	1.80	7 (12%)
3	LMT	B	600	-	36,36,36	0.61	1 (2%)	47,47,47	1.32	5 (10%)
5	PS9	B	800	1	3,5,8	1.60	0	2,4,8	0.74	0
6	MES	B	900	-	12,12,12	2.12	1 (8%)	14,16,16	2.75	7 (50%)
7	SO4	C	431	-	4,4,4	0.12	0	6,6,6	0.15	0
7	SO4	C	432	-	4,4,4	0.10	0	6,6,6	0.18	0
7	SO4	C	433	-	4,4,4	0.19	0	6,6,6	0.34	0
7	SO4	C	434	-	4,4,4	0.18	0	6,6,6	0.21	0
2	FAD	C	441	4	51,58,58	1.00	1 (1%)	54,89,89	1.91	6 (11%)
3	LMT	C	600	-	36,36,36	0.52	0	47,47,47	1.10	4 (8%)
5	PS9	C	800	1	3,5,8	1.61	0	2,4,8	1.08	0
6	MES	C	900	-	12,12,12	2.17	1 (8%)	14,16,16	2.77	5 (35%)
7	SO4	D	432	-	4,4,4	0.16	0	6,6,6	0.17	0
7	SO4	D	433	-	4,4,4	0.14	0	6,6,6	0.11	0
7	SO4	D	434	-	4,4,4	0.15	0	6,6,6	0.20	0
2	FAD	D	441	4	51,58,58	1.04	3 (5%)	54,89,89	1.89	7 (12%)
3	LMT	D	600	-	36,36,36	0.48	0	47,47,47	0.92	1 (2%)
5	PS9	D	802	-	2,4,8	1.61	0	1,3,8	1.38	0
6	MES	D	900	-	12,12,12	2.21	1 (8%)	14,16,16	2.57	4 (28%)
7	SO4	E	431	-	4,4,4	0.19	0	6,6,6	0.15	0
7	SO4	E	432	-	4,4,4	0.14	0	6,6,6	0.12	0
7	SO4	E	433	-	4,4,4	0.15	0	6,6,6	0.14	0
7	SO4	E	434	-	4,4,4	0.17	0	6,6,6	0.15	0
7	SO4	E	435	-	4,4,4	0.13	0	6,6,6	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	E	441	4	51,58,58	1.10	3 (5%)	54,89,89	2.00	7 (12%)
3	LMT	E	600	-	36,36,36	0.53	0	47,47,47	1.03	4 (8%)
5	PS9	E	800	1	8,8,8	0.72	0	8,8,8	1.11	0
6	MES	E	900	-	12,12,12	2.15	1 (8%)	14,16,16	2.80	7 (50%)
7	SO4	F	431	-	4,4,4	0.15	0	6,6,6	0.09	0
7	SO4	F	433	-	4,4,4	0.17	0	6,6,6	0.27	0
7	SO4	F	434	-	4,4,4	0.15	0	6,6,6	0.11	0
2	FAD	F	441	4	51,58,58	1.00	2 (3%)	54,89,89	1.85	5 (9%)
3	LMT	F	600	-	36,36,36	0.49	0	47,47,47	1.02	1 (2%)
5	PS9	F	800	1	8,8,8	0.76	0	8,8,8	1.12	0
6	MES	F	900	-	12,12,12	2.10	1 (8%)	14,16,16	2.82	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SO4	A	431	-	-	0/0/0/0	0/0/0/0
7	SO4	A	432	-	-	0/0/0/0	0/0/0/0
7	SO4	A	433	-	-	0/0/0/0	0/0/0/0
7	SO4	A	434	-	-	0/0/0/0	0/0/0/0
2	FAD	A	441	4	-	0/28/50/50	0/6/6/6
3	LMT	A	600	-	-	0/21/61/61	0/2/2/2
5	PS9	A	800	1	-	0/0/8/8	0/1/1/1
6	MES	A	900	-	-	0/6/14/14	0/1/1/1
7	SO4	B	431	-	-	0/0/0/0	0/0/0/0
7	SO4	B	432	-	-	0/0/0/0	0/0/0/0
7	SO4	B	433	-	-	0/0/0/0	0/0/0/0
7	SO4	B	434	-	-	0/0/0/0	0/0/0/0
7	SO4	B	435	-	-	0/0/0/0	0/0/0/0
2	FAD	B	441	4	-	0/28/50/50	0/6/6/6
3	LMT	B	600	-	-	0/21/61/61	0/2/2/2
5	PS9	B	800	1	-	0/3/3/8	0/0/0/1
6	MES	B	900	-	-	0/6/14/14	0/1/1/1
7	SO4	C	431	-	-	0/0/0/0	0/0/0/0
7	SO4	C	432	-	-	0/0/0/0	0/0/0/0
7	SO4	C	433	-	-	0/0/0/0	0/0/0/0
7	SO4	C	434	-	-	0/0/0/0	0/0/0/0
2	FAD	C	441	4	-	0/28/50/50	0/6/6/6
3	LMT	C	600	-	-	0/21/61/61	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PS9	C	800	1	-	0/3/3/8	0/0/0/1
6	MES	C	900	-	-	0/6/14/14	0/1/1/1
7	SO4	D	432	-	-	0/0/0/0	0/0/0/0
7	SO4	D	433	-	-	0/0/0/0	0/0/0/0
7	SO4	D	434	-	-	0/0/0/0	0/0/0/0
2	FAD	D	441	4	-	0/28/50/50	0/6/6/6
3	LMT	D	600	-	-	0/21/61/61	0/2/2/2
5	PS9	D	802	-	-	0/2/2/8	0/0/0/1
6	MES	D	900	-	-	0/6/14/14	0/1/1/1
7	SO4	E	431	-	-	0/0/0/0	0/0/0/0
7	SO4	E	432	-	-	0/0/0/0	0/0/0/0
7	SO4	E	433	-	-	0/0/0/0	0/0/0/0
7	SO4	E	434	-	-	0/0/0/0	0/0/0/0
7	SO4	E	435	-	-	0/0/0/0	0/0/0/0
2	FAD	E	441	4	-	0/28/50/50	0/6/6/6
3	LMT	E	600	-	-	0/21/61/61	0/2/2/2
5	PS9	E	800	1	-	0/0/8/8	0/1/1/1
6	MES	E	900	-	-	0/6/14/14	0/1/1/1
7	SO4	F	431	-	-	0/0/0/0	0/0/0/0
7	SO4	F	433	-	-	0/0/0/0	0/0/0/0
7	SO4	F	434	-	-	0/0/0/0	0/0/0/0
2	FAD	F	441	4	-	0/28/50/50	0/6/6/6
3	LMT	F	600	-	-	0/21/61/61	0/2/2/2
5	PS9	F	800	1	-	0/0/8/8	0/1/1/1
6	MES	F	900	-	-	0/6/14/14	0/1/1/1

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	900	MES	C8-S	-7.42	1.66	1.77
6	C	900	MES	C8-S	-7.21	1.66	1.77
6	E	900	MES	C8-S	-7.18	1.66	1.77
6	A	900	MES	C8-S	-7.07	1.66	1.77
6	B	900	MES	C8-S	-7.03	1.67	1.77

The worst 5 of 93 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	441	FAD	N3A-C2A-N1A	-9.79	120.33	128.86
2	F	441	FAD	N3A-C2A-N1A	-9.16	120.88	128.86
2	A	441	FAD	N3A-C2A-N1A	-9.09	120.94	128.86
2	C	441	FAD	N3A-C2A-N1A	-9.03	121.00	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	441	FAD	N3A-C2A-N1A	-8.90	121.11	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

17 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	441	FAD	1	0
5	A	800	PS9	2	0
7	B	435	SO4	1	0
5	B	800	PS9	2	0
7	C	432	SO4	2	0
3	C	600	LMT	2	0
5	C	800	PS9	1	0
6	C	900	MES	1	0
7	D	432	SO4	1	0
2	D	441	FAD	1	0
5	D	802	PS9	1	0
2	E	441	FAD	1	0
3	E	600	LMT	8	0
7	F	431	SO4	1	0
2	F	441	FAD	1	0
3	F	600	LMT	9	0
5	F	800	PS9	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/430 (99%)	0.13	10 (2%) 61 67	31, 38, 45, 51	0
1	B	428/430 (99%)	0.12	12 (2%) 53 61	31, 38, 45, 54	0
1	C	428/430 (99%)	0.11	7 (1%) 72 77	30, 38, 44, 49	0
1	D	428/430 (99%)	0.36	39 (9%) 10 14	30, 38, 44, 47	0
1	E	428/430 (99%)	0.10	15 (3%) 44 51	31, 38, 45, 47	0
1	F	428/430 (99%)	0.09	12 (2%) 53 61	32, 38, 45, 54	0
All	All	2568/2580 (99%)	0.15	95 (3%) 42 49	30, 38, 44, 54	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	430	CYS	6.5
1	A	2	ALA	6.5
1	C	430	CYS	6.2
1	A	430	CYS	6.1
1	B	430	CYS	4.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSS	D	156[B]	7/8	0.97	0.09	-	34,35,35,35	2
1	CSS	C	156[B]	7/8	0.93	0.12	-	38,39,40,40	2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CSS	C	156[A]	6/8	0.93	0.12	-	39,39,40,43	1
1	CSS	A	156[A]	6/8	0.95	0.18	-	37,38,39,42	1
1	CSS	E	156[A]	6/8	0.95	0.14	-	37,38,38,40	1
1	CSS	B	156[B]	7/8	0.93	0.14	-	35,36,36,36	2
1	CSS	A	156[B]	7/8	0.95	0.18	-	35,38,38,39	2
1	CSS	F	156[B]	7/8	0.95	0.15	-	35,37,37,37	2
1	CSS	E	156[B]	7/8	0.95	0.14	-	37,38,38,38	2
1	CSS	B	156[A]	6/8	0.93	0.14	-	36,36,36,36	1
1	CSS	F	156[A]	6/8	0.95	0.15	-	37,37,37,38	1
1	CSS	D	156[A]	6/8	0.97	0.09	-	35,35,35,37	1

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	PS9	D	802	5/8	0.47	0.96	32.42	48,48,48,49	5
4	H2S	F	700	1/1	0.95	0.29	30.66	30,30,30,30	1
5	PS9	C	800	6/8	0.47	0.43	24.58	37,38,38,39	6
5	PS9	E	800	8/8	0.62	0.46	17.10	39,39,40,40	8
5	PS9	B	800	6/8	0.66	0.51	17.00	36,37,37,37	6
5	PS9	F	800	8/8	0.61	0.47	15.61	35,35,36,37	8
4	H2S	A	700	1/1	0.96	0.35	11.54	31,31,31,31	1
5	PS9	A	800	8/8	0.66	0.45	11.39	34,35,35,35	8
3	LMT	F	600	35/35	0.41	0.45	9.88	36,45,52,53	35
4	H2S	E	700	1/1	0.94	0.26	9.73	32,32,32,32	1
4	H2S	C	700	1/1	0.98	0.33	9.64	34,34,34,34	1
3	LMT	C	600	35/35	0.55	0.40	8.99	34,46,51,52	35
3	LMT	A	600	35/35	0.30	0.48	8.65	40,44,49,50	35
4	H2S	B	700	1/1	0.94	0.21	7.30	32,32,32,32	1
3	LMT	D	600	35/35	0.58	0.37	6.79	43,49,50,51	35

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	LMT	B	600	35/35	0.50	0.38	6.54	41,47,51,52	35
3	LMT	E	600	35/35	0.61	0.40	6.52	40,43,47,47	35
4	H2S	D	700	1/1	0.93	0.16	2.28	37,37,37,37	1
6	MES	D	900	12/12	0.91	0.22	1.58	48,53,57,58	0
6	MES	A	900	12/12	0.92	0.20	0.78	56,60,63,63	0
6	MES	C	900	12/12	0.92	0.19	0.61	50,57,61,62	0
7	SO4	B	433	5/5	0.86	0.16	0.06	65,66,66,67	0
6	MES	E	900	12/12	0.96	0.14	-0.42	55,59,63,63	0
6	MES	F	900	12/12	0.96	0.15	-0.50	47,51,55,55	0
2	FAD	D	441	53/53	0.95	0.11	-0.66	31,36,38,40	0
7	SO4	F	433	5/5	0.93	0.15	-0.67	62,63,63,64	0
2	FAD	E	441	53/53	0.95	0.10	-0.95	26,29,42,43	0
7	SO4	C	433	5/5	0.94	0.11	-1.19	60,61,61,62	0
7	SO4	E	433	5/5	0.95	0.09	-1.24	72,72,73,73	0
2	FAD	B	441	53/53	0.97	0.09	-1.61	23,30,30,32	0
2	FAD	A	441	53/53	0.96	0.10	-1.61	23,28,30,32	0
7	SO4	A	433	5/5	0.98	0.07	-1.71	42,42,43,43	0
6	MES	B	900	12/12	0.97	0.10	-1.74	47,52,55,55	0
2	FAD	F	441	53/53	0.97	0.09	-1.76	25,30,34,35	0
7	SO4	D	433	5/5	0.91	0.12	-1.77	101,101,101,101	0
2	FAD	C	441	53/53	0.97	0.08	-1.78	22,30,32,32	0
7	SO4	B	435	5/5	0.95	0.15	-	70,70,70,71	0
7	SO4	C	434	5/5	0.92	0.21	-	62,62,63,63	0
7	SO4	E	431	5/5	0.86	0.22	-	69,69,70,70	0
7	SO4	F	431	5/5	0.93	0.16	-	72,72,73,73	0
7	SO4	B	431	5/5	0.96	0.12	-	77,77,78,78	0
7	SO4	C	431	5/5	0.95	0.20	-	71,71,71,71	0
7	SO4	E	435	5/5	0.94	0.14	-	75,75,76,76	0
7	SO4	B	434	5/5	0.95	0.33	-	64,64,64,65	0
7	SO4	B	432	5/5	0.96	0.17	-	54,54,55,55	0
7	SO4	A	434	5/5	0.96	0.19	-	61,62,62,63	0
5	PS9	D	800	1/8	0.71	0.51	-	33,33,33,33	1
7	SO4	D	434	5/5	0.90	0.28	-	69,69,70,70	0
7	SO4	E	432	5/5	0.94	0.32	-	66,66,67,67	0
7	SO4	F	434	5/5	0.95	0.26	-	63,64,64,65	0
7	SO4	A	432	5/5	0.90	0.21	-	61,61,62,62	0
7	SO4	E	434	5/5	0.91	0.20	-	68,68,68,69	0
7	SO4	A	431	5/5	0.89	0.24	-	75,76,76,76	0
7	SO4	C	432	5/5	0.93	0.17	-	66,66,66,66	0
7	SO4	D	432	5/5	0.88	0.17	-	71,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.