



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 9, 2017 – 11:10 am GMT

PDB ID : 5HY8  
Title : Glycation restrains allosteric transition in hemoglobin: The molecular basis of oxidative stress under hyperglycemic conditions in diabetes  
Authors : Saraswathi, N.T.; Pannu, N.S.; Syakhovich, V.E.; Saurabh, A.; Bokut, S.B.; Moras, D.; Ruff, M.  
Deposited on : 2016-02-01  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	recalc28906
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28906


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div>10%</div> <div>96%</div> <div>.</div> </div>
1	C	141	<div> <div>13%</div> <div>98%</div> <div>.</div> </div>
1	E	141	<div> <div>81%</div> <div>79%</div> <div>13%</div> <div>.</div> <div>.</div> </div>
1	G	141	<div> <div>38%</div> <div>87%</div> <div>9%</div> <div>.</div> <div>.</div> </div>
1	S	141	<div> <div>48%</div> <div>96%</div> <div>.</div> </div>
2	B	146	<div> <div>8%</div> <div>96%</div> <div>.</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	D	146	
2	F	146	
2	H	146	
2	T	146	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OXY	A	202	-	-	-	X
4	OXY	B	202	-	-	-	X
4	OXY	G	202	-	-	-	X
5	FRU	A	203	-	-	-	X
5	FRU	C	203	-	-	-	X
5	FRU	F	203	-	-	-	X
6	GLC	C	204	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1068	685	187	193	3			
1	C	141	Total	C	N	O	S	0	0	0
			1068	685	187	193	3			
1	E	137	Total	C	N	O	S	0	0	0
			1030	659	180	188	3			
1	G	138	Total	C	N	O	S	0	0	0
			1042	668	181	190	3			
1	S	141	Total	C	N	O	S	0	0	0
			1068	685	187	193	3			

- Molecule 2 is a protein called Hemoglobin subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1122	724	195	200	3			
2	D	146	Total	C	N	O	S	0	0	0
			1122	724	195	200	3			
2	F	146	Total	C	N	O	S	0	0	0
			1122	724	195	200	3			
2	H	146	Total	C	N	O	S	0	0	0
			1122	724	195	200	3			
2	T	146	Total	C	N	O	S	0	0	0
			1122	724	195	200	3			

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	G	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	H	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	S	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	T	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is OXYGEN MOLECULE (three-letter code: OXY) (formula:  $O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 2 2	0	0
4	B	1	Total O 2 2	0	0
4	C	1	Total O 2 2	0	0
4	D	1	Total O 2 2	0	0
4	F	1	Total O 2 2	0	0
4	G	1	Total O 2 2	0	0
4	S	1	Total O 2 2	0	0
4	T	1	Total O 2 2	0	0

- Molecule 5 is FRUCTOSE (three-letter code: FRU) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	C	1	Total	C	O	0	0
			11	6	5		
5	F	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			12	6	6		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	30	Total 30	O 30	0	0
7	B	36	Total 36	O 36	0	0
7	C	39	Total 39	O 39	0	0
7	D	32	Total 32	O 32	0	0
7	E	12	Total 12	O 12	0	0
7	F	26	Total 26	O 26	0	0
7	G	18	Total 18	O 18	0	0
7	H	10	Total 10	O 10	0	0
7	S	15	Total 15	O 15	0	0
7	T	14	Total 14	O 14	0	0







- Molecule 1: Hemoglobin subunit alpha




- Molecule 2: Hemoglobin subunit beta




- Molecule 2: Hemoglobin subunit beta




- Molecule 2: Hemoglobin subunit beta



- Molecule 2: Hemoglobin subunit beta



- Molecule 2: Hemoglobin subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	237.99Å 59.27Å 137.02Å 90.00° 125.36° 90.00°	Depositor
Resolution (Å)	112.13 – 2.30 14.89 – 2.30	Depositor EDS
% Data completeness (in resolution range)	87.0 (112.13-2.30) 87.3 (14.89-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.203 , 0.244 0.265 , 0.293	Depositor DCC
$R_{free}$ test set	3017 reflections (4.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.0	Xtriage
Anisotropy	0.351	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h-2*k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	11609	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	109.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, GLC, OXY, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.01	1/1096 (0.1%)	0.91	1/1491 (0.1%)
1	C	0.95	0/1096	0.87	0/1491
1	E	0.77	1/1057 (0.1%)	1.13	14/1438 (1.0%)
1	G	0.90	0/1070	0.99	4/1456 (0.3%)
1	S	0.78	0/1096	0.84	1/1491 (0.1%)
2	B	1.10	2/1152 (0.2%)	0.89	1/1566 (0.1%)
2	D	1.03	1/1152 (0.1%)	0.87	2/1566 (0.1%)
2	F	0.99	2/1152 (0.2%)	1.00	9/1566 (0.6%)
2	H	0.81	0/1152	0.86	3/1566 (0.2%)
2	T	0.84	1/1152 (0.1%)	0.81	1/1566 (0.1%)
All	All	0.93	8/11175 (0.1%)	0.92	36/15197 (0.2%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	37	TRP	CE3-CZ3	7.08	1.50	1.38
1	A	49	SER	CA-CB	6.00	1.61	1.52
2	D	130	TYR	CE1-CZ	-5.67	1.31	1.38
1	E	116	GLU	CD-OE1	-5.59	1.19	1.25
2	B	101	GLU	CG-CD	5.36	1.59	1.51

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	80	LEU	CB-CG-CD2	10.33	128.56	111.00
1	E	129	LEU	CB-CG-CD2	9.12	126.50	111.00
2	F	68	LEU	CB-CG-CD2	8.13	124.83	111.00
1	E	92	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	E	92	ARG	NE-CZ-NH2	-7.82	116.39	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1068	0	1071	2	0
1	C	1068	0	1070	1	0
1	E	1030	0	1028	24	0
1	G	1042	0	1037	7	0
1	S	1068	0	1073	2	0
2	B	1122	0	1118	2	0
2	D	1122	0	1118	3	0
2	F	1122	0	1116	17	0
2	H	1122	0	1118	16	0
2	T	1122	0	1118	2	0
3	A	43	0	30	0	0
3	B	43	0	30	1	0
3	C	43	0	30	1	0
3	D	43	0	30	1	0
3	E	43	0	30	3	0
3	F	43	0	30	2	0
3	G	43	0	30	0	0
3	H	43	0	30	1	0
3	S	43	0	30	1	0
3	T	43	0	30	2	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
4	F	2	0	0	0	0
4	G	2	0	0	0	0
4	S	2	0	0	0	0
4	T	2	0	0	0	0
5	A	11	0	9	1	0
5	C	11	0	9	1	0
5	F	11	0	9	2	0
6	C	12	0	12	0	0
7	A	30	0	0	0	0
7	B	36	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	39	0	0	0	0
7	D	32	0	0	1	0
7	E	12	0	0	0	0
7	F	26	0	0	1	0
7	G	18	0	0	0	0
7	H	10	0	0	0	0
7	S	15	0	0	0	0
7	T	14	0	0	0	0
All	All	11609	0	11206	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:93:CYS:SG	2:H:145:TYR:HE1	1.78	1.05
2:H:93:CYS:SG	2:H:145:TYR:CE1	2.49	1.04
2:H:35:TYR:O	2:H:38:THR:HG22	1.64	0.96
1:E:32:MET:HE1	1:E:101:LEU:HB2	1.50	0.92
2:F:24:GLY:N	2:F:68:LEU:HD12	1.97	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	138 (99%)	1 (1%)	0	100	100
1	C	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
1	E	135/141 (96%)	133 (98%)	2 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	136/141 (96%)	134 (98%)	2 (2%)	0	100	100
1	S	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
2	B	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
2	D	144/146 (99%)	140 (97%)	4 (3%)	0	100	100
2	F	144/146 (99%)	137 (95%)	6 (4%)	1 (1%)	25	30
2	H	144/146 (99%)	140 (97%)	4 (3%)	0	100	100
2	T	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
All	All	1408/1435 (98%)	1379 (98%)	28 (2%)	1 (0%)	55	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	21	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	112 (99%)	1 (1%)	82	91
1	C	113/113 (100%)	112 (99%)	1 (1%)	82	91
1	E	109/113 (96%)	106 (97%)	3 (3%)	49	65
1	G	110/113 (97%)	106 (96%)	4 (4%)	40	55
1	S	113/113 (100%)	111 (98%)	2 (2%)	64	79
2	B	118/118 (100%)	117 (99%)	1 (1%)	85	93
2	D	118/118 (100%)	116 (98%)	2 (2%)	66	81
2	F	118/118 (100%)	116 (98%)	2 (2%)	66	81
2	H	118/118 (100%)	115 (98%)	3 (2%)	53	70
2	T	118/118 (100%)	117 (99%)	1 (1%)	85	93
All	All	1148/1155 (99%)	1128 (98%)	20 (2%)	66	81



5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	117	HIS
1	G	52	SER
2	H	120	LYS
1	E	75	ASP
2	F	87	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	87	HIS
2	H	63	HIS
2	H	143	HIS
2	H	146	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HEM	A	201	1,4	28,50,50	1.08	2 (7%)	17,82,82	1.62	4 (23%)
4	OXY	A	202	3	1,1,1	0.22	0	0,0,0	0.00	-
5	FRU	A	203	1	11,11,12	0.77	0	12,17,18	1.05	0
3	HEM	B	201	2,4	28,50,50	1.24	3 (10%)	17,82,82	1.55	3 (17%)
4	OXY	B	202	3	1,1,1	0.41	0	0,0,0	0.00	-
3	HEM	C	201	1,4	28,50,50	1.08	2 (7%)	17,82,82	1.79	3 (17%)
4	OXY	C	202	3	1,1,1	0.21	0	0,0,0	0.00	-
5	FRU	C	203	1	11,11,12	0.85	1 (9%)	12,17,18	1.39	2 (16%)
6	GLC	C	204	-	12,12,12	1.80	3 (25%)	17,17,17	1.90	7 (41%)
3	HEM	D	201	2,4	28,50,50	0.94	1 (3%)	17,82,82	1.62	4 (23%)
4	OXY	D	202	3	1,1,1	0.33	0	0,0,0	0.00	-
3	HEM	E	201	1	28,50,50	1.53	4 (14%)	17,82,82	1.65	4 (23%)
3	HEM	F	201	2,4	28,50,50	1.12	1 (3%)	17,82,82	1.85	5 (29%)
4	OXY	F	202	3	1,1,1	0.29	0	0,0,0	0.00	-
5	FRU	F	203	2	11,11,12	0.71	0	12,17,18	1.35	3 (25%)
3	HEM	G	201	1,4	28,50,50	0.71	0	17,82,82	1.68	2 (11%)
4	OXY	G	202	3	1,1,1	0.26	0	0,0,0	0.00	-
3	HEM	H	201	2	28,50,50	0.94	2 (7%)	17,82,82	1.38	2 (11%)
3	HEM	S	201	1,4	28,50,50	1.19	1 (3%)	17,82,82	1.56	4 (23%)
4	OXY	S	202	3	1,1,1	0.23	0	0,0,0	0.00	-
3	HEM	T	201	2,4	28,50,50	0.78	1 (3%)	17,82,82	1.25	2 (11%)
4	OXY	T	202	3	1,1,1	0.33	0	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	201	1,4	-	0/6/54/54	0/0/8/8
4	OXY	A	202	3	-	0/0/0/0	0/0/0/0
5	FRU	A	203	1	-	0/2/21/24	0/1/1/1
3	HEM	B	201	2,4	-	0/6/54/54	0/0/8/8
4	OXY	B	202	3	-	0/0/0/0	0/0/0/0
3	HEM	C	201	1,4	-	0/6/54/54	0/0/8/8
4	OXY	C	202	3	-	0/0/0/0	0/0/0/0
5	FRU	C	203	1	-	0/2/21/24	0/1/1/1
6	GLC	C	204	-	-	0/2/22/22	0/1/1/1
3	HEM	D	201	2,4	-	0/6/54/54	0/0/8/8
4	OXY	D	202	3	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	E	201	1	-	0/6/54/54	0/0/8/8
3	HEM	F	201	2,4	-	0/6/54/54	0/0/8/8
4	OXY	F	202	3	-	0/0/0/0	0/0/0/0
5	FRU	F	203	2	-	0/2/21/24	0/1/1/1
3	HEM	G	201	1,4	-	0/6/54/54	0/0/8/8
4	OXY	G	202	3	-	0/0/0/0	0/0/0/0
3	HEM	H	201	2	-	0/6/54/54	0/0/8/8
3	HEM	S	201	1,4	-	0/6/54/54	0/0/8/8
4	OXY	S	202	3	-	0/0/0/0	0/0/0/0
3	HEM	T	201	2,4	-	0/6/54/54	0/0/8/8
4	OXY	T	202	3	-	0/0/0/0	0/0/0/0

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	201	HEM	C1B-NB	-5.32	1.30	1.36
3	S	201	HEM	C1B-NB	-3.84	1.32	1.36
3	B	201	HEM	C3B-C2B	-3.50	1.35	1.40
3	B	201	HEM	C1B-NB	-3.22	1.33	1.36
3	A	201	HEM	C1B-NB	-3.22	1.33	1.36

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	201	HEM	CAA-CBA-CGA	-4.69	104.64	112.66
3	D	201	HEM	CBD-CAD-C3D	-4.51	103.87	112.47
3	A	201	HEM	CBD-CAD-C3D	-3.88	105.07	112.47
3	H	201	HEM	CAA-CBA-CGA	-3.55	106.59	112.66
3	F	201	HEM	CBD-CAD-C3D	-3.54	105.72	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	203	FRU	1	0
3	B	201	HEM	1	0
3	C	201	HEM	1	0
5	C	203	FRU	1	0
3	D	201	HEM	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	201	HEM	3	0
3	F	201	HEM	2	0
5	F	203	FRU	2	0
3	H	201	HEM	1	0
3	S	201	HEM	1	0
3	T	201	HEM	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	141/141 (100%)	0.60	14 (9%) 8 11	39, 70, 99, 119	0
1	C	141/141 (100%)	0.92	19 (13%) 3 5	36, 77, 119, 139	0
1	E	137/141 (97%)	4.22	114 (83%) 0 0	89, 209, 324, 372	0
1	G	138/141 (97%)	1.93	54 (39%) 0 0	47, 109, 194, 234	0
1	S	141/141 (100%)	2.12	68 (48%) 0 0	61, 116, 178, 193	0
2	B	146/146 (100%)	0.42	12 (8%) 12 17	28, 61, 109, 132	0
2	D	146/146 (100%)	1.00	22 (15%) 3 4	40, 80, 116, 137	0
2	F	146/146 (100%)	1.23	37 (25%) 1 1	35, 90, 185, 232	0
2	H	146/146 (100%)	3.03	94 (64%) 0 0	65, 155, 240, 273	0
2	T	146/146 (100%)	2.51	75 (51%) 0 0	69, 150, 211, 241	0
All	All	1428/1435 (99%)	1.79	509 (35%) 0 0	28, 98, 232, 372	0

The worst 5 of 509 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	83	LEU	12.9
1	E	80	LEU	12.1
1	E	136	LEU	11.5
2	H	48	LEU	11.2
2	T	141	LEU	10.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	GLC	C	204	12/12	0.81	0.37	5.87	47,56,60,62	0
4	OXY	B	202	2/2	0.98	0.25	5.74	35,35,35,56	0
4	OXY	A	202	2/2	0.96	0.35	3.69	46,46,46,72	0
5	FRU	C	203	11/12	0.88	0.33	2.66	79,83,95,115	0
4	OXY	G	202	2/2	0.84	0.50	2.60	80,80,80,117	0
5	FRU	A	203	11/12	0.81	0.33	2.17	77,89,100,105	0
5	FRU	F	203	11/12	0.66	0.46	1.48	99,107,119,123	0
4	OXY	S	202	2/2	0.93	0.35	0.67	63,63,63,88	0
3	HEM	B	201	43/43	0.94	0.15	0.09	23,28,44,55	0
3	HEM	T	201	43/43	0.84	0.32	0.06	71,98,114,124	0
4	OXY	D	202	2/2	0.97	0.20	0.06	44,44,44,76	0
3	HEM	D	201	43/43	0.91	0.18	-0.17	36,47,66,70	0
3	HEM	A	201	43/43	0.90	0.17	-0.21	30,40,53,59	0
4	OXY	C	202	2/2	0.98	0.19	-0.31	45,45,45,71	0
3	HEM	F	201	43/43	0.94	0.14	-0.48	28,35,60,71	0
3	HEM	G	201	43/43	0.91	0.18	-0.53	33,48,79,84	0
3	HEM	C	201	43/43	0.93	0.14	-0.63	23,39,82,100	0
3	HEM	H	201	43/43	0.82	0.25	-1.10	52,71,99,110	0
3	HEM	S	201	43/43	0.91	0.16	-1.17	33,45,66,70	0
3	HEM	E	201	43/43	0.86	0.20	-1.57	52,64,87,92	0
4	OXY	T	202	2/2	0.97	0.25	-1.88	56,56,56,84	0
4	OXY	F	202	2/2	0.98	0.18	-	78,78,78,81	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.