



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 27, 2017 – 06:09 PM EST

PDB ID : 5HYI  
Title : Glycosylated, disulfide-linked Hole-Hole Fc fragment  
Authors : Kuglstatter, A.; Stihle, M.; Benz, J.  
Deposited on : unknown  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	
1	B	227	
1	C	227	
1	D	227	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7251 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	0	0
			1656	1050	281	318	7			
1	B	209	Total	C	N	O	S	0	0	0
			1653	1047	281	318	7			
1	C	208	Total	C	N	O	S	0	0	0
			1649	1045	280	317	7			
1	D	209	Total	C	N	O	S	0	0	0
			1653	1047	281	318	7			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	349	CYS	TYR	engineered mutation	UNP P01857
A	366	SER	THR	engineered mutation	UNP P01857
A	368	ALA	LEU	engineered mutation	UNP P01857
A	407	VAL	TYR	engineered mutation	UNP P01857
B	349	CYS	TYR	engineered mutation	UNP P01857
B	366	SER	THR	engineered mutation	UNP P01857
B	368	ALA	LEU	engineered mutation	UNP P01857
B	407	VAL	TYR	engineered mutation	UNP P01857
C	349	CYS	TYR	engineered mutation	UNP P01857
C	366	SER	THR	engineered mutation	UNP P01857
C	368	ALA	LEU	engineered mutation	UNP P01857
C	407	VAL	TYR	engineered mutation	UNP P01857
D	349	CYS	TYR	engineered mutation	UNP P01857
D	366	SER	THR	engineered mutation	UNP P01857
D	368	ALA	LEU	engineered mutation	UNP P01857
D	407	VAL	TYR	engineered mutation	UNP P01857

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	C	1	Total	C	O	0	0
			11	6	5		
3	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		
4	C	1	Total	C	O	0	0
			11	6	5		
4	C	1	Total	C	O	0	0
			11	6	5		
4	D	1	Total	C	O	0	0
			11	6	5		
4	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	C	1	Total	C	O	0	0
			11	6	5		
5	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula:  $C_6H_{12}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		
6	B	1	Total	C	O	0	0
			10	6	4		
6	C	1	Total	C	O	0	0
			10	6	4		
6	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	66	Total	O	0	0
			66	66		
7	B	29	Total	O	0	0
			29	29		
7	C	33	Total	O	0	0
			33	33		
7	D	72	Total	O	0	0
			72	72		







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.28Å 105.66Å 134.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.15 – 2.90 49.17 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.15-2.90) 99.5 (49.17-2.90)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.29 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, $R_{free}$	0.175 , 0.235 0.197 , 0.254	Depositor DCC
$R_{free}$ test set	1677 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	75.9	Xtriage
Anisotropy	0.604	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 67.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.027 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7251	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, GAL, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/1701	0.79	0/2318
1	B	0.52	0/1697	0.75	0/2311
1	C	0.53	0/1693	0.72	0/2306
1	D	0.56	0/1697	0.78	0/2311
All	All	0.55	0/6788	0.76	0/9246

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1656	0	1624	14	0
1	B	1653	0	1620	25	0
1	C	1649	0	1617	24	0
1	D	1653	0	1620	21	0
2	A	56	0	48	1	0
2	B	56	0	48	0	0
2	C	56	0	48	0	0
2	D	56	0	48	1	0
3	A	11	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	11	0	8	0	0
3	C	11	0	8	0	0
3	D	11	0	8	1	0
4	A	22	0	18	0	0
4	B	22	0	18	0	0
4	C	22	0	18	0	0
4	D	22	0	18	1	0
5	A	11	0	10	0	0
5	B	11	0	10	0	0
5	C	11	0	10	0	0
5	D	11	0	10	0	0
6	A	10	0	10	0	0
6	B	10	0	10	0	0
6	C	10	0	10	1	0
6	D	10	0	10	0	0
7	A	66	0	0	0	0
7	B	29	0	0	1	0
7	C	33	0	0	1	0
7	D	72	0	0	1	0
All	All	7251	0	6857	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:422:VAL:HG22	1:B:442:SER:HB3	1.63	0.79
1:B:386:GLN:HG2	1:D:355:ARG:HG2	1.70	0.74
1:D:259:VAL:HG13	1:D:336:ILE:HD11	1.75	0.69
1:C:278:TYR:HB2	1:C:320:LYS:HB2	1.76	0.67
1:A:278:TYR:HB2	1:A:320:LYS:HB3	1.79	0.64

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/227 (91%)	197 (95%)	9 (4%)	1 (0%)	32	68
1	B	207/227 (91%)	190 (92%)	11 (5%)	6 (3%)	5	21
1	C	206/227 (91%)	186 (90%)	18 (9%)	2 (1%)	18	51
1	D	207/227 (91%)	197 (95%)	8 (4%)	2 (1%)	18	51
All	All	827/908 (91%)	770 (93%)	46 (6%)	11 (1%)	14	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	HIS
1	B	376	ASP
1	B	254	SER
1	B	310	HIS
1	D	310	HIS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/209 (93%)	176 (91%)	18 (9%)	10	31
1	B	193/209 (92%)	175 (91%)	18 (9%)	10	31
1	C	193/209 (92%)	170 (88%)	23 (12%)	6	18
1	D	193/209 (92%)	172 (89%)	21 (11%)	7	22
All	All	773/836 (92%)	693 (90%)	80 (10%)	8	25

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	250	THR
1	C	282	VAL
1	D	370	LYS
1	C	252	MET
1	C	264	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	276	ASN
1	D	435	HIS
1	D	429	HIS
1	C	347	GLN
1	D	311	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	501	1,2,6	14,14,15	0.26	0	15,19,21	0.73	1 (6%)
2	NAG	A	502	3,2	14,14,15	0.32	0	15,19,21	0.97	1 (6%)
3	BMA	A	503	2,4	11,11,12	0.44	0	13,15,17	0.76	0
4	MAN	A	504	3,2	11,11,12	0.36	0	13,15,17	0.91	2 (15%)
2	NAG	A	505	5,4	14,14,15	0.24	0	15,19,21	1.03	1 (6%)
4	MAN	A	506	3,2	11,11,12	0.73	1 (9%)	13,15,17	1.89	2 (15%)
5	GAL	A	507	2	11,11,12	0.46	0	13,15,17	0.75	0
6	FUC	A	508	2	9,10,11	0.48	0	13,14,16	0.93	1 (7%)
2	NAG	A	509	4	14,14,15	0.38	0	15,19,21	1.69	2 (13%)
2	NAG	B	501	1,2,6	14,14,15	0.29	0	15,19,21	0.83	1 (6%)
2	NAG	B	502	3,2	14,14,15	0.34	0	15,19,21	0.71	1 (6%)
3	BMA	B	503	2,4	11,11,12	0.27	0	13,15,17	0.90	1 (7%)
4	MAN	B	504	3,2	11,11,12	0.57	0	13,15,17	1.09	2 (15%)
2	NAG	B	505	5,4	14,14,15	0.33	0	15,19,21	1.15	2 (13%)
4	MAN	B	506	3,2	11,11,12	0.51	0	13,15,17	1.43	2 (15%)
5	GAL	B	507	2	11,11,12	0.44	0	13,15,17	0.69	0
6	FUC	B	508	2	9,10,11	0.50	0	13,14,16	0.95	1 (7%)
2	NAG	B	509	4	14,14,15	0.25	0	15,19,21	1.29	2 (13%)
2	NAG	C	501	1,2,6	14,14,15	0.28	0	15,19,21	1.07	2 (13%)
2	NAG	C	502	3,2	14,14,15	0.37	0	15,19,21	0.67	0
3	BMA	C	503	2,4	11,11,12	0.51	0	13,15,17	0.71	0
4	MAN	C	504	3,2	11,11,12	0.39	0	13,15,17	1.18	1 (7%)
2	NAG	C	505	5,4	14,14,15	0.27	0	15,19,21	1.75	2 (13%)
4	MAN	C	506	3,2	11,11,12	0.43	0	13,15,17	0.84	1 (7%)
5	GAL	C	507	2	11,11,12	0.39	0	13,15,17	0.62	0
6	FUC	C	508	2	9,10,11	0.41	0	13,14,16	1.46	2 (15%)
2	NAG	C	509	4	14,14,15	0.35	0	15,19,21	0.61	0
2	NAG	D	501	1,2,6	14,14,15	0.29	0	15,19,21	0.62	0
2	NAG	D	502	3,2	14,14,15	0.29	0	15,19,21	0.58	0
3	BMA	D	503	2,4	11,11,12	0.37	0	13,15,17	0.74	0
4	MAN	D	504	3,2	11,11,12	0.34	0	13,15,17	0.93	1 (7%)
2	NAG	D	505	5,4	14,14,15	0.37	0	15,19,21	1.03	1 (6%)
4	MAN	D	506	3,2	11,11,12	0.61	0	13,15,17	1.74	2 (15%)
5	GAL	D	507	2	11,11,12	0.50	0	13,15,17	0.73	0
6	FUC	D	508	2	9,10,11	0.37	0	13,14,16	0.80	1 (7%)
2	NAG	D	509	4	14,14,15	0.31	0	15,19,21	0.77	1 (6%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	A	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	503	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	504	3,2	-	0/2/19/22	0/1/1/1
2	NAG	A	505	5,4	-	0/6/23/26	0/1/1/1
4	MAN	A	506	3,2	-	0/2/19/22	0/1/1/1
5	GAL	A	507	2	-	0/2/19/22	0/1/1/1
6	FUC	A	508	2	-	0/0/17/20	0/1/1/1
2	NAG	A	509	4	-	0/6/23/26	0/1/1/1
2	NAG	B	501	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	B	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	503	2,4	-	0/2/19/22	0/1/1/1
4	MAN	B	504	3,2	-	0/2/19/22	0/1/1/1
2	NAG	B	505	5,4	-	0/6/23/26	0/1/1/1
4	MAN	B	506	3,2	-	0/2/19/22	0/1/1/1
5	GAL	B	507	2	-	0/2/19/22	0/1/1/1
6	FUC	B	508	2	-	0/0/17/20	0/1/1/1
2	NAG	B	509	4	-	0/6/23/26	0/1/1/1
2	NAG	C	501	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	C	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	C	503	2,4	-	0/2/19/22	0/1/1/1
4	MAN	C	504	3,2	-	0/2/19/22	0/1/1/1
2	NAG	C	505	5,4	-	0/6/23/26	0/1/1/1
4	MAN	C	506	3,2	-	0/2/19/22	0/1/1/1
5	GAL	C	507	2	-	0/2/19/22	0/1/1/1
6	FUC	C	508	2	-	0/0/17/20	0/1/1/1
2	NAG	C	509	4	-	0/6/23/26	0/1/1/1
2	NAG	D	501	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	D	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	D	503	2,4	-	0/2/19/22	0/1/1/1
4	MAN	D	504	3,2	-	0/2/19/22	0/1/1/1
2	NAG	D	505	5,4	-	0/6/23/26	0/1/1/1
4	MAN	D	506	3,2	-	0/2/19/22	0/1/1/1
5	GAL	D	507	2	-	0/2/19/22	0/1/1/1
6	FUC	D	508	2	-	0/0/17/20	0/1/1/1
2	NAG	D	509	4	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	506	MAN	C1-C2	2.00	1.56	1.52

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	505	NAG	O5-C1-C2	-4.67	104.98	111.47
2	B	509	NAG	O5-C1-C2	-3.75	106.25	111.47
2	A	502	NAG	O5-C1-C2	-3.17	107.06	111.47
2	D	505	NAG	O5-C1-C2	-2.92	107.41	111.47
2	A	505	NAG	O5-C1-C2	-2.67	107.76	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	NAG	1	0
6	C	508	FUC	1	0
3	D	503	BMA	1	0
2	D	505	NAG	1	0
4	D	506	MAN	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/227 (92%)	-0.11	0 <span>100</span> <span>100</span>	59, 80, 110, 127	0
1	B	209/227 (92%)	0.28	7 (3%) 47 40	56, 97, 143, 166	0
1	C	208/227 (91%)	0.38	14 (6%) 19 14	60, 103, 163, 178	0
1	D	209/227 (92%)	-0.04	0 <span>100</span> <span>100</span>	58, 81, 116, 137	0
All	All	835/908 (91%)	0.13	21 (2%) 58 53	56, 86, 145, 178	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	324	SER	5.4
1	B	310	HIS	5.3
1	C	275	PHE	5.1
1	C	276	ASN	4.3
1	C	277	TRP	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	505	14/15	0.97	0.19	-0.44	104,107,114,114	0
5	GAL	C	507	11/12	0.92	0.16	-0.60	103,111,114,114	0
5	GAL	A	507	11/12	0.95	0.16	-0.71	104,109,114,115	0
2	NAG	A	501	14/15	0.96	0.14	-0.79	98,110,118,120	0
2	NAG	B	501	14/15	0.89	0.19	-1.00	128,137,142,145	0
5	GAL	D	507	11/12	0.97	0.17	-1.04	91,96,98,99	0
2	NAG	B	505	14/15	0.92	0.14	-1.23	118,120,124,124	0
2	NAG	C	505	14/15	0.97	0.11	-1.44	101,104,106,107	0
5	GAL	B	507	11/12	0.94	0.12	-1.79	129,132,136,137	0
2	NAG	A	502	14/15	0.98	0.14	-1.94	85,91,97,98	0
2	NAG	C	502	14/15	0.94	0.20	-2.11	120,130,137,137	0
2	NAG	D	502	14/15	0.96	0.14	-2.22	101,106,108,114	0
2	NAG	D	505	14/15	0.97	0.16	-2.69	98,102,106,107	0
3	BMA	C	503	11/12	0.93	0.11	-	114,130,135,142	0
2	NAG	B	509	14/15	0.91	0.16	-	147,151,154,155	0
4	MAN	D	506	11/12	0.91	0.16	-	140,147,151,153	0
2	NAG	A	509	14/15	0.93	0.21	-	135,145,149,149	0
2	NAG	C	509	14/15	0.90	0.10	-	167,174,177,177	0
4	MAN	B	504	11/12	0.93	0.11	-	121,126,127,129	0
2	NAG	D	509	14/15	0.92	0.14	-	135,141,143,144	0
4	MAN	C	504	11/12	0.96	0.10	-	100,110,115,115	0
3	BMA	B	503	11/12	0.96	0.09	-	115,117,122,128	0
3	BMA	A	503	11/12	0.95	0.12	-	96,102,108,118	0
4	MAN	A	506	11/12	0.90	0.16	-	127,130,135,142	0
4	MAN	D	504	11/12	0.92	0.19	-	112,119,126,127	0
3	BMA	D	503	11/12	0.94	0.11	-	114,116,124,132	0
2	NAG	C	501	14/15	0.94	0.18	-	138,146,155,160	0
4	MAN	C	506	11/12	0.92	0.10	-	154,159,160,168	0
4	MAN	B	506	11/12	0.94	0.13	-	135,140,142,145	0
6	FUC	C	508	10/11	0.86	0.19	-	142,146,149,149	0
6	FUC	A	508	10/11	0.93	0.23	-	124,129,134,136	0
6	FUC	B	508	10/11	0.90	0.18	-	142,149,153,156	0
2	NAG	B	502	14/15	0.95	0.10	-	108,113,119,120	0
6	FUC	D	508	10/11	0.92	0.18	-	115,119,123,123	0
4	MAN	A	504	11/12	0.96	0.17	-	105,111,122,123	0
2	NAG	D	501	14/15	0.94	0.13	-	107,111,127,128	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.