



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2017 – 04:48 AM EDT

PDB ID : 2HZ4  
Title : Abl kinase domain unligated and in complex with tetrahydrostaurosporine  
Authors : Cowan-Jacob, S.W.; Fendrich, G.; Liebetanz, J.; Fabbro, D.; Manley, P.  
Deposited on : unknown  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*




The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	B	273	
1	C	273	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6518 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2141	1380	348	396	17			
1	B	261	Total	C	N	O	S	0	0	0
			2136	1378	348	393	17			
1	C	259	Total	C	N	O	S	0	0	0
			2117	1366	345	389	17			

- Molecule 2 is 1,2,3,4-TETRAHYDROGEN-STAUROSPORINE (three-letter code: 4ST) (formula: C<sub>28</sub>H<sub>30</sub>N<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			35	28	4	3		
2	B	1	Total	C	N	O	0	0
			35	28	4	3		
2	C	1	Total	C	N	O	0	0
			35	28	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	O 2	0	0
3	B	9	Total 9	O 9	0	0
3	C	8	Total 8	O 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proto-oncogene tyrosine-protein kinase ABL1



- Molecule 1: Proto-oncogene tyrosine-protein kinase ABL1



- Molecule 1: Proto-oncogene tyrosine-protein kinase ABL1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	185.45Å 58.85Å 103.98Å 90.00° 119.02° 90.00°	Depositor
Resolution (Å)	40.00 – 2.80 39.81 – 2.80	Depositor EDS
% Data completeness (in resolution range)	96.2 (40.00-2.80) 96.2 (39.81-2.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.216 , 0.285 0.213 , 0.279	Depositor DCC
$R_{free}$ test set	1203 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.8	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 37.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6518	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 4ST

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.94	2/2195 (0.1%)	1.00	5/2969 (0.2%)
1	B	0.99	2/2190 (0.1%)	1.01	7/2963 (0.2%)
1	C	0.89	0/2169	0.97	7/2932 (0.2%)
All	All	0.94	4/6554 (0.1%)	0.99	19/8864 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	2
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	450	GLU	CD-OE2	6.65	1.32	1.25
1	A	437	MET	SD-CE	6.31	2.13	1.77
1	B	425	PHE	CE2-CZ	5.51	1.47	1.37
1	A	405	TRP	CB-CG	-5.49	1.40	1.50

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	482	ASP	CB-CG-OD2	8.36	125.82	118.30
1	A	276	ASP	CB-CG-OD2	6.28	123.95	118.30
1	C	367	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	B	233	ASP	CB-CG-OD2	6.12	123.80	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	ASP	CB-CG-OD2	6.02	123.72	118.30
1	B	386	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	C	325	ASP	CB-CG-OD2	5.97	123.68	118.30
1	B	332	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	A	381	ASP	CB-CG-OD2	5.86	123.57	118.30
1	C	421	ASP	CB-CG-OD1	5.79	123.51	118.30
1	B	276	ASP	CB-CG-OD2	5.73	123.45	118.30
1	B	381	ASP	CB-CG-OD2	5.68	123.42	118.30
1	C	473	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	C	457	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	B	241	ASP	CB-CG-OD2	5.38	123.14	118.30
1	B	391	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	328	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	C	278	MET	N-CA-C	5.10	124.76	111.00
1	C	445	LEU	CA-CB-CG	5.04	126.89	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	278	MET	Peptide
1	A	309	PRO	Peptide
1	C	309	PRO	Peptide
1	C	440	TYR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2141	0	2098	6	0
1	B	2136	0	2100	9	0
1	C	2117	0	2074	4	0
2	A	35	0	30	0	0
2	B	35	0	30	3	0
2	C	35	0	30	0	0
3	A	2	0	0	0	0
3	B	9	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	8	0	0	0	0
All	All	6518	0	6362	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:437:MET:CE	1:A:437:MET:SD	2.13	1.37
1:A:388:MET:HA	1:B:374:ASN:HD21	1.71	0.56
1:A:360:ILE:HD12	1:A:362:ARG:HG3	1.89	0.53
1:B:290:MET:CE	1:B:313:ILE:HG21	2.40	0.52
1:B:367:ARG:HB2	2:B:501:4ST:HAA1	1.97	0.46
1:A:294:LYS:HG3	1:A:300:GLN:HE22	1.82	0.45
2:B:501:4ST:HAB2	2:B:501:4ST:CBC	2.47	0.45
1:B:484:PRO:HG2	1:B:489:ILE:HD11	1.99	0.44
1:C:294:LYS:HG3	1:C:300:GLN:HE22	1.82	0.44
2:B:501:4ST:HAH	2:B:501:4ST:HAC2	1.99	0.43
1:A:359:PHE:CE1	1:A:387:LEU:HD12	2.55	0.42
1:C:290:MET:HB3	1:C:301:LEU:HB2	2.00	0.42
1:B:428:LEU:O	1:B:428:LEU:HD12	2.20	0.42
1:B:326:TYR:OH	1:B:375:HIS:HD2	2.03	0.41
1:B:316:GLU:OE1	1:B:378:LYS:NZ	2.53	0.41
1:A:395:ALA:HB3	1:A:414:ASN:HB3	2.03	0.40
1:B:396:HIS:ND1	1:B:397:ALA:O	2.43	0.40
1:C:396:HIS:CD2	1:C:397:ALA:O	2.74	0.40
1:B:322:ASN:HD21	1:B:324:LEU:HB2	1.87	0.40
1:C:349:SER:HB3	1:C:490:HIS:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/273 (94%)	236 (92%)	17 (7%)	5 (2%)	9	30
1	B	257/273 (94%)	237 (92%)	19 (7%)	1 (0%)	38	72
1	C	251/273 (92%)	225 (90%)	23 (9%)	3 (1%)	15	44
All	All	766/819 (94%)	698 (91%)	59 (8%)	9 (1%)	15	44

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	275	GLU
1	B	275	GLU
1	C	278	MET
1	A	239	ARG
1	A	455	ASP
1	A	265	SER
1	C	466	GLU
1	A	495	THR
1	C	448	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/239 (96%)	220 (96%)	10 (4%)	33	67
1	B	230/239 (96%)	224 (97%)	6 (3%)	51	83
1	C	227/239 (95%)	218 (96%)	9 (4%)	36	70
All	All	687/717 (96%)	662 (96%)	25 (4%)	40	74

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	248	LEU
1	A	276	ASP
1	A	285	LYS
1	A	308	GLU
1	A	362	ARG
1	A	376	LEU
1	A	387	LEU
1	A	481	SER
1	A	500	SER
1	B	233	ASP
1	B	272	THR
1	B	308	GLU
1	B	341	LEU
1	B	370	LEU
1	B	387	LEU
1	C	305	CYS
1	C	404	LYS
1	C	418	ILE
1	C	437	MET
1	C	438	SER
1	C	446	SER
1	C	447	GLN
1	C	455	ASP
1	C	459	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	300	GLN
1	B	374	ASN
1	B	375	HIS
1	B	479	ASN
1	C	300	GLN
1	C	331	ASN
1	C	396	HIS
1	C	490	HIS
1	C	491	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	4ST	A	501	-	30,42,42	1.49	6 (20%)	25,68,68	2.40	9 (36%)
2	4ST	B	501	-	30,42,42	1.55	8 (26%)	25,68,68	2.53	9 (36%)
2	4ST	C	501	-	30,42,42	1.73	10 (33%)	25,68,68	2.18	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4ST	A	501	-	-	0/4/49/49	0/0/8/8
2	4ST	B	501	-	-	0/4/49/49	0/0/8/8
2	4ST	C	501	-	-	0/4/49/49	0/0/8/8

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	4ST	CAX-CAY	-2.72	1.36	1.41
2	C	501	4ST	CAS-NAP	-2.68	1.32	1.35
2	C	501	4ST	CAX-CAY	-2.57	1.36	1.41
2	B	501	4ST	CAX-CAY	-2.54	1.36	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	4ST	CBB-CBC	-2.41	1.36	1.42
2	C	501	4ST	CBB-CBC	-2.15	1.36	1.42
2	C	501	4ST	CAV-CAT	-2.11	1.34	1.38
2	B	501	4ST	CAN-CBE	2.01	1.54	1.51
2	C	501	4ST	CAF-CAH	2.02	1.41	1.36
2	C	501	4ST	CAN-CBE	2.05	1.54	1.51
2	C	501	4ST	CAT-CAZ	2.09	1.45	1.41
2	A	501	4ST	CAT-CAZ	2.10	1.45	1.41
2	B	501	4ST	CAE-CAG	2.14	1.41	1.36
2	B	501	4ST	CBF-CBD	2.20	1.55	1.52
2	B	501	4ST	CAF-CAH	2.24	1.41	1.36
2	C	501	4ST	CAE-CAG	2.33	1.42	1.36
2	A	501	4ST	CAN-CBE	2.40	1.55	1.51
2	A	501	4ST	CAC-CBI	2.41	1.54	1.51
2	B	501	4ST	OAQ-CBF	2.51	1.47	1.42
2	A	501	4ST	CAE-CAG	2.70	1.43	1.36
2	A	501	4ST	CAF-CAH	3.11	1.44	1.36
2	C	501	4ST	CBF-CBD	3.34	1.56	1.52
2	B	501	4ST	CAC-CBI	3.73	1.56	1.51
2	C	501	4ST	CAC-CBI	3.93	1.56	1.51

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	4ST	CAU-CAM-NAP	-3.48	97.86	101.86
2	A	501	4ST	CAH-CAY-CAX	-2.64	116.92	120.73
2	B	501	4ST	CAU-CAM-NAP	-2.61	98.86	101.86
2	C	501	4ST	OAR-CBI-CAC	-2.32	95.98	106.42
2	C	501	4ST	CAG-CAX-CBA	-2.17	128.46	134.81
2	C	501	4ST	CAU-CAM-NAP	-2.08	99.47	101.86
2	B	501	4ST	CAT-CAZ-CBB	-2.08	106.08	109.68
2	B	501	4ST	CAG-CAX-CBA	-2.07	128.77	134.81
2	A	501	4ST	CAG-CAX-CBA	-2.07	128.77	134.81
2	B	501	4ST	CAH-CAY-CAX	-2.01	117.83	120.73
2	C	501	4ST	OAR-CBE-CAN	2.18	115.45	112.31
2	C	501	4ST	CBA-CAX-CAY	2.18	108.75	106.37
2	A	501	4ST	CAM-NAP-CAS	2.19	116.14	113.78
2	B	501	4ST	CAM-NAP-CAS	2.53	116.51	113.78
2	C	501	4ST	CAG-CAX-CAY	2.69	122.79	119.39
2	A	501	4ST	CAB-OAQ-CBF	2.94	119.72	114.29
2	A	501	4ST	CAM-CAU-CAW	2.95	112.25	109.61
2	B	501	4ST	CAG-CAX-CAY	3.22	123.46	119.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	4ST	CAG-CAX-CAY	3.49	123.81	119.39
2	A	501	4ST	OAR-CBE-CAN	4.45	118.73	112.31
2	C	501	4ST	CAB-OAQ-CBF	4.94	123.42	114.29
2	B	501	4ST	OAR-CBE-CAN	4.97	119.50	112.31
2	B	501	4ST	CAT-CAV-NBG	5.63	110.24	106.66
2	A	501	4ST	CAT-CAV-NBG	6.20	110.60	106.66
2	C	501	4ST	CAT-CAV-NBG	6.67	110.90	106.66
2	B	501	4ST	CAB-OAQ-CBF	6.80	126.86	114.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	4ST	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/273 (95%)	0.04	8 (3%) 49 38	42, 59, 88, 92	0
1	B	261/273 (95%)	-0.03	5 (1%) 67 58	40, 53, 80, 91	0
1	C	259/273 (94%)	0.60	32 (12%) 4 2	59, 75, 101, 109	0
All	All	782/819 (95%)	0.20	45 (5%) 24 15	40, 66, 92, 109	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	248	LEU	5.2
1	C	303	GLY	4.6
1	B	276	ASP	4.6
1	C	257	TYR	4.5
1	C	454	LYS	4.4
1	C	304	VAL	4.2
1	C	391	ASP	4.2
1	C	306	THR	4.0
1	B	275	GLU	3.9
1	C	273	LEU	3.9
1	C	313	ILE	3.8
1	B	278	MET	3.7
1	C	311	PHE	3.7
1	C	276	ASP	3.6
1	C	256	VAL	3.4
1	C	482	ASP	3.2
1	A	259	GLY	3.2
1	C	307	ARG	3.1
1	C	244	MET	3.0
1	C	312	TYR	2.9
1	B	311	PHE	2.9
1	C	267	THR	2.9
1	A	278	MET	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	450	GLU	2.8
1	A	271	LYS	2.8
1	C	447	GLN	2.8
1	A	306	THR	2.7
1	C	272	THR	2.6
1	C	292	GLU	2.6
1	C	478	TRP	2.6
1	C	274	LYS	2.5
1	C	310	PRO	2.4
1	A	274	LYS	2.4
1	C	320	TYR	2.4
1	C	329	GLU	2.4
1	C	278	MET	2.4
1	A	308	GLU	2.3
1	A	311	PHE	2.3
1	C	453	GLU	2.2
1	C	255	GLU	2.1
1	A	307	ARG	2.1
1	C	262	LYS	2.1
1	C	246	HIS	2.1
1	C	372	GLY	2.0
1	B	244	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	4ST	B	501	35/35	0.95	0.18	-0.58	42,47,47,48	0
2	4ST	C	501	35/35	0.91	0.19	-0.87	76,78,78,78	0
2	4ST	A	501	35/35	0.94	0.18	-1.05	64,65,70,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.