



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:19 pm GMT

PDB ID : 3I0G  
Title : Crystal structure of GTB C80S/C196S + DA + UDP-Gal  
Authors : Schuman, B.; Persson, M.; Landry, R.C.; Polakowski, R.; Weadge, J.T.; Seto, N.O.L.; Borisova, S.; Palcic, M.M.; Evans, S.V.  
Deposited on : 2009-06-25  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1307 (1.40-1.40)
Clashscore	112137	1411 (1.40-1.40)
Ramachandran outliers	110173	1373 (1.40-1.40)
Sidechain outliers	110143	1372 (1.40-1.40)
RSRZ outliers	101464	1315 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	<div> <div>7%</div> <div> <div></div> <div>76%</div> <div>15%</div> <div>7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FUC	A	356	-	-	-	X
4	UDP	A	357	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GAL	A	359	X	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2634 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABO glycosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	4	0
			2201	1426	378	388	9			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	80	SER	CYS	ENGINEERED	UNP Q70V26
A	196	SER	CYS	ENGINEERED	UNP Q70V26
A	355	GLU	-	SEE REMARK 999	UNP Q70V26


- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	2	Total	C	O	0	0
			27	18	9		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
4	A	1	Total	C	N	O	P	0	0
			24	9	2	11	2		

- Molecule 5 is SUGAR (BETA-D-GALACTOSE) (three-letter code: GAL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	344	Total 344	O 344	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ABO glycosyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.55Å 149.28Å 79.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.87 – 1.40 19.86 – 1.40	Depositor EDS
% Data completeness (in resolution range)	95.3 (19.87-1.40) 95.3 (19.86-1.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.195 , 0.230 0.189 , 0.222	Depositor DCC
$R_{free}$ test set	2993 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2634	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, MN, GAL, BHG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.36	4/2278 (0.2%)	1.48	29/3092 (0.9%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	LYS	CE-NZ	7.21	1.67	1.49
1	A	127	VAL	CB-CG1	-6.31	1.39	1.52
1	A	325	TRP	CZ3-CH2	5.33	1.48	1.40
1	A	300	TRP	CE3-CZ3	5.01	1.47	1.38

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	213	ASP	CB-CG-OD2	17.41	133.96	118.30
1	A	282	ARG	NE-CZ-NH1	16.49	128.54	120.30
1	A	282	ARG	NE-CZ-NH2	-15.75	112.43	120.30
1	A	213	ASP	CB-CG-OD1	-13.95	105.75	118.30
1	A	302	ASP	CB-CG-OD1	11.22	128.40	118.30
1	A	241	ARG	NE-CZ-NH2	10.05	125.33	120.30
1	A	217	ARG	NE-CZ-NH1	-9.71	115.45	120.30
1	A	146[A]	ARG	NE-CZ-NH1	-8.87	115.86	120.30
1	A	146[B]	ARG	NE-CZ-NH1	-8.87	115.86	120.30
1	A	340	ARG	NE-CZ-NH1	8.71	124.65	120.30
1	A	241	ARG	NE-CZ-NH1	-8.31	116.14	120.30
1	A	154	ASP	CB-CG-OD1	7.90	125.41	118.30
1	A	269	PHE	CB-CG-CD2	7.23	125.86	120.80
1	A	218	ASP	CB-CG-OD1	7.04	124.64	118.30
1	A	211	ASP	CB-CG-OD2	-6.64	112.32	118.30
1	A	110	ARG	NE-CZ-NH1	-6.52	117.04	120.30
1	A	244	PHE	CB-CG-CD1	-6.26	116.42	120.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	A	282	ARG	CD-NE-CZ	5.97	131.96	123.60
1	A	269	PHE	CB-CG-CD1	-5.83	116.72	120.80
1	A	340	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	259	ASP	CB-CG-OD1	5.74	123.46	118.30
1	A	146[A]	ARG	NE-CZ-NH2	5.55	123.08	120.30
1	A	146[B]	ARG	NE-CZ-NH2	5.55	123.08	120.30
1	A	341	PHE	CB-CG-CD2	-5.38	117.03	120.80
1	A	195	PHE	CB-CG-CD2	-5.36	117.05	120.80
1	A	211	ASP	CB-CG-OD1	5.33	123.09	118.30
1	A	244	PHE	CB-CG-CD2	5.04	124.33	120.80
1	A	286	GLN	CB-CA-C	5.03	120.46	110.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2201	0	2170	50	0
2	A	27	0	31	1	0
3	A	1	0	0	0	0
4	A	49	0	22	0	0
5	A	12	0	11	0	0
6	A	344	0	0	24	1
All	All	2634	0	2234	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:LYS:HE3	6:A:558:HOH:O	1.29	1.23
1:A:288:MET:CE	6:A:499:HOH:O	1.94	1.16

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:ARG:HD2	6:A:377:HOH:O	1.51	1.10
1:A:240:SER:CA	6:A:521:HOH:O	1.98	1.08
1:A:197:GLU:HG2	6:A:436:HOH:O	1.55	1.07
1:A:82:LYS:H	1:A:82:LYS:HE2	1.20	1.05
1:A:199:ARG:HD3	1:A:199:ARG:H	1.16	1.05
1:A:198:ARG:NH1	1:A:199:ARG:HE	1.59	1.00
1:A:199:ARG:HH11	1:A:199:ARG:HG2	1.25	0.98
1:A:199:ARG:H	1:A:199:ARG:CD	1.76	0.96
1:A:188:ARG:CD	6:A:488:HOH:O	2.18	0.92
1:A:82:LYS:H	1:A:82:LYS:CE	1.84	0.91
1:A:288:MET:HE3	6:A:499:HOH:O	1.61	0.89
1:A:188:ARG:CG	6:A:488:HOH:O	2.19	0.89
1:A:199:ARG:N	1:A:199:ARG:HD3	1.90	0.87
1:A:296:ILE:CA	6:A:543:HOH:O	2.23	0.85
1:A:197:GLU:CG	6:A:436:HOH:O	2.15	0.84
1:A:233:HIS:HD2	1:A:235:SER:H	1.23	0.84
1:A:282:ARG:HD3	6:A:565:HOH:O	1.80	0.82
1:A:297:GLU:OE1	1:A:301:HIS:HD2	1.64	0.81
1:A:198:ARG:HH11	1:A:199:ARG:HE	1.31	0.79
1:A:203:GLU:HG3	6:A:562:HOH:O	1.83	0.78
1:A:188:ARG:HG2	6:A:488:HOH:O	1.83	0.76
1:A:82:LYS:HE2	1:A:82:LYS:N	1.99	0.76
1:A:288:MET:HE1	6:A:499:HOH:O	1.69	0.74
1:A:199:ARG:HG2	1:A:199:ARG:NH1	2.02	0.63
1:A:242:GLU:H	1:A:242:GLU:CD	2.02	0.62
1:A:301:HIS:HE1	6:A:561:HOH:O	1.85	0.59
1:A:233:HIS:CD2	1:A:235:SER:H	2.14	0.58
1:A:188:ARG:HD2	6:A:488:HOH:O	1.97	0.57
1:A:258:LYS:NZ	1:A:258:LYS:H	2.02	0.57
1:A:146[A]:ARG:NH2	6:A:552:HOH:O	2.17	0.54
1:A:233:HIS:HE1	2:A:1:BHG:O4	1.93	0.52
1:A:305:HIS:HE1	6:A:44:HOH:O	1.92	0.52
1:A:251:GLN:H	1:A:251:GLN:CD	2.15	0.50
1:A:113:ASN:HD21	1:A:146[B]:ARG:HH11	1.59	0.50
1:A:82:LYS:CD	1:A:82:LYS:H	2.24	0.49
1:A:258:LYS:H	1:A:258:LYS:HZ2	1.60	0.49
1:A:188:ARG:NE	6:A:488:HOH:O	2.43	0.49
1:A:297:GLU:OE1	1:A:301:HIS:CD2	2.55	0.48
1:A:305:HIS:HD2	6:A:45:HOH:O	1.96	0.47
1:A:175:VAL:HA	1:A:176:GLY:HA3	1.57	0.47
1:A:113:ASN:HD21	1:A:146[B]:ARG:NH1	2.14	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:GLU:CD	6:A:436:HOH:O	2.48	0.46
1:A:198:ARG:NH1	1:A:199:ARG:NE	2.44	0.45
1:A:333:PRO:HD2	1:A:336:LEU:HD12	1.98	0.45
1:A:261:GLY:HA2	1:A:311:LEU:HD13	2.00	0.42
1:A:82:LYS:NZ	6:A:434:HOH:O	2.53	0.41
1:A:285:HIS:HD2	6:A:389:HOH:O	2.04	0.41
1:A:251:GLN:N	1:A:251:GLN:CD	2.75	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:497:HOH:O	6:A:497:HOH:O[4_555]	2.00	0.20

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/287 (93%)	260 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	239/255 (94%)	229 (96%)	10 (4%)	34 6

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	LYS
1	A	146[A]	ARG
1	A	146[B]	ARG
1	A	185[A]	SER
1	A	185[B]	SER
1	A	199	ARG
1	A	227	PRO
1	A	251	GLN
1	A	258	LYS
1	A	269	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	155	GLN
1	A	233	HIS
1	A	278	GLN
1	A	285	HIS
1	A	301	HIS
1	A	305	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BHG	A	1	2	16,17,18	1.15	2 (12%)	16,21,23	1.26	2 (12%)
2	FUC	A	356	2	9,10,11	1.16	0	13,14,16	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BHG	A	1	2	-	0/9/25/29	0/1/1/1
2	FUC	A	356	2	-	0/0/17/20	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	BHG	O1-C1'	2.02	1.48	1.43
2	A	1	BHG	C4-C5	2.87	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	BHG	C1'-O1-C1	-3.03	108.67	113.87
2	A	1	BHG	C3-C4-C5	-2.30	107.81	110.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	BHG	1	0

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	UDP	A	357	-	21,26,26	2.06	6 (28%)	22,40,40	1.68	1 (4%)
4	UDP	A	358	3	19,25,26	1.77	4 (21%)	18,37,40	1.95	2 (11%)
5	GAL	A	359	-	12,12,12	1.39	1 (8%)	17,17,17	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UDP	A	357	-	-	0/12/32/32	0/2/2/2
4	UDP	A	358	3	-	0/9/31/32	0/2/2/2
5	GAL	A	359	-	1/1/5/5	0/2/22/22	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	357	UDP	PB-O2B	2.04	1.63	1.54
4	A	358	UDP	PA-O1A	2.16	1.59	1.50
4	A	357	UDP	O4'-C1'	2.18	1.44	1.41
4	A	358	UDP	O4'-C1'	2.46	1.44	1.41
4	A	357	UDP	PA-O1A	2.88	1.61	1.50
5	A	359	GAL	O1-C1	3.84	1.52	1.39
4	A	357	UDP	C4-N3	4.12	1.40	1.33
4	A	358	UDP	C6-N1	4.41	1.41	1.35
4	A	357	UDP	PB-O1B	4.43	1.65	1.50
4	A	358	UDP	C4-N3	4.69	1.41	1.33
4	A	357	UDP	C6-N1	4.94	1.42	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	358	UDP	C4'-O4'-C1'	-2.79	106.80	109.77
4	A	357	UDP	C4-N3-C2	6.77	119.95	114.13
4	A	358	UDP	C4-N3-C2	7.23	120.34	114.13

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	359	GAL	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/287 (93%)	0.59	19 (7%) 17 16	9, 15, 28, 38	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	GLY	7.1
1	A	175	VAL	4.6
1	A	186	MET	4.2
1	A	203	GLU	3.7
1	A	71	TYR	3.7
1	A	161	ARG	3.5
1	A	292	GLN	3.4
1	A	70	VAL	2.8
1	A	345	PRO	2.6
1	A	157	ALA	2.5
1	A	187	ARG	2.5
1	A	199	ARG	2.5
1	A	202	SER	2.4
1	A	240	SER	2.3
1	A	251	GLN	2.3
1	A	256	ILE	2.3
1	A	111	LEU	2.1
1	A	124	LYS	2.1
1	A	212	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FUC	A	356	10/11	0.87	0.11	2.05	14,15,16,17	0
2	BHG	A	1	17/18	0.92	0.10	-0.23	12,15,23,27	0

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	UDP	A	357	25/25	0.82	0.18	3.57	19,21,37,39	8
5	GAL	A	359	12/12	0.81	0.16	3.32	19,21,25,26	0
4	UDP	A	358	24/25	0.90	0.13	0.15	14,20,22,25	0
3	MN	A	2	1/1	0.92	0.15	-	25,25,25,25	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.