



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:29 am GMT

PDB ID : 4I09
Title : structure of the mutant Catabolite gene activator protein V132L
Authors : Pohl, E.; Townsend, P.D.; Rodgers, T.; Burnell, D.; McLeish, T.C.B.; Wilson, M.R.; Cann, M.J.
Deposited on : 2012-11-16
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1316 (2.04-2.04)
Clashscore	112137	1394 (2.04-2.04)
Ramachandran outliers	110173	1383 (2.04-2.04)
Sidechain outliers	110143	1383 (2.04-2.04)
RSRZ outliers	101464	1319 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>5%</div> <div> <div></div> <div>73%</div> <div>17%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	222	<div> <div>3%</div> <div> <div></div> <div>71%</div> <div>18%</div> <div>•</div> <div>9%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catabolite gene activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	1	0
			1600	1012	277	303	8			
1	B	201	Total	C	N	O	S	0	1	0
			1560	988	275	289	8			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP P0ACJ8
A	-10	ARG	-	EXPRESSION TAG	UNP P0ACJ8
A	-9	GLY	-	EXPRESSION TAG	UNP P0ACJ8
A	-8	SER	-	EXPRESSION TAG	UNP P0ACJ8
A	-7	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-6	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-5	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-4	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-3	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-2	HIS	-	EXPRESSION TAG	UNP P0ACJ8
A	-1	GLY	-	EXPRESSION TAG	UNP P0ACJ8
A	0	SER	-	EXPRESSION TAG	UNP P0ACJ8
A	132	LEU	VAL	ENGINEERED MUTATION	UNP P0ACJ8
B	-11	MET	-	EXPRESSION TAG	UNP P0ACJ8
B	-10	ARG	-	EXPRESSION TAG	UNP P0ACJ8
B	-9	GLY	-	EXPRESSION TAG	UNP P0ACJ8
B	-8	SER	-	EXPRESSION TAG	UNP P0ACJ8
B	-7	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-6	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-5	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-4	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-3	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-2	HIS	-	EXPRESSION TAG	UNP P0ACJ8
B	-1	GLY	-	EXPRESSION TAG	UNP P0ACJ8
B	0	SER	-	EXPRESSION TAG	UNP P0ACJ8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	132	LEU	VAL	ENGINEERED MUTATION	UNP P0ACJ8

- Molecule 2 is ADENOSINE-3',5'-CYCLIC-MONOPHOSPHATE (three-letter code: CMP) (formula: $C_{10}H_{12}N_5O_6P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		
2	B	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		
3	B	66	Total	O	0	0
			66	66		

- Molecule 1: Catabolite gene activator



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.26Å 93.09Å 105.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.37 – 2.05 45.37 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.37-2.05) 99.0 (45.37-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.201 , 0.256 0.204 , 0.263	Depositor DCC
R_{free} test set	1560 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3328	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	3/1628 (0.2%)	0.50	0/2198
1	B	0.94	0/1587	0.48	0/2143
All	All	0.97	3/3215 (0.1%)	0.49	0/4341

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	TRP	CD2-CE2	6.75	1.49	1.41
1	A	124	ARG	CZ-NH1	5.45	1.40	1.33
1	A	14	TRP	CD2-CE2	5.37	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1600	0	1617	28	0
1	B	1560	0	1568	53	0
2	A	22	0	11	1	0
2	B	22	0	11	1	0
3	A	58	0	0	5	0
3	B	66	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3328	0	3207	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:301:CMP:C2	2:B:301:CMP:H2	0.97	1.50
2:A:301:CMP:H2	2:A:301:CMP:C2	0.97	1.47
1:A:181:ARG:NH1	3:A:405:HOH:O	1.82	1.10
1:B:27:LYS:HD3	1:B:88:ARG:HH12	1.25	0.98
1:B:159:THR:HG22	1:B:160:HIS:H	1.32	0.94
1:B:27:LYS:CD	1:B:88:ARG:HH12	1.83	0.92
1:B:160:HIS:HD2	1:B:162:ASP:H	1.13	0.92
1:A:126:GLN:HB3	3:A:418:HOH:O	1.72	0.89
1:B:154:GLN:HB2	1:B:155:PRO:HD2	1.55	0.87
1:B:27:LYS:HD3	1:B:88:ARG:NH1	1.91	0.84
1:A:78:GLU:HG3	3:A:435:HOH:O	1.80	0.81
1:B:160:HIS:CD2	1:B:162:ASP:H	1.99	0.76
1:B:160:HIS:HB3	1:B:163:GLY:O	1.87	0.75
1:B:159:THR:HG22	1:B:160:HIS:N	2.05	0.69
1:B:26:SER:O	1:B:27:LYS:HB2	1.92	0.69
1:B:172:GLU:O	1:B:176:ILE:CD1	2.41	0.68
1:B:165:GLN:HG3	1:B:205:VAL:HG22	1.75	0.66
1:B:88:ARG:HD2	3:B:434:HOH:O	1.96	0.65
1:B:170:ARG:HH12	1:B:181:ARG:HD2	1.61	0.65
1:B:154:GLN:CB	1:B:155:PRO:HD2	2.27	0.64
1:B:88:ARG:CD	3:B:434:HOH:O	2.44	0.63
1:B:41:TYR:HB2	1:B:71:ILE:HG13	1.79	0.63
1:A:200:HIS:O	1:A:203:THR:HG23	2.02	0.60
1:A:160:HIS:CD2	1:A:161:PRO:HD2	2.37	0.59
1:A:30:LEU:N	1:A:30:LEU:HD12	2.18	0.58
1:A:160:HIS:CG	1:A:161:PRO:HD2	2.39	0.57
1:B:168:ILE:HG13	1:B:172:GLU:HB2	1.85	0.57
1:A:90:LYS:NZ	3:A:453:HOH:O	2.37	0.57
1:A:208:GLY:O	1:A:209:THR:C	2.44	0.56
1:B:172:GLU:O	1:B:176:ILE:HD13	2.04	0.56
1:A:46:GLY:HA2	1:A:91:THR:HG21	1.85	0.56
1:B:159:THR:CG2	1:B:160:HIS:H	2.10	0.56
1:B:203:THR:HG22	3:B:448:HOH:O	2.05	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:LYS:HE3	3:A:433:HOH:O	2.05	0.55
1:B:51:LEU:CD2	1:B:61:ILE:HG12	2.37	0.55
1:B:27:LYS:HD2	1:B:88:ARG:HH12	1.71	0.54
1:A:111:PRO:HG3	1:B:111:PRO:HG3	1.89	0.54
1:B:26:SER:O	1:B:27:LYS:CB	2.57	0.53
1:B:160:HIS:CB	1:B:163:GLY:O	2.57	0.53
1:A:160:HIS:HD2	1:A:162:ASP:H	1.56	0.52
1:A:30:LEU:H	1:A:30:LEU:HD12	1.74	0.52
1:B:176:ILE:H	1:B:176:ILE:HD12	1.74	0.51
1:A:47:SER:OG	1:A:90:LYS:HE2	2.10	0.51
1:B:176:ILE:N	1:B:176:ILE:HD12	2.26	0.50
1:B:208:GLY:O	1:B:209:THR:CB	2.59	0.50
1:B:171:GLN:HG2	1:B:181:ARG:HH21	1.75	0.50
1:A:9:ASP:HB2	1:A:10:PRO:HD2	1.94	0.50
1:B:154:GLN:HB2	1:B:155:PRO:CD	2.34	0.50
1:A:160:HIS:HB3	1:A:163:GLY:O	2.13	0.49
1:B:168:ILE:CG1	1:B:172:GLU:HB2	2.42	0.49
1:B:53:LYS:HG3	1:B:86:TRP:CZ3	2.48	0.49
1:A:73:GLU:HG3	1:A:117:LEU:HD11	1.94	0.48
1:B:154:GLN:CB	1:B:155:PRO:CD	2.92	0.48
1:B:51:LEU:HD23	1:B:61:ILE:HG12	1.96	0.48
1:B:33:GLN:NE2	1:B:86:TRP:CZ3	2.82	0.47
1:B:51:LEU:HD23	1:B:61:ILE:CG1	2.44	0.47
1:A:62:LEU:HD11	1:B:129:SER:HB3	1.97	0.47
1:B:139:ASP:O	1:B:143[B]:ARG:HG3	2.15	0.46
1:A:26:SER:O	1:A:27:LYS:HB2	2.17	0.45
1:B:51:LEU:HD21	1:B:61:ILE:HG12	1.99	0.45
1:B:172:GLU:O	1:B:176:ILE:HD12	2.14	0.45
1:B:58:LYS:HD2	1:B:178:GLY:O	2.17	0.44
1:A:136:ALA:HB1	1:B:178:GLY:HA3	1.98	0.44
1:B:58:LYS:HD2	1:B:179:CYS:HA	2.00	0.43
1:B:9:ASP:HA	1:B:10:PRO:HD2	1.88	0.42
1:B:169:THR:OG1	1:B:172:GLU:HG3	2.20	0.42
1:B:170:ARG:HH11	1:B:170:ARG:HB2	1.84	0.42
1:A:137:PHE:CG	1:B:54:ASP:HB3	2.55	0.41
1:A:189:LYS:HB3	1:A:189:LYS:HE2	1.74	0.41
1:A:73:GLU:CB	1:A:117:LEU:HD11	2.51	0.41
1:B:170:ARG:NH1	1:B:181:ARG:HG2	2.35	0.41
1:B:150:ASN:HA	1:B:150:ASN:HD22	1.64	0.41
1:B:53:LYS:CG	1:B:86:TRP:CZ3	3.02	0.41
1:A:90:LYS:HB3	1:A:90:LYS:HE2	1.88	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:HIS:HD2	1:A:102:LYS:NZ	2.18	0.40
1:A:137:PHE:CD1	1:B:54:ASP:HB3	2.57	0.40
1:B:51:LEU:CD2	1:B:61:ILE:CG1	3.00	0.40
1:A:110:ASN:HA	1:A:111:PRO:HD2	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/222 (91%)	197 (97%)	6 (3%)	0	100	100
1	B	200/222 (90%)	190 (95%)	9 (4%)	1 (0%)	32	21
All	All	403/444 (91%)	387 (96%)	15 (4%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	168	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/191 (91%)	161 (93%)	12 (7%)	18	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	164/191 (86%)	157 (96%)	7 (4%)	33	25
All	All	337/382 (88%)	318 (94%)	19 (6%)	24	15

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	SER
1	A	28	SER
1	A	36	LYS
1	A	60	MET
1	A	71	ILE
1	A	81	GLN
1	A	91	THR
1	A	101	LYS
1	A	153	LYS
1	A	170	ARG
1	A	189	LYS
1	A	203	THR
1	B	13	GLU
1	B	28	SER
1	B	53	LYS
1	B	78	GLU
1	B	126	GLN
1	B	192	GLU
1	B	203	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	105	GLN
1	A	108	GLN
1	A	134	ASN
1	A	160	HIS
1	A	194	GLN
1	B	33	GLN
1	B	126	GLN
1	B	150	ASN
1	B	154	GLN
1	B	160	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CMP	A	301	-	21,25,25	1.91	5 (23%)	22,39,39	2.75	8 (36%)
2	CMP	B	301	-	21,25,25	1.35	3 (14%)	22,39,39	2.48	6 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CMP	A	301	-	-	0/0/31/31	0/4/4/4
2	CMP	B	301	-	-	0/0/31/31	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	CMP	O3'-C3'	-2.47	1.40	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	CMP	C5-N7	-2.27	1.31	1.39
2	B	301	CMP	C2-N3	2.16	1.35	1.32
2	A	301	CMP	C2-N1	2.55	1.38	1.33
2	A	301	CMP	C8-N7	2.56	1.39	1.34
2	B	301	CMP	P-O3'	3.70	1.64	1.58
2	A	301	CMP	C2-N3	3.74	1.38	1.32
2	A	301	CMP	P-O3'	5.03	1.66	1.58

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	CMP	N3-C2-N1	-8.60	121.36	128.86
2	B	301	CMP	N3-C2-N1	-7.99	121.90	128.86
2	A	301	CMP	O5'-P-O3'	-5.59	98.29	105.81
2	B	301	CMP	O5'-P-O3'	-3.81	100.69	105.81
2	A	301	CMP	C4'-O4'-C1'	-3.34	106.21	109.77
2	A	301	CMP	O3'-C3'-C4'	-2.95	108.42	110.69
2	B	301	CMP	O3'-C3'-C4'	-2.62	108.67	110.69
2	B	301	CMP	C4'-O4'-C1'	-2.47	107.14	109.77
2	B	301	CMP	C4-C5-N7	-2.25	107.23	109.41
2	A	301	CMP	C4-C5-N7	-2.10	107.38	109.41
2	A	301	CMP	C1'-N9-C4	-2.03	123.14	126.64
2	A	301	CMP	O3'-C3'-C2'	2.05	117.72	115.67
2	A	301	CMP	O2P-P-O1P	3.99	121.58	108.63
2	B	301	CMP	O2P-P-O1P	5.15	125.36	108.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	CMP	1	0
2	B	301	CMP	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/222 (91%)	0.15	12 (5%) 23 25	21, 38, 76, 87	0
1	B	201/222 (90%)	0.12	7 (3%) 44 49	22, 40, 87, 97	0
All	All	405/444 (91%)	0.14	19 (4%) 32 35	21, 39, 79, 97	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	THR	4.7
1	A	207	TYR	4.1
1	A	163	GLY	3.6
1	A	161	PRO	3.2
1	A	206	VAL	3.0
1	B	206	VAL	2.9
1	A	6	PRO	2.7
1	A	162	ASP	2.6
1	B	208	GLY	2.6
1	B	209	THR	2.5
1	B	152	ALA	2.4
1	B	205	VAL	2.3
1	A	164	MET	2.2
1	A	208	GLY	2.2
1	B	155	PRO	2.2
1	A	195	ASN	2.1
1	A	160	HIS	2.0
1	A	200	HIS	2.0
1	B	200	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CMP	B	301	22/22	0.99	0.11	-0.51	18,21,24,26	0
2	CMP	A	301	22/22	0.99	0.10	-0.56	20,23,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.