



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2017 – 08:11 AM EST

PDB ID : 4I1I
Title : Crystal structure of a putative Cytosolic malate dehydrogenase from Leishmania major Friedlin in complex with NAD
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : unknown
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>5%</div> <div>7%</div> </div> </div>
1	B	345	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>•</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	401	-	-	-	X
3	PO4	B	402	-	-	-	X
4	EDO	A	403	-	-	-	X
4	EDO	A	404	-	-	-	X
4	EDO	A	405	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	12	0
			2360	1491	411	445	13			
1	B	313	Total	C	N	O	S	0	3	0
			2281	1440	406	424	11			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q4Q7X6
A	-19	ALA	-	EXPRESSION TAG	UNP Q4Q7X6
A	-18	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-17	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-16	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-15	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-14	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-13	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
A	-12	MET	-	EXPRESSION TAG	UNP Q4Q7X6
A	-11	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
A	-10	THR	-	EXPRESSION TAG	UNP Q4Q7X6
A	-9	LEU	-	EXPRESSION TAG	UNP Q4Q7X6
A	-8	GLU	-	EXPRESSION TAG	UNP Q4Q7X6
A	-7	ALA	-	EXPRESSION TAG	UNP Q4Q7X6
A	-6	GLN	-	EXPRESSION TAG	UNP Q4Q7X6
A	-5	THR	-	EXPRESSION TAG	UNP Q4Q7X6
A	-4	GLN	-	EXPRESSION TAG	UNP Q4Q7X6
A	-3	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
A	-2	PRO	-	EXPRESSION TAG	UNP Q4Q7X6
A	-1	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
A	0	SER	-	EXPRESSION TAG	UNP Q4Q7X6
B	-20	MET	-	EXPRESSION TAG	UNP Q4Q7X6
B	-19	ALA	-	EXPRESSION TAG	UNP Q4Q7X6
B	-18	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
B	-17	HIS	-	EXPRESSION TAG	UNP Q4Q7X6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
B	-15	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
B	-14	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
B	-13	HIS	-	EXPRESSION TAG	UNP Q4Q7X6
B	-12	MET	-	EXPRESSION TAG	UNP Q4Q7X6
B	-11	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
B	-10	THR	-	EXPRESSION TAG	UNP Q4Q7X6
B	-9	LEU	-	EXPRESSION TAG	UNP Q4Q7X6
B	-8	GLU	-	EXPRESSION TAG	UNP Q4Q7X6
B	-7	ALA	-	EXPRESSION TAG	UNP Q4Q7X6
B	-6	GLN	-	EXPRESSION TAG	UNP Q4Q7X6
B	-5	THR	-	EXPRESSION TAG	UNP Q4Q7X6
B	-4	GLN	-	EXPRESSION TAG	UNP Q4Q7X6
B	-3	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
B	-2	PRO	-	EXPRESSION TAG	UNP Q4Q7X6
B	-1	GLY	-	EXPRESSION TAG	UNP Q4Q7X6
B	0	SER	-	EXPRESSION TAG	UNP Q4Q7X6

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	312	Total	O	0	0
			312	312		
5	B	311	Total	O	0	0
			311	311		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malate dehydrogenase



- Molecule 1: Malate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.82Å 65.13Å 77.49Å 90.00° 110.67° 90.00°	Depositor
Resolution (Å)	48.50 – 1.50 48.45 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.50-1.50) 99.3 (48.45-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 1.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.145 , 0.166 0.145 , 0.166	Depositor DCC
R_{free} test set	4794 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5343	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2434	0.83	5/3315 (0.2%)
1	B	0.73	0/2328	0.77	2/3173 (0.1%)
All	All	0.74	0/4762	0.80	7/6488 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	25	ARG	NE-CZ-NH1	10.07	125.34	120.30
1	B	25	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	A	25[A]	ARG	NE-CZ-NH1	8.01	124.31	120.30
1	A	25[B]	ARG	NE-CZ-NH1	8.01	124.31	120.30
1	A	25[A]	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	A	25[B]	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	A	91	ARG	NE-CZ-NH2	-5.87	117.36	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2360	0	2389	16	0
1	B	2281	0	2294	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	26	1	0
3	A	5	0	0	1	0
3	B	10	0	0	0	0
4	A	16	0	24	3	0
4	B	4	0	6	0	0
5	A	312	0	0	3	0
5	B	311	0	0	2	0
All	All	5343	0	4739	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82[B]:ILE:HD11	1:A:253[B]:ILE:HD11	1.52	0.90
1:A:82[B]:ILE:CD1	1:A:253[B]:ILE:HD11	2.14	0.76
1:B:165:SER:OG	1:B:169:ARG:NH1	2.23	0.70
1:A:165:SER:OG	1:A:169:ARG:NH1	2.25	0.69
1:A:29:LEU:HD12	1:A:35[A]:VAL:HG12	1.76	0.67
1:A:277[A]:SER:OG	4:A:403:EDO:H21	1.97	0.64
1:A:14[B]:GLN:NE2	5:A:705:HOH:O	2.36	0.58
1:B:14:GLN:NE2	5:B:774:HOH:O	2.37	0.58
1:A:85[B]:MET:SD	1:A:139:LEU:HD11	2.48	0.54
1:A:85[B]:MET:SD	1:A:139:LEU:CD1	2.97	0.52
4:A:404:EDO:H12	5:A:564:HOH:O	2.10	0.50
1:A:128:VAL:O	2:A:400:NAD:H2N	2.12	0.49
1:A:177:GLN:HG3	5:A:679:HOH:O	2.12	0.49
1:A:82[B]:ILE:HD11	1:A:253[B]:ILE:CD1	2.33	0.48
1:A:29:LEU:HD12	1:A:35[A]:VAL:CG1	2.44	0.47
1:A:277[A]:SER:OG	4:A:403:EDO:C2	2.62	0.46
1:A:112:GLY:HA3	1:A:142[B]:SER:OG	2.16	0.46
1:A:25[B]:ARG:NH2	3:A:401:PO4:O3	2.48	0.46
1:B:105:ALA:HA	1:B:138:ILE:HD13	2.00	0.44
1:B:233[B]:ARG:NH2	5:B:731:HOH:O	2.02	0.43
1:B:130:ASN:OD1	1:B:187:HIS:HD2	2.04	0.40
1:A:85[B]:MET:CE	1:A:111:GLN:OE1	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/345 (95%)	325 (99%)	3 (1%)	0	100	100
1	B	312/345 (90%)	310 (99%)	2 (1%)	0	100	100
All	All	640/690 (93%)	635 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/266 (92%)	245 (100%)	0	100	100
1	B	231/266 (87%)	231 (100%)	0	100	100
All	All	476/532 (90%)	476 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	400	-	41,48,48	1.09	5 (12%)	43,73,73	1.76	6 (13%)
3	PO4	A	401	-	4,4,4	1.36	1 (25%)	6,6,6	0.68	0
4	EDO	A	402	-	3,3,3	0.68	0	2,2,2	0.24	0
4	EDO	A	403	-	3,3,3	1.02	0	2,2,2	0.63	0
4	EDO	A	404	-	3,3,3	0.92	0	2,2,2	0.29	0
4	EDO	A	405	-	3,3,3	0.32	0	2,2,2	0.16	0
3	PO4	B	401	-	4,4,4	2.14	1 (25%)	6,6,6	0.42	0
3	PO4	B	402	-	4,4,4	0.85	0	6,6,6	0.81	0
4	EDO	B	403	-	3,3,3	0.29	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	400	-	-	0/22/62/62	0/5/5/5
3	PO4	A	401	-	-	0/0/0/0	0/0/0/0
4	EDO	A	402	-	-	0/1/1/1	0/0/0/0
4	EDO	A	403	-	-	0/1/1/1	0/0/0/0
4	EDO	A	404	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	405	-	-	0/1/1/1	0/0/0/0
3	PO4	B	401	-	-	0/0/0/0	0/0/0/0
3	PO4	B	402	-	-	0/0/0/0	0/0/0/0
4	EDO	B	403	-	-	0/1/1/1	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAD	C5N-C4N	-2.52	1.34	1.38
2	A	400	NAD	O4D-C1D	2.01	1.44	1.41
2	A	400	NAD	C2A-N3A	2.10	1.35	1.32
3	A	401	PO4	P-O1	2.33	1.55	1.50
2	A	400	NAD	C5A-C4A	2.34	1.45	1.40
2	A	400	NAD	O7N-C7N	2.80	1.30	1.24
3	B	401	PO4	P-O1	4.00	1.59	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	NAD	N3A-C2A-N1A	-5.86	123.75	128.86
2	A	400	NAD	C3N-C7N-N7N	-4.86	112.22	117.77
2	A	400	NAD	C1B-N9A-C4A	-3.55	120.50	126.64
2	A	400	NAD	C4B-O4B-C1B	-2.74	106.85	109.77
2	A	400	NAD	N6A-C6A-N1A	2.46	123.65	118.77
2	A	400	NAD	O7N-C7N-C3N	4.49	124.88	119.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	NAD	1	0
3	A	401	PO4	1	0
4	A	403	EDO	2	0
4	A	404	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/345 (92%)	-0.09	7 (2%) 62 68	10, 17, 37, 51	0
1	B	313/345 (90%)	-0.18	5 (1%) 72 77	12, 19, 38, 52	0
All	All	633/690 (91%)	-0.13	12 (1%) 67 72	10, 18, 38, 52	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	301	GLY	4.9
1	A	304	GLY	4.2
1	B	93	ALA	3.5
1	B	301	GLY	3.5
1	A	-1	GLY	3.3
1	B	303	LEU	2.9
1	B	94	GLY	2.5
1	A	0	SER	2.3
1	A	300	ASN	2.3
1	A	202	THR	2.2
1	B	144	GLN	2.0
1	A	302	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PO4	B	402	5/5	0.91	0.20	10.89	14,18,21,22	5
4	EDO	A	404	4/4	0.87	0.14	4.59	24,26,30,30	0
4	EDO	A	403	4/4	0.91	0.18	4.31	23,25,27,27	0
4	EDO	A	405	4/4	0.95	0.17	2.13	34,37,37,38	0
3	PO4	A	401	5/5	0.95	0.12	2.07	24,25,28,34	0
4	EDO	B	403	4/4	0.92	0.12	1.80	28,34,35,40	0
4	EDO	A	402	4/4	0.92	0.11	1.02	25,27,27,28	0
2	NAD	A	400	44/44	0.97	0.07	-0.76	13,14,16,17	0
3	PO4	B	401	5/5	0.85	0.20	-	18,21,21,22	5

6.5 Other polymers [i](#)

There are no such residues in this entry.