



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2017 – 04:13 PM EDT

PDB ID : 3I2C
Title : Crystal structure of anti-IL-23 antibody CNTO4088
Authors : Teplyakov, A.; Obmolova, G.; Gilliland, G.L.
Deposited on : unknown
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	218	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 27%, green 70%);"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; text-align: center;"> </div> </div> </div>
2	H	222	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background: linear-gradient(to right, green 77%, yellow 17%, orange 6%);"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; text-align: center;"> </div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CNTO4088 LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	218	Total	C	N	O	S	0	0	0
			1673	1048	283	337	5			

- Molecule 2 is a protein called CNTO4088 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1678	1065	281	323	9			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CNTO4088 LIGHT CHAIN



- Molecule 2: CNTO4088 HEAVY CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	99.43Å 99.43Å 115.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 15.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	94.9 (15.00-2.80) 99.9 (15.00-2.80)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.196 , 0.235 0.194 , 0.231	Depositor DCC
R_{free} test set	685 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	67.3	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.000 for -h,k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3359	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.40	0/1714	0.59	0/2330
2	H	0.41	0/1722	0.64	3/2351 (0.1%)
All	All	0.40	0/3436	0.62	3/4681 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	168	SER	N-CA-C	6.51	128.59	111.00
2	H	4	LEU	CA-CB-CG	5.58	128.14	115.30
2	H	180	ASP	CB-CG-OD2	5.20	122.98	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1673	0	1600	35	0
2	H	1678	0	1639	31	0
3	H	6	0	8	0	0
4	H	2	0	0	0	0
All	All	3359	0	3247	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:35:HIS:HD2	2:H:47:TRP:HE1	1.22	0.82
2:H:35:HIS:HE1	2:H:101:GLU:OE1	1.72	0.72
1:L:55:ALA:HB2	1:L:75:PHE:HE1	1.56	0.70
2:H:73:SER:HB3	2:H:82:TYR:HB2	1.74	0.70
2:H:112:GLN:NE2	2:H:112:GLN:H	1.90	0.70
2:H:35:HIS:CD2	2:H:47:TRP:HE1	2.08	0.70
1:L:119:VAL:HA	1:L:139:PHE:O	1.93	0.69
2:H:6:GLN:NE2	2:H:96:TYR:O	2.29	0.65
2:H:219:PRO:O	2:H:220:ARG:HB2	1.96	0.64
1:L:189:GLU:HA	1:L:192:ARG:HG2	1.80	0.63
1:L:199:GLU:HG3	1:L:210:VAL:HG22	1.83	0.60
1:L:13:VAL:HG21	1:L:82:VAL:HG21	1.82	0.60
1:L:41:GLN:HB2	1:L:51:LEU:HD11	1.84	0.58
1:L:55:ALA:HB2	1:L:75:PHE:CE1	2.38	0.58
1:L:40:TYR:HE2	1:L:93:GLN:HE21	1.50	0.58
1:L:112:ARG:HH21	1:L:115:ALA:HB2	1.69	0.57
1:L:35:SER:HA	1:L:96:GLY:HA3	1.87	0.57
1:L:151:LYS:HE2	1:L:199:GLU:HB3	1.90	0.54
1:L:65:ARG:HB3	1:L:81:PRO:HD2	1.90	0.53
2:H:112:GLN:HE21	2:H:112:GLN:H	1.56	0.53
2:H:196:PRO:HB3	2:H:219:PRO:HG3	1.90	0.53
1:L:17:GLN:HE21	1:L:17:GLN:HA	1.75	0.51
2:H:6:GLN:H	2:H:112:GLN:HE22	1.59	0.50
2:H:23:ALA:HA	2:H:80:LEU:HD23	1.94	0.50
1:L:194:ASN:ND2	1:L:214:ASN:OD1	2.46	0.49
1:L:42:GLN:NE2	1:L:48:PRO:HD3	2.27	0.49
2:H:219:PRO:O	2:H:220:ARG:CB	2.61	0.49
2:H:28:ASN:HD22	2:H:31:THR:HG23	1.78	0.48
2:H:22:CYS:HB3	2:H:81:LEU:HB3	1.96	0.48
1:L:40:TYR:OH	1:L:93:GLN:NE2	2.46	0.47
2:H:35:HIS:HD2	2:H:47:TRP:NE1	2.00	0.47
2:H:178:GLN:O	2:H:180:ASP:N	2.47	0.47
2:H:37:VAL:HG23	2:H:99:MET:HE1	1.96	0.47
1:L:141:ASN:HB3	1:L:142:ASN:HD22	1.81	0.46
1:L:167:TRP:CE2	1:L:179:MET:HG3	2.50	0.46
1:L:42:GLN:HE22	2:H:39:GLN:HE22	1.62	0.46
2:H:64:ASP:N	2:H:65:PRO:HD2	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:62:TYR:HE2	2:H:72:ILE:HG13	1.80	0.46
2:H:38:ARG:HB3	2:H:48:ILE:HD11	1.98	0.46
1:L:217:GLU:HB2	2:H:135:SER:HB3	1.98	0.45
2:H:103:ILE:HG13	2:H:104:TYR:CD1	2.52	0.45
1:L:174:ASP:C	1:L:174:ASP:OD1	2.55	0.45
1:L:153:LYS:HB2	1:L:197:THR:HB	1.99	0.45
2:H:28:ASN:ND2	2:H:31:THR:HG23	2.32	0.45
2:H:99:MET:HG3	2:H:107:PHE:HB3	1.98	0.45
1:L:38:HIS:ND1	2:H:106:SER:HB3	2.32	0.44
1:L:128:GLN:HG2	1:L:133:GLY:O	2.18	0.43
1:L:197:THR:HG23	1:L:210:VAL:HG13	2.01	0.43
1:L:122:PHE:HA	1:L:123:PRO:HD3	1.90	0.43
1:L:43:LYS:HB3	1:L:44:PRO:HD2	2.01	0.43
1:L:94:HIS:HE1	1:L:97:GLU:O	2.02	0.43
1:L:2:ILE:HD11	1:L:28:SER:HB2	2.02	0.42
1:L:38:HIS:CE1	2:H:106:SER:HB3	2.54	0.42
2:H:36:TRP:CE2	2:H:83:LEU:HB2	2.54	0.42
1:L:119:VAL:HG22	1:L:211:LYS:HG2	2.02	0.41
1:L:112:ARG:NH2	1:L:115:ALA:HB2	2.33	0.41
1:L:38:HIS:CE1	1:L:54:LEU:H	2.38	0.41
2:H:130:PRO:HD3	2:H:215:LYS:HE2	2.01	0.41
1:L:36:PHE:HB3	1:L:54:LEU:HD13	2.03	0.41
2:H:69:ARG:C	2:H:70:PHE:HD1	2.25	0.41
1:L:125:SER:HB2	1:L:127:GLU:HG3	2.04	0.40
2:H:37:VAL:HG23	2:H:99:MET:CE	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	216/218 (99%)	196 (91%)	17 (8%)	3 (1%)	13	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	220/222 (99%)	200 (91%)	15 (7%)	5 (2%)	7	25
All	All	436/440 (99%)	396 (91%)	32 (7%)	8 (2%)	10	32

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	44	PRO
2	H	220	ARG
2	H	178	GLN
2	H	179	SER
1	L	32	SER
1	L	72	GLY
2	H	103	ILE
2	H	65	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	188/190 (99%)	167 (89%)	21 (11%)	7	21
2	H	185/188 (98%)	167 (90%)	18 (10%)	9	27
All	All	373/378 (99%)	334 (90%)	39 (10%)	8	23

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	10	SER
1	L	17	GLN
1	L	27	LYS
1	L	65	ARG
1	L	93	GLN
1	L	94	HIS
1	L	95	SER
1	L	104	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	108	LEU
1	L	109	GLU
1	L	112	ARG
1	L	120	SER
1	L	127	GLU
1	L	130	THR
1	L	140	LEU
1	L	157	SER
1	L	185	LEU
1	L	191	GLU
1	L	207	SER
1	L	212	SER
1	L	217	GLU
2	H	1	GLU
2	H	4	LEU
2	H	6	GLN
2	H	11	LEU
2	H	37	VAL
2	H	38	ARG
2	H	60	THR
2	H	73	SER
2	H	81	LEU
2	H	83	LEU
2	H	85	MET
2	H	99	MET
2	H	112	GLN
2	H	123	THR
2	H	139	THR
2	H	160	THR
2	H	199	THR
2	H	221	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	17	GLN
1	L	42	GLN
1	L	46	GLN
1	L	93	GLN
1	L	94	HIS
1	L	142	ASN
1	L	149	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	193	HIS
2	H	28	ASN
2	H	35	HIS
2	H	78	GLN
2	H	86	ASN
2	H	112	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	H	1001	-	5,5,5	0.32	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	H	1001	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	218/218 (100%)	-0.51	3 (1%) 75 69	37, 59, 94, 114	0
2	H	222/222 (100%)	-0.67	0 100 100	37, 55, 76, 96	0
All	All	440/440 (100%)	-0.59	3 (0%) 87 83	37, 56, 87, 114	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	31	SER	3.0
1	L	32	SER	2.9
1	L	30	SER	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	H	1001	6/6	0.88	0.16	1.02	53,64,70,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.