



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 26, 2017 – 03:41 PM EDT

PDB ID : 3I5C  
Title : Crystal structure of a fusion protein containing the leucine zipper of GCN4 and the GGDEF domain of WspR from *Pseudomonas aeruginosa*  
Authors : Navarro, M.V.A.S.; De, N.; Sondermann, H.  
Deposited on : unknown  
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3233 (1.96-1.92)
Clashscore	112137	3430 (1.96-1.92)
Ramachandran outliers	110173	3395 (1.96-1.92)
Sidechain outliers	110143	3395 (1.96-1.92)
RSRZ outliers	101464	3250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• 5%</div> </div> </div>
1	B	206	<div> <div>11%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• •</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fusion of General control protein GCN4 and WSPR response regulator protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1533	947	284	295	7			
1	B	198	Total	C	N	O	S	0	2	0
			1543	955	286	296	6			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (three-letter code: C2E) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>10</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 46	C 20	N 10	O 14	P 2	0	0
3	A	1	Total 46	C 20	N 10	O 14	P 2	0	0
3	B	1	Total 46	C 20	N 10	O 14	P 2	0	0
3	B	1	Total 46	C 20	N 10	O 14	P 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	191	Total 191	O 191	0	0
4	B	101	Total 101	O 101	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fusion of General control protein GCN4 and WSPR response regulator protein



- Molecule 1: Fusion of General control protein GCN4 and WSPR response regulator protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.81Å 92.81Å 133.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.87 – 1.94 46.83 – 1.94	Depositor EDS
% Data completeness (in resolution range)	96.9 (31.87-1.94) 96.9 (46.83-1.94)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.18 (at 1.94Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.220 , 0.259 0.214 , 0.251	Depositor DCC
$R_{free}$ test set	3117 reflections (7.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.9	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3553	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/1555	0.54	0/2090
1	B	0.31	0/1571	0.48	0/2114
All	All	0.33	0/3126	0.51	0/4204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1533	0	1522	28	0
1	B	1543	0	1520	32	0
2	A	1	0	0	0	0
3	A	92	0	44	1	0
3	B	92	0	44	0	0
4	A	191	0	0	7	0
4	B	101	0	0	5	0
All	All	3553	0	3130	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:HIS:NE2	4:A:443:HOH:O	2.14	0.79
1:B:311:THR:O	1:B:314:VAL:HG12	1.91	0.71
1:B:152:GLU:HG2	1:B:156:LYS:HE2	1.73	0.70
1:B:311:THR:H	1:B:314:VAL:HG12	1.56	0.69
1:A:204:SER:OG	1:A:302:THR:HB	1.93	0.68
4:A:378:HOH:O	1:B:163:GLU:HG2	1.97	0.65
1:A:268:ARG:HB2	1:A:336:LEU:HD21	1.77	0.64
1:B:302:THR:HG23	4:B:47:HOH:O	2.01	0.60
1:A:336:LEU:O	1:A:337:MET:HB2	2.02	0.59
1:B:253:GLU:O	1:B:253:GLU:HG2	2.01	0.59
1:B:217:ASN:HD22	1:B:222:HIS:HA	1.68	0.59
1:A:195:ARG:HD3	4:A:423:HOH:O	2.01	0.58
1:B:217:ASN:ND2	1:B:222:HIS:HA	2.19	0.57
1:B:215:SER:O	1:B:219:THR:HG23	2.07	0.55
1:A:217:ASN:OD1	1:A:225:GLY:HA3	2.07	0.54
1:A:302:THR:HG23	4:A:20:HOH:O	2.07	0.53
1:A:311:THR:O	1:A:314:VAL:HG12	2.08	0.53
1:A:317:GLU:O	1:A:321:GLN:HG3	2.09	0.52
1:B:195:ARG:HD3	4:B:93:HOH:O	2.09	0.52
1:A:192:GLU:OE1	1:B:195:ARG:NH2	2.41	0.51
1:A:180:SER:OG	1:A:184:HIS:HD2	1.92	0.51
1:B:209:ASP:OD1	1:B:254:GLU:HG2	2.10	0.51
1:B:240:CYS:HB2	1:B:245:ASP:HB3	1.93	0.50
1:B:204:SER:OG	1:B:302:THR:HB	2.12	0.49
1:B:283:ILE:O	1:B:293:HIS:HA	2.12	0.49
1:A:238:GLU:O	1:A:273:LYS:NZ	2.45	0.49
1:B:272:GLU:HB3	4:B:37:HOH:O	2.13	0.48
1:A:336:LEU:O	1:A:337:MET:CB	2.61	0.47
1:B:180:SER:OG	1:B:184:HIS:HD2	1.98	0.47
1:B:264:PRO:HA	1:B:302:THR:HG21	1.97	0.47
1:A:145:GLN:OE1	1:A:145:GLN:HA	2.15	0.46
1:A:254:GLU:HG3	4:A:132:HOH:O	2.15	0.46
1:B:240:CYS:HB2	1:B:245:ASP:CB	2.45	0.46
1:A:177:THR:HB	1:A:233:ALA:CB	2.45	0.46
1:A:302:THR:CG2	4:A:20:HOH:O	2.62	0.46
1:B:286:ASP:OD2	1:B:287:GLN:HG2	2.16	0.46
1:A:191:MET:CE	1:A:194:ARG:HH21	2.29	0.46
1:B:227:GLU:HB2	1:B:230:ARG:HH21	1.81	0.46
1:A:240:CYS:HB2	1:A:245:ASP:CB	2.46	0.45
1:A:240:CYS:HB2	1:A:245:ASP:HB3	1.98	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:GLU:HG2	1:A:191:MET:CE	2.46	0.45
1:B:302:THR:CG2	4:B:47:HOH:O	2.62	0.45
1:A:313:ARG:HG2	4:A:128:HOH:O	2.17	0.45
1:A:283:ILE:HB	1:A:294:LEU:HB2	1.99	0.44
1:B:191:MET:CE	4:B:74:HOH:O	2.66	0.44
1:A:329:ASN:HD22	1:A:329:ASN:N	2.15	0.44
1:B:311:THR:H	1:B:314:VAL:CG1	2.26	0.43
1:A:257:MET:HB2	1:A:257:MET:HE2	1.82	0.43
1:B:163:GLU:HA	1:B:163:GLU:OE1	2.18	0.43
1:A:147:GLU:HG3	1:B:146:LEU:HD11	2.02	0.42
1:B:240:CYS:HB3	1:B:257:MET:HE3	2.01	0.42
1:B:183:ARG:HH11	1:B:183:ARG:HG2	1.85	0.42
1:B:268:ARG:HG2	1:B:336:LEU:HD21	2.01	0.42
1:A:163:GLU:OE2	1:A:163:GLU:HA	2.20	0.41
1:A:194:ARG:HD2	3:A:3:C2E:O2P	2.21	0.41
1:B:202:GLN:NE2	1:B:264:PRO:HD3	2.36	0.41
1:B:144:LYS:HE3	1:B:148:ASP:OD1	2.20	0.40
1:B:182:ARG:NH1	1:B:250:TYR:CZ	2.90	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/206 (94%)	192 (99%)	2 (1%)	0	100	100
1	B	198/206 (96%)	194 (98%)	4 (2%)	0	100	100
All	All	392/412 (95%)	386 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/171 (96%)	159 (97%)	5 (3%)	46	32
1	B	162/171 (95%)	157 (97%)	5 (3%)	45	31
All	All	326/342 (95%)	316 (97%)	10 (3%)	45	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	LYS
1	A	154	LEU
1	A	237	ARG
1	A	294	LEU
1	A	302	THR
1	B	163	GLU
1	B	176	LEU
1	B	253	GLU
1	B	287	GLN
1	B	302	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	184	HIS
1	A	287	GLN
1	A	329	ASN
1	B	184	HIS
1	B	202	GLN
1	B	217	ASN
1	B	231	GLN
1	B	287	GLN
1	B	329	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	C2E	A	1	-	44,52,52	0.92	3 (6%)	50,82,82	1.91	13 (26%)
3	C2E	A	3	-	44,52,52	0.89	1 (2%)	50,82,82	1.92	12 (24%)
3	C2E	B	2	-	44,52,52	0.91	0	50,82,82	1.96	12 (24%)
3	C2E	B	4	-	44,52,52	0.89	1 (2%)	50,82,82	1.88	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C2E	A	1	-	-	0/22/62/62	0/6/7/7
3	C2E	A	3	-	-	0/22/62/62	0/6/7/7
3	C2E	B	2	-	-	0/22/62/62	0/6/7/7
3	C2E	B	4	-	-	0/22/62/62	0/6/7/7

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	C2E	C81-N71	-2.29	1.30	1.34
3	A	3	C2E	C8-N7	-2.25	1.30	1.34
3	A	1	C2E	C8-N7	-2.22	1.30	1.34
3	B	4	C2E	C8-N7	-2.17	1.30	1.34
3	A	1	C2E	O4A-C1A	2.04	1.44	1.41

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	C2E	C51-C61-N11	-4.56	117.00	123.48
3	A	3	C2E	N31-C21-N11	-4.27	121.22	127.46
3	B	2	C2E	N31-C21-N11	-4.19	121.33	127.46
3	B	4	C2E	N3-C2-N1	-4.13	121.43	127.46
3	B	4	C2E	C51-C61-N11	-4.03	117.74	123.48
3	B	4	C2E	N31-C21-N11	-3.93	121.72	127.46
3	A	3	C2E	C51-C61-N11	-3.91	117.91	123.48
3	B	2	C2E	N3-C2-N1	-3.91	121.75	127.46
3	A	1	C2E	N3-C2-N1	-3.91	121.75	127.46
3	B	2	C2E	C51-C61-N11	-3.81	118.06	123.48
3	A	3	C2E	N3-C2-N1	-3.73	122.01	127.46
3	A	3	C2E	C5-C6-N1	-3.68	118.24	123.48
3	A	1	C2E	N31-C21-N11	-3.66	122.11	127.46
3	B	2	C2E	C61-C51-C41	-3.65	117.22	120.84
3	A	1	C2E	C5-C6-N1	-3.62	118.33	123.48
3	A	1	C2E	C6-C5-C4	-3.58	117.28	120.84
3	B	4	C2E	C5-C6-N1	-3.39	118.66	123.48
3	B	2	C2E	C5-C6-N1	-3.35	118.71	123.48
3	B	2	C2E	C6-C5-C4	-3.05	117.81	120.84
3	A	3	C2E	C61-C51-C41	-3.03	117.83	120.84
3	B	4	C2E	C6-C5-C4	-2.88	117.98	120.84
3	A	1	C2E	C41-C51-N71	-2.83	106.68	109.41
3	B	4	C2E	C61-C51-C41	-2.61	118.25	120.84
3	A	3	C2E	C6-C5-C4	-2.55	118.31	120.84
3	A	3	C2E	C4-C5-N7	-2.48	107.02	109.41
3	B	2	C2E	C41-C51-N71	-2.38	107.11	109.41
3	A	1	C2E	C61-C51-C41	-2.37	118.49	120.84
3	A	3	C2E	C41-C51-N71	-2.35	107.14	109.41
3	B	4	C2E	C4-C5-N7	-2.29	107.20	109.41
3	B	4	C2E	C41-C51-N71	-2.16	107.32	109.41
3	A	1	C2E	C1'-N9-C4	-2.04	123.12	126.64
3	A	1	C2E	C4A-O4A-C1A	-2.03	107.61	109.77
3	B	2	C2E	N2-C2-N1	2.08	120.56	117.24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	C2E	C6-N1-C2	2.78	120.05	116.06
3	B	4	C2E	C6-N1-C2	3.05	120.44	116.06
3	A	3	C2E	C6-N1-C2	3.16	120.60	116.06
3	A	1	C2E	C6-N1-C2	3.56	121.18	116.06
3	B	4	C2E	C61-N11-C21	3.77	121.48	116.06
3	A	1	C2E	C21-N31-C41	3.82	119.62	115.16
3	B	2	C2E	C61-N11-C21	3.89	121.66	116.06
3	A	3	C2E	C61-N11-C21	4.00	121.81	116.06
3	A	1	C2E	C2-N3-C4	4.04	119.87	115.16
3	A	1	C2E	C61-N11-C21	4.07	121.91	116.06
3	B	4	C2E	C21-N31-C41	4.34	120.22	115.16
3	A	3	C2E	C21-N31-C41	4.80	120.77	115.16
3	B	2	C2E	C21-N31-C41	4.84	120.81	115.16
3	A	3	C2E	C2-N3-C4	4.98	120.97	115.16
3	B	2	C2E	C2-N3-C4	5.03	121.04	115.16
3	B	4	C2E	C2-N3-C4	5.35	121.41	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3	C2E	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	196/206 (95%)	0.72	12 (6%) 22 29	20, 32, 55, 97	0
1	B	198/206 (96%)	0.77	22 (11%) 6 9	22, 41, 70, 86	0
All	All	394/412 (95%)	0.74	34 (8%) 11 16	20, 36, 65, 97	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	LYS	4.5
1	A	142	ARG	3.9
1	A	240	CYS	3.7
1	B	325	GLN	3.6
1	B	142	ARG	3.5
1	B	294	LEU	3.4
1	B	321	GLN	3.2
1	B	222	HIS	3.2
1	B	143	MET	3.2
1	B	292	SER	3.1
1	B	288	PRO	3.1
1	B	146	LEU	3.0
1	A	336	LEU	2.9
1	B	324	TYR	2.9
1	A	154	LEU	2.9
1	A	143	MET	2.7
1	B	220	PHE	2.7
1	A	239	GLY	2.7
1	B	224	ALA	2.6
1	A	146	LEU	2.5
1	B	290	PRO	2.4
1	A	222	HIS	2.4
1	B	145	GLN	2.4
1	B	322	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	219	THR	2.3
1	A	150	VAL	2.3
1	A	145	GLN	2.3
1	B	213	PHE	2.2
1	B	282	GLN	2.2
1	B	336	LEU	2.2
1	B	329	ASN	2.2
1	A	152	GLU	2.1
1	B	323	LEU	2.1
1	B	214	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	C2E	B	2	46/46	0.98	0.13	-0.23	18,21,24,26	0
3	C2E	B	4	46/46	0.96	0.13	-0.53	29,41,48,51	0
3	C2E	A	1	46/46	0.98	0.13	-0.63	18,23,26,27	0
3	C2E	A	3	46/46	0.97	0.13	-0.67	25,30,39,42	0
2	MG	A	400	1/1	0.95	0.18	-	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.