



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:05 am GMT

PDB ID : 4I5E  
Title : Crystal structure of Ralstonia sp. alcohol dehydrogenase in complex with NADP+  
Authors : Jarasch, A.; Lerchner, A.; Meining, W.; Schiefner, A.; Skerra, A.  
Deposited on : 2012-11-28  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	<div> <div>10%</div> <div> <div>83%</div> <div>10%</div> <div>5%</div> </div> </div>
1	B	262	<div> <div>19%</div> <div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
1	C	262	<div> <div>16%</div> <div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
1	D	262	<div> <div>11%</div> <div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>
1	E	262	<div> <div>11%</div> <div> <div>84%</div> <div>10%</div> <div>5%</div> </div> </div>
1	F	262	<div> <div>10%</div> <div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	262	
1	H	262	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	302	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15560 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase/short-chain dehydrogenase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	B	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	C	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	D	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	E	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	F	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	G	249	Total	C	N	O	0	0	0
			1882	1181	336	365			
1	H	249	Total	C	N	O	0	0	0
			1882	1181	336	365			

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	EXPRESSION TAG	UNP C0IR58
A	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
A	-10	SER	-	EXPRESSION TAG	UNP C0IR58
A	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
A	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
A	-7	SER	-	EXPRESSION TAG	UNP C0IR58
A	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
A	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
A	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
A	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
A	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
A	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
A	0	GLY	-	EXPRESSION TAG	UNP C0IR58

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	EXPRESSION TAG	UNP C0IR58
B	-12	MET	-	EXPRESSION TAG	UNP C0IR58
B	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
B	-10	SER	-	EXPRESSION TAG	UNP C0IR58
B	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
B	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
B	-7	SER	-	EXPRESSION TAG	UNP C0IR58
B	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
B	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
B	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
B	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
B	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
B	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
B	0	GLY	-	EXPRESSION TAG	UNP C0IR58
B	1	ALA	-	EXPRESSION TAG	UNP C0IR58
C	-12	MET	-	EXPRESSION TAG	UNP C0IR58
C	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
C	-10	SER	-	EXPRESSION TAG	UNP C0IR58
C	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
C	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
C	-7	SER	-	EXPRESSION TAG	UNP C0IR58
C	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
C	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
C	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
C	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
C	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
C	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
C	0	GLY	-	EXPRESSION TAG	UNP C0IR58
C	1	ALA	-	EXPRESSION TAG	UNP C0IR58
D	-12	MET	-	EXPRESSION TAG	UNP C0IR58
D	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
D	-10	SER	-	EXPRESSION TAG	UNP C0IR58
D	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
D	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
D	-7	SER	-	EXPRESSION TAG	UNP C0IR58
D	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
D	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
D	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
D	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
D	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
D	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
D	0	GLY	-	EXPRESSION TAG	UNP C0IR58

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
D	1	ALA	-	EXPRESSION TAG	UNP C0IR58
E	-12	MET	-	EXPRESSION TAG	UNP C0IR58
E	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
E	-10	SER	-	EXPRESSION TAG	UNP C0IR58
E	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
E	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
E	-7	SER	-	EXPRESSION TAG	UNP C0IR58
E	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
E	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
E	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
E	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
E	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
E	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
E	0	GLY	-	EXPRESSION TAG	UNP C0IR58
E	1	ALA	-	EXPRESSION TAG	UNP C0IR58
F	-12	MET	-	EXPRESSION TAG	UNP C0IR58
F	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
F	-10	SER	-	EXPRESSION TAG	UNP C0IR58
F	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
F	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
F	-7	SER	-	EXPRESSION TAG	UNP C0IR58
F	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
F	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
F	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
F	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
F	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
F	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
F	0	GLY	-	EXPRESSION TAG	UNP C0IR58
F	1	ALA	-	EXPRESSION TAG	UNP C0IR58
G	-12	MET	-	EXPRESSION TAG	UNP C0IR58
G	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
G	-10	SER	-	EXPRESSION TAG	UNP C0IR58
G	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
G	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
G	-7	SER	-	EXPRESSION TAG	UNP C0IR58
G	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
G	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
G	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
G	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
G	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
G	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
G	0	GLY	-	EXPRESSION TAG	UNP C0IR58

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	1	ALA	-	EXPRESSION TAG	UNP C0IR58
H	-12	MET	-	EXPRESSION TAG	UNP C0IR58
H	-11	ALA	-	EXPRESSION TAG	UNP C0IR58
H	-10	SER	-	EXPRESSION TAG	UNP C0IR58
H	-9	ARG	-	EXPRESSION TAG	UNP C0IR58
H	-8	GLY	-	EXPRESSION TAG	UNP C0IR58
H	-7	SER	-	EXPRESSION TAG	UNP C0IR58
H	-6	HIS	-	EXPRESSION TAG	UNP C0IR58
H	-5	HIS	-	EXPRESSION TAG	UNP C0IR58
H	-4	HIS	-	EXPRESSION TAG	UNP C0IR58
H	-3	HIS	-	EXPRESSION TAG	UNP C0IR58
H	-2	HIS	-	EXPRESSION TAG	UNP C0IR58
H	-1	HIS	-	EXPRESSION TAG	UNP C0IR58
H	0	GLY	-	EXPRESSION TAG	UNP C0IR58
H	1	ALA	-	EXPRESSION TAG	UNP C0IR58

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	H	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total 14	O 14	0	0
4	B	15	Total 15	O 15	0	0
4	C	10	Total 10	O 10	0	0
4	D	11	Total 11	O 11	0	0
4	E	12	Total 12	O 12	0	0
4	F	16	Total 16	O 16	0	0
4	G	8	Total 8	O 8	0	0
4	H	10	Total 10	O 10	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase




- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase



- Molecule 1: Alcohol dehydrogenase/short-chain dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.34Å 74.55Å 132.69Å 80.98° 85.99° 64.60°	Depositor
Resolution (Å)	29.60 – 2.80 29.56 – 2.80	Depositor EDS
% Data completeness (in resolution range)	93.7 (29.60-2.80) 89.3 (29.56-2.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.261 , 0.305 0.268 , 0.312	Depositor DCC
$R_{free}$ test set	2942 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.5	Xtriage
Anisotropy	0.266	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 54.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	15560	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.28 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.8141e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.69	4/1905 (0.2%)	0.96	7/2581 (0.3%)
1	B	0.54	0/1905	0.69	3/2581 (0.1%)
1	C	0.49	1/1905 (0.1%)	0.63	0/2581
1	D	0.51	2/1905 (0.1%)	0.64	1/2581 (0.0%)
1	E	0.57	2/1905 (0.1%)	0.95	4/2581 (0.2%)
1	F	0.56	2/1905 (0.1%)	0.67	1/2581 (0.0%)
1	G	0.52	2/1905 (0.1%)	0.64	1/2581 (0.0%)
1	H	0.53	1/1905 (0.1%)	0.65	1/2581 (0.0%)
All	All	0.55	14/15240 (0.1%)	0.74	18/20648 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127	ARG	CZ-NH1	-13.63	1.15	1.33
1	G	127	ARG	CZ-NH1	-8.06	1.22	1.33
1	H	127	ARG	CZ-NH1	-7.58	1.23	1.33
1	E	127	ARG	CZ-NH1	-7.57	1.23	1.33
1	F	127	ARG	CZ-NH1	-7.29	1.23	1.33

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	127	ARG	NE-CZ-NH2	23.25	131.93	120.30
1	E	127	ARG	NE-CZ-NH1	-23.18	108.71	120.30
1	A	127	ARG	NE-CZ-NH2	22.71	131.66	120.30
1	A	148	ASP	CB-CG-OD2	15.64	132.37	118.30
1	E	127	ARG	CD-NE-CZ	12.53	141.14	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	127	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1882	0	1922	21	0
1	B	1882	0	1922	25	0
1	C	1882	0	1922	23	0
1	D	1882	0	1922	22	0
1	E	1882	0	1922	22	0
1	F	1882	0	1922	27	2
1	G	1882	0	1922	23	2
1	H	1882	0	1922	25	0
2	A	48	0	25	1	0
2	B	48	0	25	5	0
2	C	48	0	25	1	0
2	D	48	0	25	4	0
2	E	48	0	25	4	0
2	F	48	0	25	7	0
2	G	48	0	25	6	0
2	H	48	0	25	6	0
3	A	6	0	8	1	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	E	6	0	8	0	0
4	A	14	0	0	0	0
4	B	15	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	10	0	0	0	0
4	D	11	0	0	2	0
4	E	12	0	0	1	0
4	F	16	0	0	0	0
4	G	8	0	0	0	0
4	H	10	0	0	1	0
All	All	15560	0	15608	173	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 173 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:183:ILE:HB	2:E:301:NAP:N7N	1.90	0.86
1:E:183:ILE:HB	2:E:301:NAP:H72N	1.41	0.83
1:G:183:ILE:HB	2:G:301:NAP:N7N	2.01	0.75
1:B:34:PHE:CD1	1:B:55:THR:HB	2.26	0.71
1:E:34:PHE:CD1	1:E:55:THR:HB	2.29	0.67

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:40:ARG:NH2	1:G:102:GLU:OE2[1_655]	1.91	0.29
1:F:102:GLU:OE2	1:G:40:ARG:NH2[1_655]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/262 (94%)	235 (95%)	10 (4%)	2 (1%)	22	55
1	B	247/262 (94%)	240 (97%)	6 (2%)	1 (0%)	38	72
1	C	247/262 (94%)	238 (96%)	9 (4%)	0	100	100
1	D	247/262 (94%)	238 (96%)	8 (3%)	1 (0%)	38	72
1	E	247/262 (94%)	240 (97%)	6 (2%)	1 (0%)	38	72
1	F	247/262 (94%)	238 (96%)	9 (4%)	0	100	100
1	G	247/262 (94%)	239 (97%)	7 (3%)	1 (0%)	38	72
1	H	247/262 (94%)	240 (97%)	6 (2%)	1 (0%)	38	72
All	All	1976/2096 (94%)	1908 (97%)	61 (3%)	7 (0%)	38	72

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	136	SER
1	A	136	SER
1	G	136	SER
1	H	136	SER
1	A	184	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/204 (95%)	191 (98%)	3 (2%)	70	92
1	B	194/204 (95%)	191 (98%)	3 (2%)	70	92
1	C	194/204 (95%)	192 (99%)	2 (1%)	80	95
1	D	194/204 (95%)	192 (99%)	2 (1%)	80	95
1	E	194/204 (95%)	192 (99%)	2 (1%)	80	95
1	F	194/204 (95%)	192 (99%)	2 (1%)	80	95
1	G	194/204 (95%)	192 (99%)	2 (1%)	80	95
1	H	194/204 (95%)	192 (99%)	2 (1%)	80	95

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1552/1632 (95%)	1534 (99%)	18 (1%)	75 94

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	148	ASP
1	D	158	ARG
1	G	148	ASP
1	C	148	ASP
1	C	158	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	87	ASN
1	H	87	ASN
1	F	87	ASN
1	D	87	ASN
1	G	87	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAP	A	301	-	44,52,52	0.98	2 (4%)	51,80,80	1.46	3 (5%)
3	GOL	A	302	-	5,5,5	0.37	0	5,5,5	0.31	0
2	NAP	B	301	-	44,52,52	1.19	4 (9%)	51,80,80	1.64	8 (15%)
3	GOL	B	302	-	5,5,5	0.33	0	5,5,5	0.36	0
2	NAP	C	301	-	44,52,52	0.90	1 (2%)	51,80,80	1.59	6 (11%)
3	GOL	C	302	-	5,5,5	0.25	0	5,5,5	0.33	0
2	NAP	D	301	-	44,52,52	1.14	4 (9%)	51,80,80	1.59	8 (15%)
2	NAP	E	301	-	44,52,52	1.13	4 (9%)	51,80,80	1.50	6 (11%)
3	GOL	E	302	-	5,5,5	0.59	0	5,5,5	0.55	0
2	NAP	F	301	-	44,52,52	1.20	5 (11%)	51,80,80	1.86	9 (17%)
2	NAP	G	301	-	44,52,52	0.88	2 (4%)	51,80,80	1.81	6 (11%)
2	NAP	H	301	-	44,52,52	0.95	2 (4%)	51,80,80	1.49	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	301	-	-	0/27/67/67	0/5/5/5
3	GOL	A	302	-	-	0/4/4/4	0/0/0/0
2	NAP	B	301	-	-	0/27/67/67	0/5/5/5
3	GOL	B	302	-	-	0/4/4/4	0/0/0/0
2	NAP	C	301	-	-	0/27/67/67	0/5/5/5
3	GOL	C	302	-	-	0/4/4/4	0/0/0/0
2	NAP	D	301	-	-	1/27/67/67	0/5/5/5
2	NAP	E	301	-	-	0/27/67/67	0/5/5/5
3	GOL	E	302	-	-	0/4/4/4	0/0/0/0
2	NAP	F	301	-	-	0/27/67/67	0/5/5/5
2	NAP	G	301	-	-	0/27/67/67	0/5/5/5
2	NAP	H	301	-	-	0/27/67/67	0/5/5/5

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	NAP	C2A-N3A	2.04	1.35	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	301	NAP	P2B-O2B	2.07	1.63	1.59
2	F	301	NAP	C2A-N3A	2.08	1.35	1.32
2	G	301	NAP	O4D-C1D	2.10	1.44	1.41
2	E	301	NAP	C2D-C1D	2.12	1.57	1.53

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	301	NAP	N3A-C2A-N1A	-7.47	122.36	128.86
2	H	301	NAP	N3A-C2A-N1A	-7.30	122.50	128.86
2	F	301	NAP	N3A-C2A-N1A	-7.13	122.64	128.86
2	B	301	NAP	N3A-C2A-N1A	-6.81	122.93	128.86
2	A	301	NAP	N3A-C2A-N1A	-6.74	122.98	128.86

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	NAP	P2B-O2B-C2B-C1B

There are no ring outliers.

9 monomers are involved in 35 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAP	1	0
3	A	302	GOL	1	0
2	B	301	NAP	5	0
2	C	301	NAP	1	0
2	D	301	NAP	4	0
2	E	301	NAP	4	0
2	F	301	NAP	7	0
2	G	301	NAP	6	0
2	H	301	NAP	6	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	249/262 (95%)	0.82	27 (10%)	6 3	25, 53, 104, 145	7 (2%)
1	B	249/262 (95%)	1.08	50 (20%)	1 1	34, 55, 101, 163	11 (4%)
1	C	249/262 (95%)	0.99	41 (16%)	2 1	31, 62, 108, 165	8 (3%)
1	D	249/262 (95%)	0.87	28 (11%)	6 3	31, 54, 98, 151	7 (2%)
1	E	249/262 (95%)	0.85	29 (11%)	5 3	28, 47, 105, 176	7 (2%)
1	F	249/262 (95%)	0.73	26 (10%)	7 4	25, 48, 99, 147	7 (2%)
1	G	249/262 (95%)	1.08	49 (19%)	1 1	25, 58, 109, 139	8 (3%)
1	H	249/262 (95%)	0.71	26 (10%)	7 4	29, 51, 99, 146	6 (2%)
All	All	1992/2096 (95%)	0.89	276 (13%)	3 2	25, 53, 105, 176	61 (3%)

The worst 5 of 276 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	191	GLN	10.1
1	C	191	GLN	9.9
1	G	193	SER	9.6
1	E	198	ALA	7.9
1	C	1	ALA	7.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	302	6/6	0.82	0.43	1.12	68,76,84,96	0
3	GOL	E	302	6/6	0.79	0.33	0.63	46,61,72,72	0
3	GOL	C	302	6/6	0.89	0.26	-0.20	63,72,74,78	0
3	GOL	A	302	6/6	0.81	0.24	-0.29	61,61,63,64	0
2	NAP	H	301	48/48	0.91	0.21	-0.34	47,73,98,109	0
2	NAP	A	301	48/48	0.85	0.23	-0.43	41,63,74,84	0
2	NAP	E	301	48/48	0.88	0.21	-0.50	43,56,66,68	0
2	NAP	G	301	48/48	0.86	0.23	-0.57	47,66,79,79	0
2	NAP	F	301	48/48	0.90	0.21	-0.60	34,48,61,74	0
2	NAP	D	301	48/48	0.88	0.22	-0.68	54,71,107,132	0
2	NAP	C	301	48/48	0.89	0.20	-0.78	55,69,78,85	0
2	NAP	B	301	48/48	0.91	0.20	-0.92	37,56,76,87	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.