



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:44 pm GMT

PDB ID : 2I67
Title : Structural Basis for the Mechanistic Understanding Human CD38 Controlled Multiple Catalysis
Authors : Liu, Q.; Kriksunov, I.A.; Graeff, R.; Munshi, C.; Lee, H.C.; Hao, Q.
Deposited on : 2006-08-28
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4385 (1.74-1.70)
Clashscore	112137	4841 (1.74-1.70)
Ramachandran outliers	110173	4771 (1.74-1.70)
Sidechain outliers	110143	4771 (1.74-1.70)
RSRZ outliers	101464	4426 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	<div> <div>13%</div> <div> <div></div> <div>73%</div> <div>22%</div> <div>• •</div> </div> </div>
1	B	262	<div> <div>9%</div> <div> <div></div> <div>71%</div> <div>21%</div> <div>• 5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	APR	A	301	X	-	-	-
2	APR	B	301	X	-	-	X
2	APR	B	302	X	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ADP-ribosyl cyclase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	252	Total	C	N	O	S	0	0	0
			2008	1266	351	375	16			
1	B	248	Total	C	N	O	S	0	0	0
			1987	1255	347	370	15			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	LYS	-	CLONING ARTIFACT	UNP P28907
A	40	ARG	-	CLONING ARTIFACT	UNP P28907
A	41	GLU	-	CLONING ARTIFACT	UNP P28907
A	42	ALA	-	CLONING ARTIFACT	UNP P28907
A	43	GLU	-	CLONING ARTIFACT	UNP P28907
A	44	ALA	-	CLONING ARTIFACT	UNP P28907
A	49	THR	GLN	ENGINEERED	UNP P28907
A	100	ASP	ASN	ENGINEERED	UNP P28907
A	164	ASP	ASN	ENGINEERED	UNP P28907
A	209	ASP	ASN	ENGINEERED	UNP P28907
A	219	ASP	ASN	ENGINEERED	UNP P28907
B	39	LYS	-	CLONING ARTIFACT	UNP P28907
B	40	ARG	-	CLONING ARTIFACT	UNP P28907
B	41	GLU	-	CLONING ARTIFACT	UNP P28907
B	42	ALA	-	CLONING ARTIFACT	UNP P28907
B	43	GLU	-	CLONING ARTIFACT	UNP P28907
B	44	ALA	-	CLONING ARTIFACT	UNP P28907
B	49	THR	GLN	ENGINEERED	UNP P28907
B	100	ASP	ASN	ENGINEERED	UNP P28907
B	164	ASP	ASN	ENGINEERED	UNP P28907
B	209	ASP	ASN	ENGINEERED	UNP P28907
B	219	ASP	ASN	ENGINEERED	UNP P28907

- Molecule 2 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula: $C_{15}H_{23}N_5O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	258	Total	O	0	0
			258	258		
3	B	213	Total	O	0	0
			213	213		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADP-ribosyl cyclase 1



• Molecule 1: ADP-ribosyl cyclase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.89Å 51.15Å 65.64Å 109.36° 91.22° 97.38°	Depositor
Resolution (Å)	20.00 – 1.71 46.03 – 1.71	Depositor EDS
% Data completeness (in resolution range)	94.4 (20.00-1.71) 84.8 (46.03-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.62 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.189 , 0.236 0.236 , 0.279	Depositor DCC
R_{free} test set	2662 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 61.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4574	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	2/2058 (0.1%)	0.77	0/2785
1	B	0.69	0/2037	0.70	0/2757
All	All	0.75	2/4095 (0.0%)	0.73	0/5542

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	72	GLU	CD-OE2	6.94	1.33	1.25
1	A	72	GLU	CB-CG	5.87	1.63	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	0	1917	42	0
1	B	1987	0	1906	50	0
2	A	36	0	21	2	0
2	B	72	0	42	0	0
3	A	258	0	0	9	0
3	B	213	0	0	7	0
All	All	4574	0	3886	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:THR:HG23	1:A:167:LYS:H	1.27	0.98
1:B:266:ILE:HD11	1:B:273:PHE:CD1	2.06	0.90
1:B:75:PRO:HA	1:B:78:ARG:HG3	1.57	0.87
1:A:203:VAL:HG21	1:B:171:GLN:HG2	1.60	0.84
1:B:115:GLN:HE22	1:B:149:LEU:H	1.24	0.83
1:A:231:GLN:HB2	1:A:234:LYS:HD3	1.60	0.82
1:A:115:GLN:HE22	1:A:149:LEU:H	1.28	0.81
1:A:45:ARG:HH12	1:A:47:ARG:HB2	1.46	0.80
1:A:90:LYS:HG2	1:A:94:ILE:HG13	1.64	0.79
1:A:142:MET:HE3	1:A:241:TRP:HH2	1.48	0.78
1:A:111:LYS:HE2	3:A:548:HOH:O	1.89	0.71
1:B:176:TRP:CZ3	1:B:181:SER:HB2	2.26	0.70
1:B:262:LEU:HD22	1:B:266:ILE:HD13	1.75	0.69
1:B:266:ILE:CD1	1:B:273:PHE:CD1	2.75	0.69
1:A:45:ARG:NH1	1:A:47:ARG:HB2	2.08	0.67
1:B:279:TYR:O	1:B:281:PRO:HD3	1.95	0.67
1:A:203:VAL:CG2	1:B:171:GLN:HG2	2.26	0.65
1:A:111:LYS:HB3	3:A:542:HOH:O	1.97	0.64
1:B:221:THR:HG23	1:B:225:VAL:CG1	2.28	0.64
1:A:244:HIS:HE1	1:A:277:ASN:OD1	1.80	0.63
1:A:106:TYR:O	1:A:110:MET:HG2	1.98	0.63
1:B:110:MET:SD	1:B:191:THR:HG22	2.39	0.62
1:A:190:LYS:HE2	3:A:556:HOH:O	1.99	0.62
1:A:142:MET:CE	1:A:241:TRP:HH2	2.13	0.61
1:B:221:THR:HG23	1:B:225:VAL:HG11	1.81	0.61
1:A:142:MET:HE3	1:A:241:TRP:CH2	2.33	0.60
1:B:171:GLN:NE2	3:B:500:HOH:O	2.34	0.60
1:A:183:ASN:ND2	1:A:186:SER:H	2.00	0.60
1:B:110:MET:HE1	1:B:192:VAL:HG12	1.83	0.59
1:A:129:LYS:NZ	1:A:156:ASP:OD1	2.34	0.59
1:A:183:ASN:HD21	1:A:186:SER:H	1.50	0.59
1:B:103:GLU:OE2	1:B:194:ARG:NH1	2.35	0.59
1:B:103:GLU:OE1	1:B:195:ARG:NH2	2.37	0.58
1:B:284:PHE:O	1:B:288:VAL:HG23	2.05	0.57
1:B:103:GLU:CD	1:B:194:ARG:HH12	2.08	0.57
1:A:228:HIS:HE1	3:A:334:HOH:O	1.86	0.56
1:A:75:PRO:O	1:A:78:ARG:HG2	2.04	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:HIS:CE1	1:A:146:GLU:HB2	2.41	0.56
1:B:106:TYR:O	1:B:110:MET:HG2	2.06	0.55
1:A:142:MET:CE	1:A:241:TRP:CH2	2.90	0.55
1:A:142:MET:HE2	3:B:511:HOH:O	2.07	0.54
1:B:183:ASN:ND2	1:B:186:SER:H	2.05	0.54
1:A:176:TRP:CZ3	1:A:181:SER:HB2	2.42	0.54
1:B:250:SER:CA	3:B:509:HOH:O	2.55	0.53
1:A:212:ARG:O	1:A:213:SER:CB	2.58	0.52
1:B:177:ARG:HG2	3:B:373:HOH:O	2.10	0.52
1:B:212:ARG:O	1:B:214:LYS:N	2.43	0.52
1:B:131:LEU:O	1:B:131:LEU:HD23	2.10	0.51
1:B:209:ASP:OD1	1:B:211:SER:HB2	2.11	0.51
1:B:45:ARG:NH2	3:B:362:HOH:O	2.43	0.51
1:B:221:THR:O	1:B:225:VAL:CG1	2.59	0.51
1:B:103:GLU:HG3	1:B:191:THR:HG23	1.93	0.51
1:B:236:GLN:HG3	3:B:442:HOH:O	2.10	0.50
1:B:90:LYS:HG2	1:B:94:ILE:HG13	1.92	0.50
1:A:90:LYS:HG2	1:A:94:ILE:CG1	2.36	0.49
1:B:90:LYS:CG	1:B:94:ILE:HG13	2.42	0.49
1:B:133:HIS:HE1	1:B:146:GLU:OE1	1.95	0.49
1:A:48:GLN:OE1	1:A:171:GLN:HB3	2.14	0.47
1:A:115:GLN:NE2	1:A:149:LEU:H	2.06	0.47
2:A:301:APR:C6	2:A:301:APR:HR'4	2.44	0.47
1:B:221:THR:O	1:B:225:VAL:HG13	2.14	0.47
1:A:138:VAL:HG11	1:A:289:LYS:HG2	1.97	0.47
1:B:46:TRP:CD1	1:B:47:ARG:HG3	2.50	0.47
1:B:246:GLY:O	1:B:248:GLU:N	2.41	0.46
1:B:45:ARG:HG3	1:B:46:TRP:H	1.81	0.46
2:A:301:APR:H'3	3:A:360:HOH:O	2.16	0.45
1:B:127:ARG:HD3	1:B:212:ARG:HD3	1.99	0.45
1:A:76:GLU:HG2	1:A:77:MET:HG2	1.98	0.45
1:A:74:HIS:HE1	3:A:414:HOH:O	2.00	0.44
1:B:50:TRP:CZ2	1:B:98:PRO:HG2	2.52	0.44
1:A:141:ASP:O	1:A:143:PHE:CE2	2.71	0.44
1:A:231:GLN:CB	1:A:234:LYS:HD3	2.40	0.43
1:B:104:GLU:HA	1:B:107:GLN:HG2	1.99	0.43
1:B:266:ILE:HG13	1:B:273:PHE:HB2	1.99	0.43
1:A:194:ARG:NH1	3:A:349:HOH:O	2.23	0.43
1:A:180:CYS:HB2	3:A:388:HOH:O	2.19	0.43
1:A:45:ARG:HH12	1:A:47:ARG:CB	2.23	0.43
1:B:69:LYS:HA	1:B:72:GLU:HG2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:PHE:HB3	1:A:60:PRO:HD3	2.01	0.42
1:B:110:MET:SD	1:B:191:THR:CG2	3.08	0.42
1:A:133:HIS:HE1	1:A:146:GLU:OE1	2.02	0.42
1:A:69:LYS:HD3	1:A:152:TYR:OH	2.20	0.42
1:B:107:GLN:HA	1:B:110:MET:HG3	2.02	0.41
1:B:228:HIS:HD2	3:B:423:HOH:O	2.03	0.41
1:A:98:PRO:O	1:A:183:ASN:HA	2.20	0.41
1:B:45:ARG:NH2	1:B:47:ARG:HD3	2.35	0.41
1:B:107:GLN:HE22	1:B:195:ARG:NH2	2.19	0.41
1:B:260:LYS:HD3	1:B:260:LYS:HA	1.91	0.41
1:A:270:ASN:ND2	3:A:541:HOH:O	2.53	0.41
1:B:45:ARG:HG3	1:B:46:TRP:N	2.36	0.41
1:B:45:ARG:HH22	1:B:47:ARG:HH11	1.69	0.40
1:B:115:GLN:NE2	1:B:149:LEU:H	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/262 (95%)	238 (95%)	6 (2%)	6 (2%)	7	1
1	B	246/262 (94%)	229 (93%)	11 (4%)	6 (2%)	7	1
All	All	496/524 (95%)	467 (94%)	17 (3%)	12 (2%)	7	1

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	248	GLU
1	A	213	SER
1	B	246	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	ASP
1	B	247	ARG
1	A	247	ARG
1	B	213	SER
1	A	245	GLY
1	B	291	PRO
1	B	290	ASN
1	B	128	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/241 (92%)	214 (96%)	9 (4%)	36	14
1	B	221/241 (92%)	207 (94%)	14 (6%)	21	5
All	All	444/482 (92%)	421 (95%)	23 (5%)	27	8

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ARG
1	A	89	PHE
1	A	103	GLU
1	A	110	MET
1	A	117	VAL
1	A	130	ASP
1	A	131	LEU
1	A	164	ASP
1	A	178	LYS
1	B	45	ARG
1	B	89	PHE
1	B	110	MET
1	B	176	TRP
1	B	177	ARG
1	B	183	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	202	ASP
1	B	233	GLU
1	B	251	ARG
1	B	262	LEU
1	B	266	ILE
1	B	282	ASP
1	B	285	LEU
1	B	286	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	133	HIS
1	A	182	ASN
1	A	183	ASN
1	A	228	HIS
1	A	244	HIS
1	A	286	GLN
1	B	107	GLN
1	B	115	GLN
1	B	133	HIS
1	B	171	GLN
1	B	183	ASN
1	B	244	HIS
1	B	270	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	APR	A	301	-	34,39,39	1.00	2 (5%)	36,60,60	2.40	11 (30%)
2	APR	B	301	-	34,39,39	1.00	1 (2%)	36,60,60	1.51	3 (8%)
2	APR	B	302	-	34,39,39	1.00	2 (5%)	36,60,60	1.57	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APR	A	301	-	2/2/10/10	0/18/54/54	0/4/4/4
2	APR	B	301	-	2/2/10/10	0/18/54/54	0/4/4/4
2	APR	B	302	-	2/2/10/10	0/18/54/54	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	APR	O4'-C1'	2.01	1.44	1.41
2	A	301	APR	C2-N3	2.18	1.35	1.32
2	B	302	APR	C5-C4	3.22	1.47	1.40
2	A	301	APR	C5-C4	3.37	1.48	1.40
2	B	301	APR	C5-C4	3.48	1.48	1.40

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	APR	O2'-C2'-C3'	-8.09	85.90	111.83
2	B	302	APR	N3-C2-N1	-6.60	123.11	128.86
2	B	301	APR	N3-C2-N1	-6.08	123.56	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	APR	N3-C2-N1	-5.73	123.87	128.86
2	B	301	APR	O1D-C1D-O4D	-3.72	106.14	111.14
2	A	301	APR	O5'-C5'-C4'	-3.53	96.50	109.00
2	A	301	APR	C4-C5-N7	-2.78	106.72	109.41
2	B	302	APR	C4-C5-N7	-2.71	106.79	109.41
2	B	301	APR	C4-C5-N7	-2.62	106.88	109.41
2	A	301	APR	O2A-PA-O5'	-2.58	95.96	108.14
2	A	301	APR	O1D-C1D-O4D	-2.57	107.68	111.14
2	A	301	APR	C1'-N9-C4	-2.21	122.81	126.64
2	A	301	APR	O4D-C1D-C2D	2.09	107.10	104.46
2	A	301	APR	C1D-C2D-C3D	2.21	105.10	102.30
2	B	302	APR	C1D-C2D-C3D	2.51	105.48	102.30
2	B	302	APR	C4'-O4'-C1'	2.52	112.45	109.77
2	A	301	APR	O2A-PA-O1A	2.77	126.62	112.28
2	A	301	APR	O3'-C3'-C4'	5.06	125.85	111.09

All (6) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	301	APR	C3'
2	B	301	APR	C1D
2	A	301	APR	C3'
2	A	301	APR	C1D
2	B	302	APR	C3'
2	B	302	APR	C1D

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	APR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	252/262 (96%)	0.92	33 (13%) 4 5	22, 29, 44, 59	0
1	B	248/262 (94%)	0.84	24 (9%) 8 10	22, 30, 45, 52	0
All	All	500/524 (95%)	0.88	57 (11%) 6 7	22, 30, 45, 59	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	247	ARG	6.9
1	B	213	SER	6.5
1	B	247	ARG	6.0
1	B	245	GLY	5.9
1	A	245	GLY	5.3
1	B	176	TRP	5.1
1	A	46	TRP	4.7
1	A	248	GLU	4.4
1	A	249	ASP	4.4
1	A	293	ASP	4.4
1	A	163	PHE	4.3
1	A	246	GLY	4.2
1	A	47	ARG	4.0
1	B	254	CYS	4.0
1	A	143	PHE	3.7
1	B	246	GLY	3.7
1	B	292	GLU	3.5
1	B	284	PHE	3.2
1	B	286	GLN	3.2
1	A	204	VAL	3.1
1	B	283	LYS	3.0
1	A	203	VAL	3.0
1	B	127	ARG	2.8
1	A	196	PHE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	270	ASN	2.7
1	A	238	LEU	2.6
1	B	195	ARG	2.6
1	B	250	SER	2.6
1	A	78	ARG	2.5
1	B	285	LEU	2.5
1	A	192	VAL	2.5
1	A	176	TRP	2.4
1	B	177	ARG	2.4
1	B	202	ASP	2.4
1	B	255	GLN	2.3
1	A	123	LEU	2.3
1	A	262	LEU	2.3
1	A	225	VAL	2.3
1	A	73	ILE	2.3
1	A	206	VAL	2.3
1	A	235	VAL	2.3
1	B	79	HIS	2.3
1	A	145	LEU	2.2
1	A	122	ILE	2.2
1	A	124	LEU	2.2
1	A	189	TRP	2.2
1	A	188	PHE	2.2
1	B	266	ILE	2.1
1	B	253	LEU	2.1
1	A	222	PHE	2.1
1	A	273	PHE	2.1
1	A	45	ARG	2.0
1	B	281	PRO	2.0
1	B	74	HIS	2.0
1	B	282	ASP	2.0
1	A	138	VAL	2.0
1	A	230	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	APR	B	302	36/36	0.35	0.47	5.61	139,143,150,150	0
2	APR	B	301	36/36	0.84	0.29	2.31	34,59,77,77	0
2	APR	A	301	36/36	0.91	0.19	1.70	20,27,54,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.