



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2017 – 08:15 PM EDT

PDB ID : 3I83
Title : Crystal structure of 2-dehydropantoate 2-reductase from *Methylococcus capsulatus*
Authors : Bonanno, J.B.; Gilmore, M.; Bain, K.T.; Chang, S.; Sampathkumar, P.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : unknown
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div> <div>16%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div>••</div> </div> </div>
1	B	320	<div> <div>33%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>10%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydropantoate 2-reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	1	0
			2309	1468	396	435	10			
1	B	287	Total	C	N	O	S	0	1	0
			2130	1359	365	396	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q604L6
A	0	SER	-	expression tag	UNP Q604L6
A	1	LEU	-	expression tag	UNP Q604L6
A	311	GLU	-	expression tag	UNP Q604L6
A	312	GLY	-	expression tag	UNP Q604L6
A	313	HIS	-	expression tag	UNP Q604L6
A	314	HIS	-	expression tag	UNP Q604L6
A	315	HIS	-	expression tag	UNP Q604L6
A	316	HIS	-	expression tag	UNP Q604L6
A	317	HIS	-	expression tag	UNP Q604L6
A	318	HIS	-	expression tag	UNP Q604L6
B	-1	MET	-	expression tag	UNP Q604L6
B	0	SER	-	expression tag	UNP Q604L6
B	1	LEU	-	expression tag	UNP Q604L6
B	311	GLU	-	expression tag	UNP Q604L6
B	312	GLY	-	expression tag	UNP Q604L6
B	313	HIS	-	expression tag	UNP Q604L6
B	314	HIS	-	expression tag	UNP Q604L6
B	315	HIS	-	expression tag	UNP Q604L6
B	316	HIS	-	expression tag	UNP Q604L6
B	317	HIS	-	expression tag	UNP Q604L6
B	318	HIS	-	expression tag	UNP Q604L6

- Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	167	Total	O	0	0
			167	167		
3	B	92	Total	O	0	0
			92	92		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 2-dehydropantoate 2-reductase



• Molecule 1: 2-dehydropantoate 2-reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.85Å 79.18Å 100.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 28.60 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-1.90) 99.4 (28.60-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.15 (at 1.91Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.192 , 0.227 0.202 , 0.237	Depositor DCC
R_{free} test set	2355 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4702	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/2353	0.79	2/3204 (0.1%)
1	B	0.80	2/2168 (0.1%)	0.72	1/2945 (0.0%)
All	All	0.82	2/4521 (0.0%)	0.76	3/6149 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	59	VAL	CB-CG2	6.28	1.66	1.52
1	B	294	ALA	CA-CB	5.24	1.63	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	LEU	CB-CG-CD2	-6.34	100.22	111.00
1	A	241	LYS	CD-CE-NZ	5.15	123.54	111.70
1	B	302	ARG	NE-CZ-NH1	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2309	0	2333	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2130	0	2144	12	0
2	A	4	0	3	0	0
3	A	167	0	0	0	0
3	B	92	0	0	1	0
All	All	4702	0	4480	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:ILE:HD11	1:A:275:ALA:HB2	1.60	0.81
1:A:69:LYS:HG2	1:A:70:PRO:CD	2.21	0.70
1:B:152:PRO:O	3:B:364:HOH:O	2.12	0.66
1:A:69:LYS:HD3	1:A:94:ALA:HB2	1.78	0.66
1:A:79:VAL:HB	1:A:270:VAL:HG11	1.80	0.64
1:A:69:LYS:HG2	1:A:70:PRO:HD2	1.80	0.63
1:A:77:ILE:HD12	1:A:83:ALA:HB1	1.83	0.59
1:B:198:VAL:HG21	1:B:256:MET:HG2	1.87	0.56
1:B:8:THR:HG22	1:B:28:VAL:HG11	1.88	0.55
1:B:272:LEU:HD21	1:B:292:VAL:HG12	1.88	0.53
1:A:69:LYS:HG2	1:A:70:PRO:HD3	1.91	0.52
1:B:3:ILE:HB	1:B:26:VAL:HG22	1.92	0.50
1:A:106:ILE:HD11	1:A:275:ALA:CB	2.37	0.49
1:A:125:LEU:HD11	1:A:186:LYS:HD3	1.96	0.48
1:A:77:ILE:HD11	1:A:85:ARG:HG2	1.96	0.47
1:A:69:LYS:CG	1:A:70:PRO:HD3	2.44	0.47
1:B:120:GLU:OE1	1:B:156:SER:OG	2.26	0.47
1:B:8:THR:CG2	1:B:28:VAL:HG11	2.45	0.46
1:B:154:GLY:HA2	1:B:177:GLU:HG2	1.97	0.46
1:A:77:ILE:HD11	1:A:85:ARG:CG	2.46	0.46
1:B:18:LEU:CD1	1:B:171:ILE:HD13	2.46	0.46
1:A:69:LYS:CG	1:A:70:PRO:CD	2.94	0.45
1:A:186:LYS:HG2	1:A:190[A]:ASN:ND2	2.32	0.45
1:A:120:GLU:OE1	1:A:156:SER:OG	2.23	0.44
1:B:34:TYR:CE1	1:B:59:VAL:HG12	2.53	0.44
1:A:106:ILE:CD1	1:A:275:ALA:HB2	2.40	0.43
1:B:137:GLU:C	1:B:138:ILE:HD13	2.39	0.43
1:A:10:ALA:HA	1:A:138:ILE:HD12	2.01	0.42
1:A:94:ALA:HB1	1:A:95:PRO:HD2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ILE:CD1	1:A:138:ILE:HB	2.50	0.41
1:A:248:LYS:HA	1:A:248:LYS:HD2	1.82	0.41
1:B:137:GLU:O	1:B:138:ILE:HD13	2.20	0.41
1:A:75:LEU:HD21	1:A:85:ARG:HG2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/320 (97%)	298 (96%)	11 (4%)	0	100	100
1	B	278/320 (87%)	255 (92%)	22 (8%)	1 (0%)	38	26
All	All	587/640 (92%)	553 (94%)	33 (6%)	1 (0%)	51	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	93	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/254 (94%)	231 (97%)	8 (3%)	43	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	218/254 (86%)	215 (99%)	3 (1%)	71	69
All	All	457/508 (90%)	446 (98%)	11 (2%)	54	47

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	SER
1	A	43	ARG
1	A	70	PRO
1	A	74	LEU
1	A	106	ILE
1	A	157	GLU
1	A	241	LYS
1	A	283	ARG
1	B	32	SER
1	B	40	LYS
1	B	107	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	264	GLN
1	B	24	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACY	A	319	-	1,3,3	2.78	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACY	A	319	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	319	ACY	CH3-C	2.78	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/320 (96%)	0.98	51 (16%) 2 2	13, 29, 44, 58	0
1	B	287/320 (89%)	2.13	107 (37%) 0 0	12, 23, 43, 54	0
All	All	597/640 (93%)	1.53	158 (26%) 1 1	12, 25, 44, 58	0

All (158) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	37	VAL	11.8
1	B	47	ALA	11.1
1	B	49	LEU	10.7
1	B	30	SER	10.0
1	B	41	GLY	9.4
1	B	91	ASP	8.5
1	B	58	ALA	7.9
1	B	60	VAL	7.8
1	B	84	ASP	7.8
1	B	51	ASP	7.5
1	B	48	THR	7.4
1	B	29	VAL	7.3
1	B	39	ALA	6.8
1	B	57	ALA	6.8
1	B	139	TRP	6.6
1	B	31	ARG	6.5
1	B	54	PHE	6.4
1	B	88	LEU	6.3
1	B	87	GLY	6.2
1	B	133	THR	6.2
1	B	34	TYR	6.0
1	B	143	TYR	5.9
1	B	32	SER	5.8
1	B	129	GLY	5.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	35	GLU	5.7
1	B	141	GLN	5.7
1	B	116	PHE	5.7
1	B	50	GLY	5.6
1	B	86	VAL	5.4
1	B	90	ARG	5.4
1	B	89	LEU	5.4
1	B	138	ILE	5.3
1	A	83	ALA	5.3
1	B	72	CYS	5.2
1	B	36	THR	5.1
1	B	131	THR	5.1
1	B	28	VAL	5.1
1	B	45	ARG	5.1
1	B	59	VAL	5.1
1	B	4	LEU	5.0
1	B	25	CYS	4.8
1	A	68	THR	4.8
1	B	137	GLU	4.8
1	B	117	PRO	4.7
1	B	53	THR	4.5
1	B	56	PRO	4.5
1	B	142	ALA	4.4
1	B	42	ILE	4.3
1	B	170	GLY	4.3
1	B	46	SER	4.2
1	B	130	VAL	4.2
1	B	134	ALA	4.2
1	B	8	THR	4.2
1	B	155	VAL	4.2
1	B	55	ARG	4.1
1	B	114	ALA	4.1
1	B	44	ILE	4.1
1	B	140	HIS	4.1
1	B	22	THR	4.1
1	B	76	CYS	4.0
1	B	115	ALA	4.0
1	A	74	LEU	3.8
1	B	92	ALA	3.8
1	B	2	ASN	3.8
1	B	132	ARG	3.8
1	B	26	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	21	LYS	3.7
1	B	20	ALA	3.6
1	A	67	GLU	3.6
1	B	33	ASP	3.6
1	A	82	GLY	3.6
1	A	191	ALA	3.6
1	B	157	GLU	3.5
1	B	7	GLY	3.5
1	B	111	GLU	3.5
1	B	113	ALA	3.5
1	A	95	PRO	3.4
1	B	165	ALA	3.4
1	B	38	LYS	3.4
1	A	196	LEU	3.4
1	B	24	HIS	3.3
1	B	6	ILE	3.3
1	B	40	LYS	3.3
1	A	66	LEU	3.3
1	B	27	SER	3.3
1	B	61	ARG	3.2
1	B	85	ARG	3.2
1	B	161	THR	3.1
1	B	17	ALA	3.1
1	A	294	ALA	3.1
1	B	136	GLY	3.0
1	A	139	TRP	3.0
1	A	215	PHE	3.0
1	B	295	LEU	3.0
1	A	195	PRO	3.0
1	B	154	GLY	3.0
1	A	94	ALA	2.9
1	A	199	LEU	2.9
1	A	76	CYS	2.9
1	A	295	LEU	2.9
1	A	61	ARG	2.8
1	B	135	PRO	2.8
1	A	219	ILE	2.7
1	B	156	SER	2.7
1	B	52	TYR	2.7
1	B	3	ILE	2.7
1	A	69	LYS	2.7
1	B	175	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	158	ARG	2.7
1	A	102	ILE	2.7
1	A	192	ALA	2.7
1	B	164	ALA	2.7
1	B	73	THR	2.6
1	B	128	ILE	2.6
1	A	5	VAL	2.6
1	A	188	VAL	2.6
1	A	75	LEU	2.6
1	A	91	ASP	2.6
1	A	131	THR	2.6
1	A	289	LEU	2.5
1	B	219	ILE	2.5
1	B	191	ALA	2.5
1	A	6	ILE	2.4
1	A	292	VAL	2.4
1	B	121	VAL	2.4
1	B	171	ILE	2.4
1	A	100	VAL	2.3
1	B	99	ILE	2.3
1	B	271	ILE	2.3
1	B	292	VAL	2.3
1	A	272	LEU	2.3
1	B	23	GLY	2.3
1	B	98	GLY	2.3
1	B	206[A]	LEU	2.3
1	A	16	GLY	2.3
1	A	189	TRP	2.3
1	A	106	ILE	2.3
1	B	199	LEU	2.3
1	A	296	MET	2.2
1	A	84	ASP	2.2
1	A	307	LEU	2.2
1	B	169	ALA	2.2
1	A	93	VAL	2.2
1	A	291	SER	2.2
1	B	289	LEU	2.1
1	A	60	VAL	2.1
1	A	193	PHE	2.1
1	B	9	GLY	2.1
1	A	30	SER	2.1
1	A	310	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	271	ILE	2.1
1	A	19	LEU	2.1
1	A	51	ASP	2.1
1	A	48	THR	2.1
1	A	59	VAL	2.1
1	A	141	GLN	2.1
1	B	102	ILE	2.0
1	A	187	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACY	A	319	4/4	0.93	0.13	-0.77	26,26,26,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.