



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2018 – 08:37 AM EST

PDB ID : 1I9V
Title : CRYSTAL STRUCTURE ANALYSIS OF A TRNA-NEOMYCIN COMPLEX
Authors : Mikkelsen, N.E.; Johansson, K.; Virtanen, A.; Kirsebom, L.A.
Deposited on : 2001-03-21
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2895 (2.60-2.60)
RNA backbone	2435	1140 (3.00-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	76	 51% 33% 11% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NMY	A	200	X	-	-	-

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Mg 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total 32	O 32	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PHENYLALANINE TRANSFER RNA



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.90 Å 32.90 Å 63.10 Å 90.00° 90.50° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60	Depositor
% Data completeness (in resolution range)	90.8 (20.00-2.60)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.228 , 0.275	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1708	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5MU, MG, YG, NMY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	5/1757 (0.3%)	1.13	15/2738 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54	5MU	O3'-P	-5.93	1.54	1.61
1	A	39	U	O3'-P	5.67	1.68	1.61
1	A	16	U	O3'-P	5.44	1.67	1.61
1	A	6	U	O3'-P	5.10	1.67	1.61
1	A	39	U	C3'-O3'	5.09	1.49	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39	U	O4'-C4'-C3'	-20.39	83.61	104.00
1	A	39	U	C1'-O4'-C4'	-13.97	98.72	109.90
1	A	39	U	C4'-C3'-C2'	-12.90	89.70	102.60
1	A	39	U	C2'-C3'-O3'	12.66	137.34	109.50
1	A	38	A	P-O3'-C3'	9.38	130.95	119.70
1	A	39	U	C3'-C2'-C1'	-8.23	94.91	101.50
1	A	55	U	C2'-C3'-O3'	7.61	126.23	109.50
1	A	6	U	OP2-P-O3'	7.16	120.95	105.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	12	U	OP1-P-O3'	7.15	120.94	105.20
1	A	39	U	C5'-C4'-C3'	6.98	127.16	116.00
1	A	55	U	C3'-C2'-C1'	6.83	106.97	101.50
1	A	39	U	C1'-C2'-O2'	6.22	129.25	110.60
1	A	40	C	O5'-P-OP1	-5.47	100.77	105.70
1	A	55	U	C5-C6-N1	-5.42	119.99	122.70
1	A	19	G	O4'-C1'-N9	-5.15	104.08	108.20

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	C	Sidechain
1	A	12	U	Sidechain
1	A	16	U	Sidechain
1	A	17	U	Sidechain
1	A	19	G	Sidechain
1	A	36	A	Sidechain
1	A	6	U	Sidechain
1	A	7	U	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	834	24	0
2	A	42	0	46	2	0
3	A	3	0	0	0	0
4	A	32	0	0	1	0
All	All	1708	0	880	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:YG:H101	1:A:37:YG:HN2	1.59	0.67
1:A:55:U:H5	1:A:58:A:OP2	1.80	0.65
1:A:55:U:O2	1:A:55:U:O5'	2.15	0.63
1:A:68:U:O2'	1:A:69:U:H5'	2.04	0.57
1:A:37:YG:C10	1:A:37:YG:HN2	2.17	0.57
1:A:3:G:H1	1:A:70:C:H42	1.55	0.55
1:A:55:U:O2'	1:A:57:G:N7	2.36	0.55
1:A:36:A:O2'	1:A:37:YG:H5''	2.11	0.50
1:A:37:YG:H101	1:A:37:YG:N20	2.24	0.50
1:A:24:G:H2'	1:A:25:C:O4'	2.12	0.49
1:A:19:G:H8	4:A:120:HOH:O	1.96	0.48
1:A:38:A:H2'	1:A:39:U:C1'	2.46	0.46
1:A:10:G:N3	1:A:10:G:H2'	2.31	0.46
1:A:19:G:C5	1:A:57:G:N2	2.85	0.44
1:A:64:A:H2'	1:A:65:G:O4'	2.18	0.44
1:A:19:G:C4	1:A:57:G:N2	2.86	0.43
1:A:7:U:O2'	1:A:49:C:OP2	2.33	0.43
1:A:17:U:H3'	1:A:18:G:H5''	2.01	0.43
1:A:51:G:H2'	1:A:52:U:O4'	2.18	0.43
1:A:23:A:OP2	2:A:200:NMY:H62	2.19	0.43
1:A:1:G:N2	1:A:2:C:C2	2.88	0.42
1:A:16:U:O2	1:A:16:U:H2'	2.19	0.41
1:A:1:G:O2'	1:A:2:C:H5'	2.20	0.41
1:A:45:G:O6	2:A:200:NMY:H61	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	75/76 (98%)	9 (12%)	0

All (9) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	16	U
1	A	17	U
1	A	18	G
1	A	21	A
1	A	39	U
1	A	40	C
1	A	47	U
1	A	61	C
1	A	76	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	YG	A	37	1	29,42,43	2.45	9 (31%)	29,62,65	2.40	9 (31%)
1	5MU	A	54	1	14,22,23	1.18	2 (14%)	16,32,35	4.13	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YG	A	37	1	-	0/20/42/43	0/4/4/4
1	5MU	A	54	1	-	0/3/25/26	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	37	YG	O3'-C3'	-4.45	1.32	1.43
1	A	37	YG	O23-C21	-3.92	1.28	1.34
1	A	37	YG	C6-N1	-3.83	1.31	1.37
1	A	37	YG	C3'-C2'	-3.36	1.44	1.53
1	A	54	5MU	C6-C5	-2.30	1.34	1.40
1	A	37	YG	C15-N20	2.41	1.51	1.45
1	A	37	YG	C10-C11	2.44	1.54	1.50
1	A	37	YG	O17-C16	2.96	1.28	1.21
1	A	54	5MU	C4-N3	3.10	1.38	1.33
1	A	37	YG	O22-C21	3.36	1.28	1.21
1	A	37	YG	O4'-C1'	7.52	1.51	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	5MU	C5-C4-N3	-8.85	115.48	125.24
1	A	37	YG	O22-C21-N20	-3.46	118.91	124.87
1	A	37	YG	O17-C16-C15	-2.85	115.43	123.91
1	A	54	5MU	C5M-C5-C6	2.16	122.98	118.67
1	A	37	YG	O18-C16-C15	2.47	118.06	111.54
1	A	37	YG	O3'-C3'-C4'	2.49	118.36	111.09
1	A	37	YG	O2'-C2'-C1'	2.59	119.70	111.61
1	A	37	YG	O23-C21-N20	2.79	116.17	110.82
1	A	37	YG	C6-C5-C4	4.51	123.04	119.92
1	A	37	YG	C2'-C3'-C4'	5.04	112.43	102.62
1	A	37	YG	C3-N3-C4	6.38	127.41	118.31
1	A	54	5MU	C4-N3-C2	13.59	127.05	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	37	YG	4	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NMY	A	200	-	45,45,45	2.14	14 (31%)	59,67,67	1.36	8 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NMY	A	200	-	11/11/19/19	0/18/94/94	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	200	NMY	O18-C15	2.02	1.48	1.43
2	A	200	NMY	C21-C22	2.15	1.57	1.53
2	A	200	NMY	C20-C21	2.22	1.58	1.52
2	A	200	NMY	O16-C13	2.31	1.45	1.41
2	A	200	NMY	C18-C19	2.50	1.57	1.52
2	A	200	NMY	O18-C18	2.59	1.48	1.41
2	A	200	NMY	C10-C11	2.87	1.58	1.52
2	A	200	NMY	C10-C9	2.99	1.59	1.53
2	A	200	NMY	C4-C3	3.04	1.60	1.52
2	A	200	NMY	C3-C2	3.62	1.58	1.53
2	A	200	NMY	O5-C5	4.08	1.54	1.44
2	A	200	NMY	O5-C1	4.44	1.52	1.41
2	A	200	NMY	O22-C22	5.61	1.58	1.44
2	A	200	NMY	O22-C18	6.71	1.58	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	200	NMY	C1-O5-C5	-3.97	106.24	113.72
2	A	200	NMY	C13-O11-C11	-3.65	109.09	118.00
2	A	200	NMY	C1-O1-C10	-3.00	110.69	118.00
2	A	200	NMY	O11-C13-C14	2.34	112.81	107.96
2	A	200	NMY	O5-C5-C6	2.49	110.72	106.01
2	A	200	NMY	C13-C14-C15	2.52	105.14	102.07
2	A	200	NMY	O22-C22-C23	2.63	110.98	106.01
2	A	200	NMY	C3-C4-C5	3.28	116.00	110.22

All (11) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	200	NMY	C18
2	A	200	NMY	C9
2	A	200	NMY	C7
2	A	200	NMY	C10
2	A	200	NMY	C11
2	A	200	NMY	C1
2	A	200	NMY	C12
2	A	200	NMY	C3
2	A	200	NMY	C20
2	A	200	NMY	C14
2	A	200	NMY	C19

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	200	NMY	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.