



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:20 am GMT

PDB ID : 5IBL
Title : Human antibody 6639 in complex with influenza hemagglutinin H1 X-181
Authors : Raymond, D.D.; Harrison, S.C.
Deposited on : 2016-02-22
Resolution : 3.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1679 (3.50-3.30)
Clashscore	112137	1832 (3.50-3.30)
Ramachandran outliers	110173	1789 (3.50-3.30)
Sidechain outliers	110143	1789 (3.50-3.30)
RSRZ outliers	101464	1709 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	<div> <div>23%</div> <div> <div>53%</div> <div>6%</div> <div>40%</div> </div> </div>
1	E	176	<div> <div>7%</div> <div> <div>56%</div> <div>7%</div> <div>35%</div> </div> </div>
2	B	328	<div> <div>13%</div> <div> <div>76%</div> <div>14%</div> <div>9%</div> </div> </div>
2	F	328	<div> <div>2%</div> <div> <div>77%</div> <div>15%</div> <div>7%</div> </div> </div>
3	C	230	<div> <div>13%</div> <div> <div>73%</div> <div>20%</div> <div>6%</div> </div> </div>
3	H	230	<div> <div>11%</div> <div> <div>78%</div> <div>15%</div> <div>6%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	215	 20% 81% 18% •
4	L	215	 20% 84% 15% •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	106	Total	C	N	O	S	0	0	0
			858	534	141	179	4			
1	E	115	Total	C	N	O	S	0	0	0
			931	584	152	191	4			

- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	298	Total	C	N	O	S	0	0	0
			2336	1480	401	445	10			
2	F	306	Total	C	N	O	S	0	0	0
			2398	1517	416	455	10			

- Molecule 3 is a protein called 6639 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	216	Total	C	N	O	S	0	1	0
			1643	1043	273	319	8			
3	H	216	Total	C	N	O	S	0	0	0
			1639	1041	271	319	8			

- Molecule 4 is a protein called 6639 Light Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	215	Total	C	N	O	S	0	0	0
			1639	1022	278	333	6			
4	L	215	Total	C	N	O	S	0	0	0
			1639	1022	278	333	6			

- Molecule 5 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 2: Hemagglutinin




- Molecule 3: 6639 Heavy Chain




- Molecule 3: 6639 Heavy Chain



- Molecule 4: 6639 Light Chain



- Molecule 4: 6639 Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.13Å 104.72Å 116.66Å 101.95° 95.33° 98.36°	Depositor
Resolution (Å)	44.81 – 3.39 46.40 – 3.39	Depositor EDS
% Data completeness (in resolution range)	95.4 (44.81-3.39) 88.4 (46.40-3.39)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 3.40Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.214 , 0.257 0.249 , 0.303	Depositor DCC
R_{free} test set	1636 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	113.2	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 107.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	13181	wwPDB-VP
Average B, all atoms (Å ²)	140.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/864	0.57	0/1159
1	E	0.41	0/942	0.58	0/1266
2	B	0.39	0/2396	0.63	0/3255
2	F	0.41	0/2458	0.67	0/3338
3	C	0.38	0/1690	0.61	0/2302
3	H	0.40	0/1682	0.64	0/2291
4	D	0.38	0/1674	0.60	0/2273
4	L	0.39	0/1674	0.60	0/2273
All	All	0.39	0/13380	0.62	0/18157

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	858	0	834	4	0
1	E	931	0	892	13	0
2	B	2336	0	2281	21	0
2	F	2398	0	2347	23	0
3	C	1643	0	1598	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	1639	0	1597	12	0
4	D	1639	0	1591	15	0
4	L	1639	0	1591	16	0
5	B	28	0	26	0	0
5	F	70	0	65	2	0
All	All	13181	0	12822	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:69:GLU:OE1	1:E:69:GLU:N	2.06	0.87
1:E:83:LYS:HD3	1:E:83:LYS:C	2.00	0.80
4:L:2:ILE:HD11	4:L:94:ASN:HB3	1.64	0.79
2:B:74:PRO:HB2	2:B:141:HIS:HB2	1.72	0.72
1:E:83:LYS:O	1:E:83:LYS:HD3	1.91	0.70

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	96/176 (54%)	93 (97%)	2 (2%)	1 (1%)	18 59
1	E	105/176 (60%)	92 (88%)	10 (10%)	3 (3%)	5 36
2	B	294/328 (90%)	261 (89%)	31 (10%)	2 (1%)	25 65
2	F	302/328 (92%)	265 (88%)	32 (11%)	5 (2%)	11 47
3	C	213/230 (93%)	194 (91%)	18 (8%)	1 (0%)	32 71

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	H	212/230 (92%)	191 (90%)	18 (8%)	3 (1%)	13	51
4	D	213/215 (99%)	194 (91%)	16 (8%)	3 (1%)	13	51
4	L	213/215 (99%)	195 (92%)	16 (8%)	2 (1%)	20	61
All	All	1648/1898 (87%)	1485 (90%)	143 (9%)	20 (1%)	15	54

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	126	SER
4	D	71	ASP
4	D	139	ASN
1	E	85	ASP
2	F	126	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/151 (65%)	90 (92%)	8 (8%)	13	47
1	E	103/151 (68%)	95 (92%)	8 (8%)	15	49
2	B	262/287 (91%)	248 (95%)	14 (5%)	26	63
2	F	268/287 (93%)	250 (93%)	18 (7%)	19	57
3	C	184/195 (94%)	172 (94%)	12 (6%)	20	58
3	H	183/195 (94%)	170 (93%)	13 (7%)	17	54
4	D	186/186 (100%)	170 (91%)	16 (9%)	12	44
4	L	186/186 (100%)	175 (94%)	11 (6%)	23	60
All	All	1470/1638 (90%)	1370 (93%)	100 (7%)	18	56

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	138	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	120	GLU
4	L	46	ARG
4	D	164	VAL
1	E	38	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	114	ASN
4	L	138	ASN
2	F	47	HIS
3	C	52	ASN
4	L	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	401	2	14,14,15	0.35	0	15,19,21	0.60	0
5	NAG	B	402	2	14,14,15	0.35	0	15,19,21	1.41	2 (13%)
5	NAG	F	401	2	14,14,15	0.32	0	15,19,21	0.56	0
5	NAG	F	402	2	14,14,15	0.33	0	15,19,21	0.71	1 (6%)
5	NAG	F	403	2	14,14,15	0.37	0	15,19,21	0.66	0
5	NAG	F	404	2	14,14,15	0.39	0	15,19,21	1.09	2 (13%)
5	NAG	F	405	2	14,14,15	0.41	0	15,19,21	2.13	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	401	2	-	0/6/23/26	0/1/1/1
5	NAG	B	402	2	-	0/6/23/26	0/1/1/1
5	NAG	F	401	2	-	0/6/23/26	0/1/1/1
5	NAG	F	402	2	-	0/6/23/26	0/1/1/1
5	NAG	F	403	2	-	0/6/23/26	0/1/1/1
5	NAG	F	404	2	-	0/6/23/26	0/1/1/1
5	NAG	F	405	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	402	NAG	O5-C1-C2	-4.15	105.69	111.47
5	F	404	NAG	C1-C2-N2	-2.91	105.51	110.49
5	F	402	NAG	C1-O5-C5	2.01	114.94	112.17
5	F	404	NAG	O5-C1-C2	2.20	114.54	111.47
5	B	402	NAG	C1-C2-N2	2.86	115.38	110.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	404	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	106/176 (60%)	1.59	40 (37%) 0 0	131, 184, 216, 242	0
1	E	115/176 (65%)	0.73	12 (10%) 7 7	107, 153, 210, 221	0
2	B	298/328 (90%)	0.93	42 (14%) 3 3	85, 126, 174, 200	0
2	F	306/328 (93%)	0.52	5 (1%) 72 67	81, 113, 140, 160	0
3	C	216/230 (93%)	0.95	31 (14%) 3 3	103, 154, 195, 210	0
3	H	216/230 (93%)	0.79	25 (11%) 5 6	79, 130, 197, 218	0
4	D	215/215 (100%)	1.19	42 (19%) 1 1	105, 143, 197, 211	0
4	L	215/215 (100%)	1.12	44 (20%) 1 1	96, 137, 202, 229	0
All	All	1687/1898 (88%)	0.92	241 (14%) 3 3	79, 136, 198, 242	0

The worst 5 of 241 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	L	182	LEU	7.1
4	D	195	CYS	7.1
4	L	118	ILE	6.9
4	D	192	VAL	6.6
1	A	19	ASP	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NAG	F	403	14/15	0.85	0.24	0.17	126,136,140,140	0
5	NAG	F	405	14/15	0.58	0.27	-0.03	159,160,161,161	0
5	NAG	F	404	14/15	0.80	0.31	-0.21	144,145,148,149	0
5	NAG	B	402	14/15	0.81	0.30	-0.40	173,176,179,179	0
5	NAG	B	401	14/15	0.80	0.26	-	160,166,175,175	0
5	NAG	F	401	14/15	0.80	0.39	-	173,180,184,186	0
5	NAG	F	402	14/15	0.71	0.28	-	147,148,154,155	0

6.5 Other polymers

There are no such residues in this entry.