



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 06:21 pm GMT

PDB ID : 1ICH  
Title : SOLUTION STRUCTURE OF THE TUMOR NECROSIS FACTOR  
RECEPTOR-1 DEATH DOMAIN  
Authors : Sukits, S.F.; Lin, L.-L.; Malakian, K.; Powers, R.; Xu, G.-Y.  
Deposited on : 2001-04-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 72%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	112	

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1442 atoms, of which 728 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called TUMOR NECROSIS FACTOR RECEPTOR-1.

Mol	Chain	Residues	Atoms						Trace
1	A	87	Total	C	H	N	O	S	0
			1442	447	728	133	129	5	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	315	MET	-	INITIATING METHIONINE	UNP P19438
A	347	LYS	ARG	ENGINEERED	UNP P19438

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

### • Molecule 1: TUMOR NECROSIS FACTOR RECEPTOR-1

Chain A: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *Distance geometry, Simulated annealing molecular dynamics*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	9.8
X-PLOR	refinement	9.8

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5018
Number of chemical shift lists	1
Total number of shifts	1102
Number of shifts mapped to atoms	1102
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	72%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	12
All	All	0	12

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	380	ARG	Sidechain
1	A	381	ARG	Sidechain
1	A	368	ARG	Sidechain
1	A	341	ARG	Sidechain
1	A	358	ARG	Sidechain
1	A	348	ARG	Sidechain
1	A	365	ARG	Sidechain
1	A	397	ARG	Sidechain
1	A	394	ARG	Sidechain
1	A	385	ARG	Sidechain
1	A	379	ARG	Sidechain
1	A	384	ARG	Sidechain

### 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	714	728	728	40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	714	728	728	40

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:332:ALA:HB3	1:A:408:ILE:HG23	0.85	1.48
1:A:329:THR:OG1	1:A:412:LEU:HD21	0.81	1.75
1:A:332:ALA:CB	1:A:408:ILE:HG23	0.75	2.12
1:A:405:LEU:O	1:A:405:LEU:HD13	0.72	1.85
1:A:329:THR:CG2	1:A:389:LEU:HD13	0.70	2.15
1:A:405:LEU:HA	1:A:408:ILE:HD12	0.69	1.64
1:A:378:TRP:CZ2	1:A:391:LEU:HD22	0.69	2.23
1:A:349:LEU:HD12	1:A:395:VAL:HG21	0.67	1.64
1:A:330:LEU:HD22	1:A:375:LEU:HB3	0.64	1.68
1:A:339:PRO:HB2	1:A:367:LEU:HD21	0.64	1.69
1:A:329:THR:HG22	1:A:389:LEU:HD13	0.62	1.71
1:A:349:LEU:HD23	1:A:378:TRP:CE3	0.62	2.30
1:A:342:TRP:CD1	1:A:346:VAL:CG2	0.56	2.89
1:A:405:LEU:C	1:A:405:LEU:HD13	0.55	2.22
1:A:334:VAL:HG11	1:A:368:ARG:HG3	0.54	1.78
1:A:345:PHE:CZ	1:A:392:LEU:HD23	0.52	2.38
1:A:329:THR:HG21	1:A:389:LEU:HB2	0.52	1.79
1:A:342:TRP:CD1	1:A:346:VAL:HG23	0.52	2.40
1:A:386:GLU:O	1:A:387:ALA:HB3	0.51	2.04
1:A:339:PRO:CB	1:A:367:LEU:HD21	0.50	2.36
1:A:348:ARG:CB	1:A:395:VAL:HG13	0.50	2.37
1:A:332:ALA:HB1	1:A:408:ILE:HA	0.48	1.84
1:A:334:VAL:HA	1:A:342:TRP:CZ3	0.47	2.45
1:A:341:ARG:N	1:A:341:ARG:CD	0.47	2.78
1:A:387:ALA:HB1	1:A:390:GLU:OE1	0.47	2.08
1:A:396:LEU:HD22	1:A:401:LEU:HB3	0.47	1.85
1:A:388:THR:O	1:A:392:LEU:HD12	0.47	2.09
1:A:343:LYS:HG3	1:A:356:ILE:HG21	0.46	1.87
1:A:378:TRP:CH2	1:A:391:LEU:HD22	0.46	2.45
1:A:378:TRP:C	1:A:378:TRP:CD1	0.45	2.90
1:A:345:PHE:HZ	1:A:392:LEU:HD23	0.44	1.72
1:A:372:TYR:CE2	1:A:376:ALA:HB2	0.44	2.47
1:A:374:MET:O	1:A:378:TRP:CB	0.43	2.66
1:A:378:TRP:HZ2	1:A:391:LEU:HD13	0.43	1.72
1:A:375:LEU:CD1	1:A:375:LEU:N	0.42	2.82

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:333:VAL:HB	1:A:375:LEU:HD21	0.42	1.92
1:A:393:GLY:HA2	1:A:396:LEU:HD12	0.42	1.91
1:A:373:SER:O	1:A:377:THR:N	0.41	2.47
1:A:346:VAL:HG13	1:A:374:MET:CE	0.40	2.46
1:A:348:ARG:HB2	1:A:395:VAL:HG13	0.40	1.92

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/112 (76%)	81 (95%)	2 (2%)	2 (2%)	11	48
All	All	85/112 (76%)	81 (95%)	2 (2%)	2 (2%)	11	48

All 2 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	363	ASN
1	A	339	PRO

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/97 (79%)	73 (95%)	4 (5%)	31	77
All	All	77/97 (79%)	73 (95%)	4 (5%)	31	77

All 4 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	363	ASN
1	A	392	LEU
1	A	375	LEU
1	A	341	ARG

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 72% for the well-defined parts and 72% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5018

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1102
Number of shifts mapped to atoms	1102
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	105	$-0.77 \pm 0.14$	Should be applied
$^{13}\text{C}_\beta$	100	$0.35 \pm 0.10$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	92	$-0.38 \pm 0.18$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	96	$0.38 \pm 0.29$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 72%, i.e. 834 atoms were assigned a chemical shift out of a possible 1154. 12 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	407/427 (95%)	163/170 (96%)	164/174 (94%)	80/83 (96%)
Sidechain	391/671 (58%)	215/393 (55%)	176/236 (75%)	0/42 (0%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	36/56 (64%)	18/29 (62%)	16/24 (67%)	2/3 (67%)
Overall	834/1154 (72%)	396/592 (67%)	356/434 (82%)	82/128 (64%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 72%, i.e. 834 atoms were assigned a chemical shift out of a possible 1154. 12 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	407/427 (95%)	163/170 (96%)	164/174 (94%)	80/83 (96%)
Sidechain	391/671 (58%)	215/393 (55%)	176/236 (75%)	0/42 (0%)
Aromatic	36/56 (64%)	18/29 (62%)	16/24 (67%)	2/3 (67%)
Overall	834/1154 (72%)	396/592 (67%)	356/434 (82%)	82/128 (64%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	397	ARG	CG	19.64	33.23 – 21.23	-6.3

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

