



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:19 am GMT

PDB ID : 3ID8  
Title : Ternary complex of human pancreatic glucokinase crystallized with activator, glucose and AMP-PNP  
Authors : Petit, P.; Gluais, L.; Lagarde, A.; Boutin, J.A.; Ferry, G.; Vuillard, L.  
Deposited on : 2009-07-20  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	<div> <div>5%</div> <div> <div></div> <div>76%</div> <div>16%</div> <div>.</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ANP	A	600	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glucokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	450	3557	2215	620	690	32	0	3	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	MET	-	EXPRESSION TAG	UNP P35557
A	-3	GLY	-	EXPRESSION TAG	UNP P35557
A	-2	HIS	-	EXPRESSION TAG	UNP P35557
A	-1	HIS	-	EXPRESSION TAG	UNP P35557
A	0	HIS	-	EXPRESSION TAG	UNP P35557
A	1	HIS	-	EXPRESSION TAG	UNP P35557
A	2	HIS	-	EXPRESSION TAG	UNP P35557
A	3	HIS	-	EXPRESSION TAG	UNP P35557
A	4	GLU	-	EXPRESSION TAG	UNP P35557
A	5	ASN	-	EXPRESSION TAG	UNP P35557
A	6	LEU	-	EXPRESSION TAG	UNP P35557
A	7	TYR	-	EXPRESSION TAG	UNP P35557
A	8	PHE	-	EXPRESSION TAG	UNP P35557
A	9	GLN	-	EXPRESSION TAG	UNP P35557
A	10	GLY	-	EXPRESSION TAG	UNP P35557
A	11	MET	-	EXPRESSION TAG	UNP P35557

- Molecule 2 is SUGAR (ALPHA-D-GLUCOSE) (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is 2-AMINO-4-FLUORO-5-[(1-METHYL-1H-IMIDAZOL-2-YL)SULFANYL]-N-(1,3-THIAZOL-2-YL)BENZAMIDE (three-letter code: MRK) (formula:  $C_{14}H_{12}FN_5OS_2$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	0
			23	14	1	5	1	2		

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	K	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	125	Total	O	0	0
			125	125		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Glucokinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.00Å 82.60Å 86.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 2.40 19.95 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.94-2.40) 99.6 (19.95-2.40)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.17 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.201 , 0.279 0.213 , 0.292	Depositor DCC
$R_{free}$ test set	966 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.6	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3750	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GLC, ANP, K, MRK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.62	0/3620	0.68	2/4861 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	47	LEU	CA-CB-CG	5.72	128.45	115.30
1	A	315	LEU	CA-CB-CG	5.21	127.28	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	192	ARG	Sidechain
1	A	394	ARG	Sidechain

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3557	0	3505	77	0
2	A	12	0	12	0	0
3	A	23	0	12	1	0
4	A	31	0	13	2	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	A	125	0	0	1	0
All	All	3750	0	3542	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ASP:HB3	1:A:195:PHE:CA	1.64	1.23
1:A:194:ASP:CB	1:A:195:PHE:HA	1.66	1.21
1:A:195:PHE:CD1	1:A:196:GLU:HB2	1.79	1.16
1:A:106:GLN:HE22	1:A:136:LYS:HE2	1.12	1.14
1:A:394:ARG:HH11	1:A:433:SER:HB3	1.09	1.10
1:A:196:GLU:OE2	1:A:196:GLU:HA	1.55	1.05
1:A:195:PHE:CD1	1:A:196:GLU:N	2.29	1.01
1:A:195:PHE:CD1	1:A:196:GLU:CB	2.49	0.94
1:A:195:PHE:HD1	1:A:196:GLU:HB2	1.15	0.89
1:A:106:GLN:NE2	1:A:136:LYS:HE2	1.88	0.88
1:A:394:ARG:NH1	1:A:433:SER:HB3	1.90	0.86
1:A:228:THR:H	4:A:600:ANP:HNB1	1.22	0.86
1:A:195:PHE:CE1	1:A:196:GLU:CB	2.60	0.84
1:A:194:ASP:HB3	1:A:195:PHE:HA	0.87	0.84
1:A:394:ARG:HH11	1:A:433:SER:CB	1.90	0.84
1:A:112:GLU:CD	1:A:112:GLU:H	1.82	0.83
1:A:195:PHE:HE1	1:A:196:GLU:CG	1.94	0.80
1:A:195:PHE:CD1	1:A:196:GLU:CA	2.64	0.80
1:A:195:PHE:CG	1:A:196:GLU:N	2.52	0.76
1:A:195:PHE:CE1	1:A:196:GLU:HB2	2.19	0.71
1:A:195:PHE:CE1	1:A:196:GLU:CG	2.74	0.71
1:A:195:PHE:CE1	1:A:196:GLU:OE2	2.46	0.68
1:A:194:ASP:HB3	1:A:195:PHE:CB	2.23	0.67
1:A:228:THR:HG22	1:A:296:LYS:HB2	1.80	0.64
1:A:394:ARG:NH1	1:A:433:SER:CB	2.56	0.64

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:LYS:O	1:A:17:GLU:HG2	1.98	0.63
1:A:394:ARG:NH1	1:A:400:ASP:O	2.32	0.62
1:A:194:ASP:N	1:A:195:PHE:HB2	2.14	0.62
3:A:501:MRK:O8	3:A:501:MRK:S1	2.58	0.61
1:A:41:MET:HE3	1:A:389:VAL:HG22	1.83	0.61
1:A:195:PHE:CE1	1:A:196:GLU:CA	2.85	0.60
1:A:43:ARG:HD2	7:A:580:HOH:O	2.03	0.59
1:A:195:PHE:CE1	1:A:196:GLU:CD	2.76	0.58
1:A:5:ASN:OD1	1:A:8:PHE:CE1	2.58	0.56
1:A:195:PHE:HE1	1:A:196:GLU:CD	2.10	0.55
1:A:193:GLY:C	1:A:195:PHE:HB2	2.26	0.55
1:A:85:ARG:NH1	1:A:107:MET:HE1	2.21	0.55
1:A:195:PHE:CD1	1:A:195:PHE:C	2.83	0.51
1:A:12:LYS:O	1:A:16:VAL:HG23	2.10	0.51
1:A:64:SER:HB3	1:A:250:ARG:HB2	1.92	0.51
1:A:229:GLY:N	4:A:600:ANP:O1G	2.34	0.51
1:A:194:ASP:CA	1:A:195:PHE:HB2	2.41	0.50
1:A:228:THR:CG2	1:A:296:LYS:HB2	2.41	0.50
1:A:194:ASP:CB	1:A:195:PHE:CA	2.40	0.50
1:A:195:PHE:HE1	1:A:196:GLU:HG2	1.73	0.50
1:A:75:LEU:O	1:A:146:LEU:HD12	2.11	0.49
1:A:5:ASN:OD1	1:A:8:PHE:CD1	2.66	0.49
1:A:94:GLY:O	1:A:98:GLN:HB3	2.13	0.49
1:A:85:ARG:NH1	1:A:107:MET:CE	2.76	0.48
1:A:194:ASP:CG	1:A:195:PHE:HA	2.34	0.48
1:A:322:GLU:O	1:A:326:THR:HG22	2.13	0.47
1:A:322:GLU:O	1:A:326:THR:CG2	2.63	0.47
1:A:75:LEU:HD23	1:A:146:LEU:HD13	1.96	0.47
1:A:337:GLN:HE21	1:A:351:ILE:HD11	1.80	0.47
1:A:190:LYS:HA	1:A:195:PHE:CE2	2.49	0.47
1:A:83:ASN:HD22	1:A:109:SER:HA	1.80	0.47
1:A:195:PHE:CE1	1:A:196:GLU:HA	2.50	0.46
1:A:424:HIS:O	1:A:428:ARG:HG3	2.15	0.46
1:A:46[A]:ARG:HB2	1:A:49:THR:OG1	2.15	0.46
1:A:420:LYS:HG2	1:A:424:HIS:CE1	2.51	0.45
1:A:44:GLY:O	1:A:50:HIS:HD2	2.00	0.45
1:A:189:ILE:HG22	1:A:195:PHE:CD2	2.51	0.45
1:A:63:ARG:O	1:A:64:SER:O	2.35	0.45
1:A:338:VAL:HG13	1:A:348:ILE:HG12	1.98	0.45
1:A:308:ARG:O	1:A:312:GLU:HG2	2.18	0.43
1:A:112:GLU:N	1:A:112:GLU:CD	2.60	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:LYS:HE3	1:A:282:ALA:HB1	1.99	0.43
1:A:190:LYS:HA	1:A:195:PHE:HE2	1.84	0.43
1:A:215:TYR:CE1	1:A:451:LEU:HD13	2.54	0.42
1:A:315:LEU:HD13	1:A:316:PHE:CE2	2.54	0.42
1:A:194:ASP:CA	1:A:195:PHE:CB	2.99	0.41
1:A:64:SER:HB3	1:A:250:ARG:CB	2.51	0.41
1:A:195:PHE:CZ	1:A:196:GLU:OE2	2.74	0.41
1:A:77:LEU:HD21	1:A:146:LEU:HD11	2.03	0.41
1:A:131:SER:OG	1:A:140:LYS:HE3	2.21	0.41
1:A:75:LEU:HD23	1:A:146:LEU:CD1	2.51	0.41
1:A:65:THR:HA	1:A:66:PRO:HD3	1.90	0.40
1:A:92:GLY:HA3	1:A:99:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	449/470 (96%)	433 (96%)	10 (2%)	6 (1%)	14 19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	SER
1	A	194	ASP
1	A	138	GLN
1	A	141	HIS
1	A	136	LYS
1	A	80	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	390/403 (97%)	355 (91%)	35 (9%)	11	16

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	6	LEU
1	A	47	LEU
1	A	70	GLU
1	A	82	THR
1	A	89	VAL
1	A	98	GLN
1	A	104	LYS
1	A	112	GLU
1	A	131	SER
1	A	132	ASP
1	A	138	GLN
1	A	140	LYS
1	A	141	HIS
1	A	144	LEU
1	A	155[A]	ARG
1	A	155[B]	ARG
1	A	166	ASN
1	A	196	GLU
1	A	209	THR
1	A	243	LEU
1	A	245	GLU
1	A	248	GLU
1	A	257	TRP
1	A	268	GLU
1	A	271	LEU
1	A	304	LEU
1	A	307	LEU
1	A	309	LEU
1	A	315	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	326	THR
1	A	353	SER
1	A	386	LEU
1	A	442	GLU
1	A	445	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	50	HIS
1	A	106	GLN
1	A	141	HIS
1	A	337	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	A	500	-	12,12,12	0.61	0	17,17,17	0.66	0
3	MRK	A	501	-	22,25,25	2.68	5 (22%)	19,35,35	2.72	7 (36%)
4	ANP	A	600	5	29,33,33	1.59	3 (10%)	28,52,52	2.55	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	500	-	-	0/2/22/22	0/1/1/1
3	MRK	A	501	-	-	0/8/12/12	0/3/3/3
4	ANP	A	600	5	-	0/13/38/38	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	MRK	C12-S16	-5.21	1.70	1.78
3	A	501	MRK	C21-N17	-4.65	1.29	1.37
3	A	501	MRK	C18-N17	-2.19	1.34	1.36
4	A	600	ANP	PG-O2G	-2.03	1.51	1.56
4	A	600	ANP	PG-O1G	3.08	1.49	1.46
3	A	501	MRK	C20-N19	5.34	1.62	1.37
4	A	600	ANP	PB-O1B	6.03	1.53	1.46
3	A	501	MRK	C21-C20	8.15	1.67	1.36

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	600	ANP	N3-C2-N1	-9.36	120.71	128.86
3	A	501	MRK	C21-C20-N19	-7.55	90.28	109.38
4	A	600	ANP	O1G-PG-N3B	-4.89	104.48	111.79
4	A	600	ANP	O1B-PB-N3B	-4.14	105.60	111.79
3	A	501	MRK	C15-C13-C12	-3.98	119.30	122.69
4	A	600	ANP	C4-C5-N7	-3.00	106.51	109.41
3	A	501	MRK	C22-N17-C21	-2.81	119.57	124.89
4	A	600	ANP	PA-O3A-PB	-2.68	122.91	132.38
3	A	501	MRK	F14-C13-C12	2.02	122.11	118.44
3	A	501	MRK	C22-N17-C18	2.27	127.78	125.62
3	A	501	MRK	C21-N17-C18	4.25	112.61	108.56
4	A	600	ANP	O2B-PB-O1B	4.42	119.06	109.87
3	A	501	MRK	C9-C10-C7	4.83	123.24	120.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	MRK	1	0
4	A	600	ANP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	450/470 (95%)	0.06	25 (5%)	25 24	17, 31, 52, 65	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68	GLY	4.6
1	A	195	PHE	4.4
1	A	93	GLU	4.2
1	A	6	LEU	4.0
1	A	82	THR	3.9
1	A	141	HIS	3.9
1	A	8	PHE	3.6
1	A	94	GLY	3.3
1	A	194	ASP	3.3
1	A	398	SER	3.2
1	A	139	MET	2.9
1	A	98	GLN	2.8
1	A	67	GLU	2.7
1	A	138	GLN	2.7
1	A	196	GLU	2.6
1	A	216	GLU	2.5
1	A	69	SER	2.5
1	A	399	GLU	2.4
1	A	99	TRP	2.3
1	A	5	ASN	2.3
1	A	401	VAL	2.2
1	A	70	GLU	2.1
1	A	349	TYR	2.0
1	A	142	LYS	2.0
1	A	346	LYS	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	ANP	A	600	31/31	0.81	0.29	7.00	19,28,35,35	31
2	GLC	A	500	12/12	0.98	0.11	-0.42	16,20,21,24	0
3	MRK	A	501	23/23	0.96	0.12	-1.35	32,35,38,39	0
6	K	A	800	1/1	0.99	0.07	-1.53	27,27,27,27	0
5	MG	A	700	1/1	0.90	0.14	-	42,42,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.