



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 12:01 pm GMT

PDB ID : 4IFV
Title : Detecting Allosteric Sites of HIV-1 Reverse Transcriptase by X-Ray Crystallographic Fragment Screening
Authors : Bauman, J.D.; Patel, D.; Fromer, M.; Arnold, E.
Deposited on : 2012-12-15
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1316 (2.04-2.04)
Clashscore	112137	1394 (2.04-2.04)
Ramachandran outliers	110173	1383 (2.04-2.04)
Sidechain outliers	110143	1383 (2.04-2.04)
RSRZ outliers	101464	1319 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	<div> <div>6%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
2	B	429	<div> <div>11%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>••</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMQ	A	602	-	-	-	X
4	FMQ	B	501	-	-	-	X
5	DMS	A	603	-	-	-	X
5	DMS	B	503	-	-	-	X
5	DMS	B	504	-	-	-	X
5	DMS	B	508	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exoribonuclease H, p66 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	556	Total	C	N	O	S	0	1	0
			4525	2928	752	837	8			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP P03366
A	0	VAL	-	EXPRESSION TAG	UNP P03366
A	172	ALA	LYS	ENGINEERED MUTATION	UNP P03366
A	173	ALA	LYS	ENGINEERED MUTATION	UNP P03366
A	280	SER	CYS	ENGINEERED MUTATION	UNP P03366


- Molecule 2 is a protein called Gag-Pol polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	412	Total	C	N	O	S	6	1	0
			3419	2228	565	619	7			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	EXPRESSION TAG	UNP P03366
B	280	SER	CYS	ENGINEERED MUTATION	UNP P03366

- Molecule 3 is 4-{[4-({4-[(E)-2-CYANOETHENYL]-2,6-DIMETHYLPHENYL}AMINO)PYRIMIDIN-2-YL]AMINO}BENZONITRILE (three-letter code: T27) (formula: C₂₂H₁₈N₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N		0	0
			28	22	6			

- Molecule 4 is [1-(4-FLUOROPHENYL)-5-METHYL-1H-PYRAZOL-4-YL]METHANOL (three-letter code: FMQ) (formula: C₁₁H₁₁FN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			15	11	1	2	1		
4	B	1	Total	C	F	N	O	0	0
			15	11	1	2	1		

- Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Mg 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	380	Total 380	O 380	0	0
7	B	236	Total 236	O 236	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Exoribonuclease H, p66 RT



• Molecule 2: Gag-Pol polyprotein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.06Å 72.82Å 108.97Å 90.00° 100.43° 90.00°	Depositor
Resolution (Å)	42.92 – 2.05 42.92 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.92-2.05) 92.9 (42.92-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1233)	Depositor
R, R_{free}	0.188 , 0.219 0.185 , 0.219	Depositor DCC
R_{free} test set	1729 reflections (2.21%)	DCC
Wilson B-factor (Å ²)	31.4	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8663	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DMS, T27, FMQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/4643	0.48	0/6311
2	B	0.28	0/3517	0.50	1/4776 (0.0%)
All	All	0.28	0/8160	0.49	1/11087 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	TRP	N-CA-C	5.17	124.96	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4525	0	4579	42	1
2	B	3419	0	3447	45	0
3	A	28	0	18	1	0
4	A	15	0	11	1	0
4	B	15	0	11	0	0
5	A	16	0	24	7	0
5	B	28	0	42	6	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	380	0	0	8	0
7	B	236	0	0	12	0
All	All	8663	0	8132	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:26:LEU:H	5:B:504:DMS:H13	1.23	1.00
2:B:180:ILE:O	7:B:798:HOH:O	1.85	0.95
2:B:360:ALA:HB2	2:B:366:LYS:HD2	1.62	0.80
2:B:185:ASP:OD1	7:B:806:HOH:O	1.99	0.79
1:A:220:LYS:HG2	1:A:221:HIS:H	1.50	0.74
1:A:426:TRP:H	5:A:603:DMS:H22	1.59	0.68
2:B:358:ARG:NH1	2:B:370:GLU:OE2	2.27	0.67
2:B:89:GLU:N	2:B:89:GLU:OE1	2.27	0.66
1:A:461:LYS:NZ	7:A:927:HOH:O	2.08	0.63
1:A:543:GLY:HA3	2:B:283:LEU:O	1.99	0.62
2:B:358:ARG:NH1	7:B:823:HOH:O	2.31	0.62
1:A:474:ASN:H	5:A:606:DMS:H22	1.64	0.61
1:A:428:GLN:H	5:A:604:DMS:H23	1.67	0.60
2:B:30:LYS:NZ	2:B:404:GLU:OE2	2.32	0.59
2:B:166:LYS:NZ	7:B:749:HOH:O	2.28	0.59
5:B:504:DMS:H12	7:B:810:HOH:O	2.04	0.56
3:A:601:T27:N3	3:A:601:T27:H15	2.24	0.53
2:B:108:VAL:HB	2:B:232:TYR:HB3	1.90	0.53
2:B:358:ARG:NH2	7:B:823:HOH:O	2.41	0.53
2:B:358:ARG:CZ	7:B:823:HOH:O	2.57	0.52
1:A:253:THR:HG22	1:A:292:VAL:HG22	1.92	0.52
2:B:64:LYS:HE3	2:B:71:TRP:CE2	2.45	0.52
1:A:428:GLN:H	5:A:604:DMS:C2	2.24	0.51
1:A:450:THR:O	7:A:1028:HOH:O	2.19	0.51
1:A:7:THR:HG22	1:A:119:PRO:HB2	1.92	0.51
1:A:181:TYR:CE2	1:A:183:TYR:HB2	2.46	0.50
2:B:424:LYS:HE3	7:B:809:HOH:O	2.12	0.50
1:A:254:VAL:HG23	1:A:293:ILE:HD11	1.92	0.50
2:B:239:TRP:HE1	5:B:505:DMS:C2	2.25	0.50
1:A:68:SER:HB3	1:A:70:LYS:HG3	1.94	0.50
2:B:65:LYS:HA	2:B:407:GLN:OE1	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:TRP:H	5:A:603:DMS:C2	2.25	0.49
2:B:29:GLU:OE2	7:B:735:HOH:O	2.20	0.49
2:B:104:LYS:HE3	2:B:192:ASP:HB3	1.94	0.49
2:B:65:LYS:HD2	2:B:72:ARG:HD2	1.93	0.49
1:A:199:ARG:HH22	1:A:220:LYS:HE3	1.78	0.49
2:B:195:ILE:HA	2:B:198:HIS:HB3	1.96	0.48
1:A:542:ILE:HG12	2:B:283:LEU:HD12	1.95	0.48
2:B:115:TYR:OH	2:B:182:GLN:NE2	2.43	0.48
2:B:209:LEU:O	2:B:212:TRP:HB3	2.13	0.48
1:A:199:ARG:NH2	1:A:220:LYS:HE3	2.29	0.47
1:A:115:TYR:HA	1:A:160:PHE:CE1	2.49	0.47
1:A:208:HIS:O	1:A:211:ARG:HG2	2.15	0.47
1:A:334:GLN:H	5:A:605:DMS:C1	2.27	0.47
1:A:90:VAL:HG11	2:B:140:PRO:HB3	1.97	0.47
1:A:428:GLN:NE2	7:A:1073:HOH:O	2.29	0.47
1:A:454:LYS:HZ2	1:A:554:ALA:HA	1.80	0.47
2:B:22:LYS:HD2	7:B:783:HOH:O	2.15	0.47
1:A:369:THR:O	1:A:373[B]:GLN:HG2	2.15	0.47
2:B:241:VAL:HG23	2:B:243:PRO:HD3	1.97	0.46
1:A:388:LYS:HE3	7:A:978:HOH:O	2.16	0.46
1:A:417:VAL:HG22	1:A:419:THR:HG23	1.97	0.46
2:B:235:HIS:H	5:B:505:DMS:H23	1.80	0.46
2:B:151:GLN:HB3	2:B:185:ASP:OD2	2.16	0.46
1:A:354:TYR:HD1	1:A:374:LYS:HD2	1.82	0.44
1:A:286:THR:OG1	1:A:287:LYS:N	2.51	0.44
2:B:89:GLU:HG2	2:B:90:VAL:H	1.82	0.44
1:A:68:SER:HB3	1:A:70:LYS:H	1.83	0.43
2:B:26:LEU:N	5:B:504:DMS:H13	2.08	0.43
2:B:374:LYS:HA	2:B:374:LYS:HD2	1.77	0.43
2:B:428:GLN:NE2	7:B:777:HOH:O	2.51	0.43
1:A:396:GLU:HG3	7:A:984:HOH:O	2.18	0.43
1:A:480:GLN:HG2	4:A:602:FMQ:C11	2.49	0.43
1:A:536:VAL:HB	1:A:542:ILE:HD13	2.00	0.43
1:A:363:ASN:HA	1:A:511:ASP:OD1	2.19	0.42
2:B:236:PRO:HA	2:B:239:TRP:CD2	2.55	0.42
2:B:255:ASN:O	2:B:259:LYS:HG2	2.20	0.42
1:A:334:GLN:H	5:A:605:DMS:H11	1.83	0.42
1:A:220:LYS:HB2	1:A:220:LYS:HE2	1.87	0.42
2:B:89:GLU:CG	2:B:90:VAL:H	2.33	0.42
1:A:287:LYS:HD3	1:A:287:LYS:HA	1.85	0.42
2:B:103:LYS:HA	2:B:103:LYS:HD3	1.89	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:281:LYS:O	2:B:284:ARG:HG2	2.20	0.42
2:B:70:LYS:HD2	2:B:70:LYS:HA	1.88	0.42
2:B:210:LEU:HD23	2:B:214:LEU:HD22	2.01	0.41
1:A:358:ARG:HD3	1:A:358:ARG:HA	1.85	0.41
2:B:107:THR:OG1	2:B:198:HIS:NE2	2.49	0.41
2:B:383:TRP:O	7:B:836:HOH:O	2.22	0.41
1:A:245:VAL:N	7:A:929:HOH:O	1.99	0.41
1:A:271:TYR:HA	1:A:272:PRO:HD3	1.87	0.41
1:A:535:TRP:CZ3	2:B:426:TRP:HH2	2.37	0.41
1:A:308:GLU:OE1	7:A:995:HOH:O	2.22	0.41
2:B:235:HIS:H	5:B:505:DMS:C2	2.34	0.41
1:A:448:ARG:NH1	7:A:1012:HOH:O	2.53	0.40
2:B:54:ASN:HA	2:B:55:PRO:HD2	1.96	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLU:OE2	1:A:448:ARG:NH2[4_546]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	555/557 (100%)	548 (99%)	7 (1%)	0	100	100
2	B	409/429 (95%)	403 (98%)	5 (1%)	1 (0%)	51	42
All	All	964/986 (98%)	951 (99%)	12 (1%)	1 (0%)	55	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	89	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/495 (100%)	488 (98%)	8 (2%)	68	65
2	B	375/390 (96%)	368 (98%)	7 (2%)	62	58
All	All	871/885 (98%)	856 (98%)	15 (2%)	66	63

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	GLU
1	A	64	LYS
1	A	67	ASP
1	A	219	LYS
1	A	221	HIS
1	A	290	THR
1	A	503	LEU
1	A	512	LYS
2	B	68	SER
2	B	90	VAL
2	B	208	HIS
2	B	307	ARG
2	B	356	ARG
2	B	374	LYS
2	B	410	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	T27	A	601	-	30,30,30	1.12	3 (10%)	38,40,40	1.84	4 (10%)
4	FMQ	A	602	-	15,16,16	2.04	5 (33%)	18,22,22	1.66	3 (16%)
5	DMS	A	603	-	3,3,3	0.56	0	3,3,3	0.44	0
5	DMS	A	604	-	3,3,3	0.54	0	3,3,3	0.52	0
5	DMS	A	605	-	3,3,3	0.60	0	3,3,3	0.47	0
5	DMS	A	606	-	3,3,3	0.66	0	3,3,3	0.56	0
4	FMQ	B	501	-	15,16,16	2.05	5 (33%)	18,22,22	1.64	3 (16%)
5	DMS	B	502	-	3,3,3	0.64	0	3,3,3	0.49	0
5	DMS	B	503	-	3,3,3	0.63	0	3,3,3	0.40	0
5	DMS	B	504	-	3,3,3	0.58	0	3,3,3	0.49	0
5	DMS	B	505	-	3,3,3	0.61	0	3,3,3	0.46	0
5	DMS	B	506	-	3,3,3	0.65	0	3,3,3	0.50	0
5	DMS	B	507	-	3,3,3	0.66	0	3,3,3	0.56	0
5	DMS	B	508	-	3,3,3	0.60	0	3,3,3	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	T27	A	601	-	-	0/13/14/14	0/3/3/3
4	FMQ	A	602	-	-	0/4/6/6	0/2/2/2
5	DMS	A	603	-	-	0/0/0/0	0/0/0/0
5	DMS	A	604	-	-	0/0/0/0	0/0/0/0
5	DMS	A	605	-	-	0/0/0/0	0/0/0/0
5	DMS	A	606	-	-	0/0/0/0	0/0/0/0
4	FMQ	B	501	-	-	0/4/6/6	0/2/2/2
5	DMS	B	502	-	-	0/0/0/0	0/0/0/0
5	DMS	B	503	-	-	0/0/0/0	0/0/0/0
5	DMS	B	504	-	-	0/0/0/0	0/0/0/0
5	DMS	B	505	-	-	0/0/0/0	0/0/0/0
5	DMS	B	506	-	-	0/0/0/0	0/0/0/0
5	DMS	B	507	-	-	0/0/0/0	0/0/0/0
5	DMS	B	508	-	-	0/0/0/0	0/0/0/0

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	FMQ	F12-C9	-4.24	1.26	1.36
4	A	602	FMQ	N4-N3	-4.19	1.32	1.39
4	A	602	FMQ	F12-C9	-4.11	1.26	1.36
4	B	501	FMQ	N4-N3	-4.04	1.32	1.39
4	B	501	FMQ	C10-C9	-2.24	1.32	1.37
4	B	501	FMQ	C2-C1	-2.20	1.35	1.39
4	A	602	FMQ	C10-C9	-2.07	1.33	1.37
4	A	602	FMQ	C2-C1	-2.04	1.35	1.39
3	A	601	T27	C13-C19	2.06	1.49	1.44
3	A	601	T27	C11-N1	3.14	1.44	1.38
4	B	501	FMQ	C8-C9	3.19	1.43	1.37
4	A	602	FMQ	C8-C9	3.19	1.43	1.37
3	A	601	T27	C12-N4	3.97	1.44	1.36

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	T27	C10-C9-N2	-5.13	118.01	123.92
3	A	601	T27	N2-C12-N3	-4.69	122.17	126.68
4	A	602	FMQ	C6-N3-C2	-3.76	126.20	130.09
4	B	501	FMQ	C2-N3-N4	-2.86	108.47	112.02
4	B	501	FMQ	C14-C1-C5	-2.61	124.00	127.67
4	A	602	FMQ	C2-N3-N4	-2.36	109.09	112.02
3	A	601	T27	C10-C11-N3	-2.36	119.27	123.21
4	A	602	FMQ	C5-N4-N3	3.70	108.68	103.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	FMQ	C5-N4-N3	4.35	109.51	103.92
3	A	601	T27	C9-N2-C12	7.11	121.36	115.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	T27	1	0
4	A	602	FMQ	1	0
5	A	603	DMS	2	0
5	A	604	DMS	2	0
5	A	605	DMS	2	0
5	A	606	DMS	1	0
5	B	504	DMS	3	0
5	B	505	DMS	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	556/557 (99%)	0.04	34 (6%) 22 24	19, 39, 88, 129	0
2	B	412/429 (96%)	0.46	47 (11%) 6 5	20, 43, 97, 148	0
All	All	968/986 (98%)	0.22	81 (8%) 12 12	19, 41, 93, 148	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	92	LEU	12.2
2	B	90	VAL	10.0
1	A	67	ASP	8.5
2	B	89	GLU	7.7
2	B	93	GLY	7.1
2	B	88	TRP	6.9
1	A	221	HIS	6.7
1	A	69	THR	6.4
1	A	289	LEU	5.6
1	A	286	THR	5.3
1	A	220	LYS	5.3
1	A	218	ASP	5.2
2	B	214	LEU	5.0
2	B	95	PRO	4.7
1	A	65	LYS	4.6
2	B	6	GLU	4.5
2	B	357	MET	4.4
1	A	68	SER	4.3
1	A	72	ARG	4.2
2	B	195	ILE	4.2
2	B	66	LYS	4.1
1	A	66	LYS	4.0
2	B	91	GLN	4.0
2	B	94	ILE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	288	ALA	3.9
1	A	91	GLN	3.9
1	A	222	GLN	3.9
2	B	168	LEU	3.9
1	A	287	LYS	3.8
2	B	193	LEU	3.8
2	B	356	ARG	3.8
1	A	219	LYS	3.7
1	A	24	TRP	3.5
2	B	358	ARG	3.5
2	B	7	THR	3.5
2	B	359	GLY	3.5
2	B	212	TRP	3.4
2	B	67	ASP	3.4
2	B	209	LEU	3.3
2	B	5	ILE	3.3
2	B	284	ARG	3.3
1	A	70	LYS	3.2
2	B	69	THR	3.2
2	B	229	TRP	3.2
1	A	74	LEU	3.1
1	A	285	GLY	3.0
2	B	196	GLY	3.0
1	A	64	LYS	3.0
1	A	290	THR	2.9
2	B	197	GLN	2.9
2	B	166	LYS	2.9
1	A	292	VAL	2.9
2	B	87	PHE	2.9
2	B	202	ILE	2.7
2	B	180	ILE	2.7
1	A	71	TRP	2.7
2	B	200	THR	2.7
2	B	173	LYS	2.7
1	A	358	ARG	2.6
2	B	198	HIS	2.6
2	B	232	TYR	2.5
2	B	171	PHE	2.5
2	B	361	HIS	2.4
1	A	554	ALA	2.4
2	B	213	GLY	2.4
2	B	231	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	174	GLN	2.4
1	A	546	GLU	2.3
1	A	284	ARG	2.3
1	A	115	TYR	2.3
2	B	194	GLU	2.2
2	B	422	LEU	2.2
2	B	426	TRP	2.2
1	A	92	LEU	2.2
1	A	282	LEU	2.2
1	A	357	MET	2.1
2	B	170	PRO	2.1
2	B	204	GLU	2.0
2	B	211	ARG	2.0
1	A	52	PRO	2.0
1	A	254	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	DMS	B	508	4/4	0.84	0.28	12.28	35,59,108,109	0
5	DMS	A	603	4/4	0.92	0.26	7.59	19,44,85,87	0
4	FMQ	B	501	15/15	0.73	0.26	6.05	42,51,57,57	0
4	FMQ	A	602	15/15	0.81	0.22	3.81	51,55,66,71	0
5	DMS	B	503	4/4	0.94	0.19	3.71	36,56,60,63	0
5	DMS	B	504	4/4	0.88	0.15	2.06	21,27,58,59	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	T27	A	601	28/28	0.95	0.13	1.82	22,28,35,38	0
5	DMS	A	604	4/4	0.96	0.13	1.80	28,42,42,46	0
5	DMS	B	506	4/4	0.92	0.26	1.65	70,78,81,98	0
5	DMS	B	502	4/4	0.98	0.12	-0.26	39,50,53,53	0
5	DMS	B	505	4/4	0.91	0.14	-0.35	33,78,79,84	0
6	MG	A	607	1/1	0.79	0.10	-1.11	55,55,55,55	0
5	DMS	B	507	4/4	0.95	0.10	-1.25	42,42,55,92	0
5	DMS	A	606	4/4	0.86	0.17	-	65,86,88,90	0
5	DMS	A	605	4/4	0.78	0.20	-	32,58,84,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.