



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 3, 2018 – 10:56 AM EST

PDB ID : 5ILT
Title : Crystal structure of bovine Fab A01
Authors : Stanfield, R.L.; Wilson, I.A.
Deposited on : 2016-03-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	
2	H	271	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7088 atoms, of which 3459 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called bovine Fab A01 light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	213	Total	C	H	N	O	S	0	0	0
			3063	959	1502	263	334	5			

- Molecule 2 is a protein called bovine Fab A01 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	271	Total	C	H	N	O	S	0	5	0
			3988	1255	1957	339	418	19			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	16	Total	O	0	0
			16	16		
3	H	21	Total	O	0	0
			21	21		

- Molecule 1: bovine Fab A01 light chain



- Molecule 2: bovine Fab A01 heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.49Å 72.49Å 87.42Å 90.00° 107.31° 90.00°	Depositor
Resolution (Å)	38.42 – 2.00 38.42 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (38.42-2.00) 98.2 (38.42-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.211 , 0.254 0.211 , 0.254	Depositor DCC
R_{free} test set	1883 reflections (5.90%)	DCC
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7088	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.26	0/1591	0.47	0/2168
2	H	0.27	0/2091	0.49	0/2854
All	All	0.26	0/3682	0.48	0/5022

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1561	1502	1502	2	0
2	H	2031	1957	1957	9	0
3	H	21	0	0	0	0
3	L	16	0	0	0	0
All	All	3629	3459	3459	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:186:SER:HB2	2:H:187:SER:HA	1.85	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:83:GLU:OE1	1:L:166:LYS:NZ	2.26	0.54
2:H:34:VAL:HG21	2:H:78:VAL:HG21	1.91	0.52
2:H:125:CYS:N	2:H:126:GLY:HA3	2.29	0.47
2:H:186:SER:CB	2:H:187:SER:CA	2.95	0.43
2:H:121:ASP:OD1	2:H:122:ALA:N	2.51	0.43
2:H:120[B]:CYS:HB3	2:H:139:CYS:SG	2.59	0.43
2:H:186:SER:HB3	2:H:187:SER:C	2.40	0.42
2:H:186:SER:HB2	2:H:187:SER:CA	2.48	0.42
1:L:119:PRO:HG3	2:H:180:CYS:SG	2.60	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	211/216 (98%)	202 (96%)	9 (4%)	0	100	100
2	H	274/271 (101%)	263 (96%)	11 (4%)	0	100	100
All	All	485/487 (100%)	465 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	182/184 (99%)	179 (98%)	3 (2%)	68	72
2	H	243/238 (102%)	241 (99%)	2 (1%)	85	88
All	All	425/422 (101%)	420 (99%)	5 (1%)	75	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	30	ASN
1	L	56	SER
1	L	206	VAL
2	H	143	ASP
2	H	184	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/216 (98%)	0.11	1 (0%) 90 90	29, 52, 77, 104	0
2	H	271/271 (100%)	0.71	27 (9%) 8 8	28, 56, 119, 194	0
All	All	484/487 (99%)	0.45	28 (5%) 24 24	28, 54, 111, 194	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	268	CYS	15.0
2	H	241	THR	6.3
2	H	242	SER	5.6
2	H	182	GLY	4.8
2	H	125	CYS	4.2
2	H	180	CYS	4.2
2	H	110	ILE	3.7
2	H	211	ALA	3.7
2	H	109	TRP	3.5
2	H	142	LEU	3.4
2	H	128	LEU	3.2
2	H	114	SER	3.1
2	H	243	GLY	3.0
2	H	105	ASP	3.0
2	H	240	SER	2.8
2	H	115	SER	2.8
2	H	188	THR	2.8
2	H	127	ASN	2.7
2	H	184	LYS	2.7
2	H	145	SER	2.5
2	H	111[A]	CYS	2.5
2	H	117	THR	2.5
2	H	1	GLN	2.5
2	H	129	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	186	SER	2.4
2	H	213	LYS	2.2
1	L	211	CYS	2.1
2	H	132	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.