



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2017 – 11:58 PM EDT

PDB ID : 3INM
Title : Crystal structure of human cytosolic NADP(+)-dependent isocitrate dehydrogenase R132H mutant in complex with NADPH, ALPHA-KETOGLUTARATE and CALCIUM(2+)
Authors : Fontano, E.; Brown, R.S.; Suto, R.K.; Bhyravbhatla, B.
Deposited on : unknown
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	<div> <div>3%</div> <div>73%</div> <div>22%</div> <div>• •</div> </div>
1	B	425	<div> <div>%</div> <div>75%</div> <div>19%</div> <div>• •</div> </div>
1	C	425	<div> <div>3%</div> <div>67%</div> <div>26%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AKG	C	511	-	-	-	X
6	GOL	A	531	-	-	X	X
6	GOL	A	532	-	-	-	X
6	GOL	C	532	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP] cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	0	0
			3225	2051	545	611	18			
1	B	411	Total	C	N	O	S	0	0	0
			3256	2071	551	616	18			
1	C	407	Total	C	N	O	S	0	0	0
			3225	2051	545	611	18			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	HIS	ARG	ENGINEERED	UNP O75874
A	415	SER	-	EXPRESSION TAG	UNP O75874
A	416	LEU	-	EXPRESSION TAG	UNP O75874
A	417	GLU	-	EXPRESSION TAG	UNP O75874
A	418	HIS	-	EXPRESSION TAG	UNP O75874
A	419	HIS	-	EXPRESSION TAG	UNP O75874
A	420	HIS	-	EXPRESSION TAG	UNP O75874
A	421	HIS	-	EXPRESSION TAG	UNP O75874
A	422	HIS	-	EXPRESSION TAG	UNP O75874
A	423	HIS	-	EXPRESSION TAG	UNP O75874
A	424	HIS	-	EXPRESSION TAG	UNP O75874
A	425	HIS	-	EXPRESSION TAG	UNP O75874
B	132	HIS	ARG	ENGINEERED	UNP O75874
B	415	SER	-	EXPRESSION TAG	UNP O75874
B	416	LEU	-	EXPRESSION TAG	UNP O75874
B	417	GLU	-	EXPRESSION TAG	UNP O75874
B	418	HIS	-	EXPRESSION TAG	UNP O75874
B	419	HIS	-	EXPRESSION TAG	UNP O75874
B	420	HIS	-	EXPRESSION TAG	UNP O75874
B	421	HIS	-	EXPRESSION TAG	UNP O75874
B	422	HIS	-	EXPRESSION TAG	UNP O75874
B	423	HIS	-	EXPRESSION TAG	UNP O75874
B	424	HIS	-	EXPRESSION TAG	UNP O75874


Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	425	HIS	-	EXPRESSION TAG	UNP O75874
C	132	HIS	ARG	ENGINEERED	UNP O75874
C	415	SER	-	EXPRESSION TAG	UNP O75874
C	416	LEU	-	EXPRESSION TAG	UNP O75874
C	417	GLU	-	EXPRESSION TAG	UNP O75874
C	418	HIS	-	EXPRESSION TAG	UNP O75874
C	419	HIS	-	EXPRESSION TAG	UNP O75874
C	420	HIS	-	EXPRESSION TAG	UNP O75874
C	421	HIS	-	EXPRESSION TAG	UNP O75874
C	422	HIS	-	EXPRESSION TAG	UNP O75874
C	423	HIS	-	EXPRESSION TAG	UNP O75874
C	424	HIS	-	EXPRESSION TAG	UNP O75874
C	425	HIS	-	EXPRESSION TAG	UNP O75874

- # NDP

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	C	1	Total 48	C 21	N 7	O 17	P 3	0	0

- 
- WORLD WIDE
PDB
PROTEIN DATA BANK



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		
5	A	1	Total	Na	0	0
			1	1		
5	C	1	Total	Na	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	112	Total	O	0	0
			112	112		
7	B	112	Total	O	0	0
			112	112		
7	C	77	Total	O	0	0
			77	77		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



E361	V362	S363	I367	E368	K372	D375	I380	K381	G382	L383	P384	N385	V386	Q387	R388	L392	E396	F397	D399	K400	N404	A410	G411	ALA	LYS	LEU	SER	LEU	GLU	HIS	HIS	HIS	HIS	HIS	HIS	HIS										
R249	D252	D253	Q257	A258	S261	L266	Y272	D273	G274	G283	G284	Y285	M291	T292	S293	E306	T313	R314	H315	Y316	R317	M318	Y319	Q320	K321	E324	I330	F334	R338	R343	A344	D347	N348	R349	K350	E351	F354	N357	A358	L359	E360					
L120	V121	S122	G123	W124	H132	V146	P147	K151	V152	E153	T155	P158	S159	D160	G161	T162	Q163	K164	M171	F172	E173	E174	M182	Q185	S188	I189	H194	F197	Q198	M199	A200	L201	S202	K203	N213	K217	E229	I230	Y231	Y235	K236	S237	K243			
MET	SER	LYS	K4	S9	E12	M13	Q14	E17	M18	T19	R20	L25	L26	K27	E28	K29	P33	L39	E47	K58	I64	K65	H67	N68	V69	K72	T75	E84	K89	Q90	M91	R100	L103	F108	R109	I112	I113	G114	K115	R119						

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	96.12Å 274.69Å 116.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.91 – 2.10 24.91 – 2.10	Depositor EDS
% Data completeness (in resolution range)	92.4 (24.91-2.10) 92.4 (24.91-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.10Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.220 , 0.262 0.215 , 0.251	Depositor DCC
R_{free} test set	4153 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10205	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, AKG, NDP, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/3294	0.55	0/4444
1	B	0.79	1/3325 (0.0%)	0.58	1/4485 (0.0%)
1	C	0.80	3/3294 (0.1%)	0.57	0/4444
All	All	0.76	4/9913 (0.0%)	0.57	1/13373 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	69	VAL	CB-CG2	-5.79	1.40	1.52
1	C	174	GLU	CB-CG	5.55	1.62	1.52
1	B	262	GLU	CG-CD	5.10	1.59	1.51
1	C	272	TYR	CD1-CE1	-5.08	1.31	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	GLY	N-CA-C	5.40	126.60	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3225	0	3183	85	0
1	B	3256	0	3220	71	0
1	C	3225	0	3183	134	0
2	A	48	0	26	1	0
2	B	48	0	26	5	0
2	C	48	0	26	4	0
3	A	10	0	4	0	0
3	B	10	0	4	1	0
3	C	10	0	4	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	A	12	0	16	13	0
6	C	6	0	8	2	0
7	A	112	0	0	1	0
7	B	112	0	0	1	0
7	C	77	0	0	7	0
All	All	10205	0	9700	286	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 286 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:395:PHE:CD1	6:A:531:GOL:H12	1.46	1.47
1:C:388:ARG:HD3	1:C:388:ARG:O	1.20	1.32
1:A:388:ARG:HD2	1:A:388:ARG:O	1.16	1.29
1:A:395:PHE:HD1	6:A:531:GOL:C1	1.47	1.26
1:B:388:ARG:CD	1:B:388:ARG:O	1.89	1.21

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/425 (95%)	391 (96%)	14 (4%)	0	100	100
1	B	409/425 (96%)	399 (98%)	10 (2%)	0	100	100
1	C	405/425 (95%)	388 (96%)	17 (4%)	0	100	100
All	All	1219/1275 (96%)	1178 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/361 (95%)	323 (94%)	21 (6%)	22	18
1	B	347/361 (96%)	322 (93%)	25 (7%)	17	13
1	C	344/361 (95%)	318 (92%)	26 (8%)	15	11
All	All	1035/1083 (96%)	963 (93%)	72 (7%)	18	14

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	185	GLN
1	B	303	VAL
1	C	350	LYS
1	B	201	LEU
1	B	222	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	323	GLN
1	C	90	GLN
1	C	194	HIS
1	B	194	HIS
1	C	277	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	501	-	43,52,52	1.47	4 (9%)	49,80,80	1.60	2 (4%)
3	AKG	A	511	-	3,9,9	0.35	0	4,11,11	1.85	1 (25%)
6	GOL	A	531	-	5,5,5	0.50	0	5,5,5	0.32	0
6	GOL	A	532	-	5,5,5	0.40	0	5,5,5	0.60	0
2	NDP	B	501	-	43,52,52	1.49	3 (6%)	49,80,80	1.60	1 (2%)
3	AKG	B	511	-	3,9,9	0.42	0	4,11,11	1.37	1 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	C	501	-	43,52,52	1.33	4 (9%)	49,80,80	1.75	2 (4%)
3	AKG	C	511	-	3,9,9	0.43	0	4,11,11	1.87	1 (25%)
6	GOL	C	532	-	5,5,5	0.67	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	501	-	-	0/30/77/77	0/5/5/5
3	AKG	A	511	-	-	0/3/9/9	0/0/0/0
6	GOL	A	531	-	-	0/4/4/4	0/0/0/0
6	GOL	A	532	-	-	0/4/4/4	0/0/0/0
2	NDP	B	501	-	-	0/30/77/77	0/5/5/5
3	AKG	B	511	-	-	0/3/9/9	0/0/0/0
2	NDP	C	501	-	-	0/30/77/77	0/5/5/5
3	AKG	C	511	-	-	0/3/9/9	0/0/0/0
6	GOL	C	532	-	-	0/4/4/4	0/0/0/0

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	NDP	C2A-N1A	2.00	1.37	1.33
2	C	501	NDP	C2A-N3A	2.11	1.35	1.32
2	A	501	NDP	C2A-N1A	2.72	1.39	1.33
2	B	501	NDP	C2A-N3A	3.09	1.37	1.32
2	C	501	NDP	C6N-C5N	3.30	1.39	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	NDP	N3A-C2A-N1A	-10.76	119.48	128.86
2	B	501	NDP	N3A-C2A-N1A	-9.95	120.19	128.86
2	A	501	NDP	N3A-C2A-N1A	-9.77	120.35	128.86
3	C	511	AKG	C3-C4-C5	-3.23	107.15	112.66
3	A	511	AKG	C3-C4-C5	-3.17	107.25	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NDP	1	0
6	A	531	GOL	13	0
2	B	501	NDP	5	0
3	B	511	AKG	1	0
2	C	501	NDP	4	0
3	C	511	AKG	2	0
6	C	532	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/425 (95%)	0.16	14 (3%) 46 53	19, 32, 45, 54	0
1	B	411/425 (96%)	0.04	5 (1%) 79 82	19, 29, 42, 50	0
1	C	407/425 (95%)	0.26	13 (3%) 48 55	23, 34, 49, 59	0
All	All	1225/1275 (96%)	0.15	32 (2%) 56 62	19, 32, 45, 59	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	174	GLU	4.0
1	A	71	VAL	3.3
1	A	205	TRP	3.2
1	C	383	LEU	3.1
1	C	354	PHE	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GOL	A	532	6/6	0.86	0.26	5.79	28,33,36,43	0
6	GOL	A	531	6/6	0.79	0.40	5.57	29,34,36,37	0
6	GOL	C	532	6/6	0.75	0.30	5.51	32,34,35,36	0
3	AKG	C	511	10/10	0.89	0.15	2.54	38,44,48,49	0
3	AKG	A	511	10/10	0.92	0.14	1.67	35,41,45,47	0
3	AKG	B	511	10/10	0.93	0.14	0.83	32,37,41,45	0
4	CA	C	521	1/1	0.99	0.12	0.23	41,41,41,41	0
2	NDP	C	501	48/48	0.97	0.12	-0.36	22,29,34,37	0
2	NDP	B	501	48/48	0.97	0.11	-0.47	19,24,32,37	0
2	NDP	A	501	48/48	0.96	0.12	-0.51	20,28,36,40	0
4	CA	B	521	1/1	0.98	0.07	-2.75	33,33,33,33	0
4	CA	A	521	1/1	0.97	0.04	-3.88	35,35,35,35	0
5	NA	C	522	1/1	0.95	0.11	-	30,30,30,30	0
5	NA	B	522	1/1	0.88	0.14	-	30,30,30,30	0
5	NA	A	522	1/1	0.95	0.09	-	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.