



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:40 pm GMT

PDB ID : 2IOL  
Title : Crystal structure of the C-terminal MA3 domain of Pdcd4 (mouse); form 1  
Authors : Wlodawer, A.; LaRonde-LeBlanc, N.A.  
Deposited on : 2006-10-10  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	<div> <div>9%</div> <div>69%</div> <div>13%</div> <div>•</div> <div>15%</div> </div>
1	B	150	<div> <div>23%</div> <div>61%</div> <div>20%</div> <div>•</div> <div>15%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Programmed Cell Death 4, Pdcd4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	Se	0	0	0
			1030	662	166	195	3	4			
1	B	127	Total	C	N	O	S	Se	0	0	0
			1030	662	166	195	3	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	333	MSE	MET	MODIFIED RESIDUE	UNP Q61823
A	370	MSE	MET	MODIFIED RESIDUE	UNP Q61823
A	382	MSE	MET	MODIFIED RESIDUE	UNP Q61823
A	401	MSE	MET	MODIFIED RESIDUE	UNP Q61823
B	333	MSE	MET	MODIFIED RESIDUE	UNP Q61823
B	370	MSE	MET	MODIFIED RESIDUE	UNP Q61823
B	382	MSE	MET	MODIFIED RESIDUE	UNP Q61823
B	401	MSE	MET	MODIFIED RESIDUE	UNP Q61823


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total	O	0	0
			36	36		
2	B	14	Total	O	0	0
			14	14		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Programmed Cell Death 4, Pdcd4



#### • Molecule 1: Programmed Cell Death 4, Pdcd4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.65Å 64.65Å 164.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 28.39 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.00-2.00) 98.0 (28.39-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.259 , 0.297 0.257 , 0.295	Depositor DCC
$R_{free}$ test set	1381 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	44.2	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 66.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2110	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/1047	0.72	1/1409 (0.1%)
1	B	0.81	5/1047 (0.5%)	0.76	2/1409 (0.1%)
All	All	0.74	5/2094 (0.2%)	0.74	3/2818 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	342	GLY	C-O	10.58	1.40	1.23
1	B	342	GLY	C-N	6.91	1.50	1.34
1	B	333	MSE	CG-SE	6.41	2.17	1.95
1	B	329	LYS	CD-CE	5.93	1.66	1.51
1	B	329	LYS	CE-NZ	5.34	1.62	1.49

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	444	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	B	444	ARG	NE-CZ-NH1	6.41	123.51	120.30
1	A	334	LEU	CA-CB-CG	5.16	127.17	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1030	0	1028	20	0
1	B	1030	0	1028	30	0
2	A	36	0	0	1	0
2	B	14	0	0	5	0
All	All	2110	0	2056	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:333:MSE:SE	1:B:333:MSE:CE	2.16	1.44
1:B:333:MSE:CG	1:B:333:MSE:SE	2.17	1.42
1:B:391:TRP:CD1	1:B:401:MSE:HE3	1.98	0.97
1:B:391:TRP:HD1	1:B:401:MSE:HE3	1.37	0.82
1:A:349:HIS:O	1:A:353:GLU:HG3	1.79	0.81
1:B:348:GLU:O	1:B:352:LYS:HG2	1.88	0.73
1:A:397:THR:HG23	2:A:39:HOH:O	1.90	0.72
1:B:359:PHE:CZ	1:B:362:GLU:HG3	2.25	0.71
1:A:403:ARG:CG	1:A:403:ARG:HH11	2.04	0.70
1:B:344:ILE:HG21	1:B:389:SER:OG	1.97	0.65
1:A:375:THR:N	1:B:410:ASN:HD21	1.95	0.64
1:A:375:THR:H	1:B:410:ASN:HD21	1.47	0.62
1:B:391:TRP:HD1	1:B:401:MSE:CE	2.11	0.62
1:A:396:ILE:CG2	1:A:401:MSE:HG2	2.30	0.62
1:B:326:HIS:O	1:B:330:GLU:HG2	1.99	0.62
1:B:388:LYS:CE	2:B:38:HOH:O	2.47	0.61
1:B:333:MSE:SE	1:B:333:MSE:CB	2.99	0.61
1:B:396:ILE:CG2	1:B:401:MSE:HE2	2.37	0.55
1:A:396:ILE:HG22	1:A:401:MSE:HG2	1.89	0.55
1:B:388:LYS:HE3	2:B:38:HOH:O	2.06	0.55
1:A:356:VAL:HG13	1:A:356:VAL:O	2.07	0.55
1:B:402:LYS:O	1:B:406:GLU:HG3	2.08	0.54
1:A:324:VAL:HG11	1:A:329:LYS:HE2	1.90	0.54
1:A:326:HIS:O	1:A:330:GLU:HG2	2.08	0.53
1:A:403:ARG:CG	1:A:403:ARG:NH1	2.73	0.50
1:A:403:ARG:HG2	1:A:403:ARG:HH11	1.77	0.50
1:A:356:VAL:HG13	1:A:359:PHE:HB3	1.93	0.49
1:A:403:ARG:HG3	1:A:403:ARG:HH11	1.76	0.49
1:B:397:THR:HB	2:B:24:HOH:O	2.13	0.49
1:A:391:TRP:HB2	1:A:401:MSE:HE3	1.95	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:356:VAL:HG13	1:B:359:PHE:HB3	1.94	0.48
1:A:396:ILE:HG21	1:A:401:MSE:HG2	1.95	0.48
1:B:388:LYS:HE2	2:B:38:HOH:O	2.13	0.48
1:B:356:VAL:O	1:B:356:VAL:CG1	2.62	0.47
1:B:384:LEU:HG	2:B:34:HOH:O	2.12	0.47
1:A:398:ILE:O	1:A:398:ILE:HD13	2.16	0.46
1:B:351:LEU:HD22	1:B:363:LEU:HD13	1.98	0.45
1:B:396:ILE:HG22	1:B:401:MSE:HE2	1.99	0.45
1:B:362:GLU:HG2	1:B:407:ARG:HH12	1.84	0.43
1:B:361:HIS:CG	1:B:403:ARG:HG2	2.53	0.43
1:B:388:LYS:HA	1:B:438:ILE:HG21	2.01	0.43
1:B:365:TYR:O	1:B:369:VAL:HG23	2.18	0.43
1:B:388:LYS:HE2	1:B:388:LYS:HB3	1.87	0.42
1:A:388:LYS:HA	1:A:438:ILE:HG21	2.02	0.42
1:A:412:ILE:HG21	1:A:448:PRO:HG2	2.03	0.41
1:B:356:VAL:HG13	1:B:359:PHE:CB	2.51	0.40
1:A:334:LEU:C	1:A:334:LEU:HD23	2.42	0.40
1:B:367:ALA:HB1	1:B:383:ILE:HG23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/150 (83%)	125 (100%)	0	0	100	100
1	B	125/150 (83%)	123 (98%)	2 (2%)	0	100	100
All	All	250/300 (83%)	248 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/133 (89%)	110 (93%)	8 (7%)	18	13
1	B	118/133 (89%)	111 (94%)	7 (6%)	23	17
All	All	236/266 (89%)	221 (94%)	15 (6%)	20	15

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	324	VAL
1	A	356	VAL
1	A	363	LEU
1	A	398	ILE
1	A	403	ARG
1	A	417	LEU
1	A	443	LEU
1	A	444	ARG
1	B	324	VAL
1	B	356	VAL
1	B	397	THR
1	B	399	ASP
1	B	403	ARG
1	B	443	LEU
1	B	444	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	410	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/150 (82%)	0.67	13 (10%) <b>7</b> <b>7</b>	43, 50, 57, 59	0
1	B	123/150 (82%)	1.58	35 (28%) <b>1</b> <b>1</b>	41, 52, 59, 61	0
All	All	246/300 (82%)	1.12	48 (19%) <b>1</b> <b>1</b>	41, 51, 59, 61	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	417	LEU	10.9
1	B	421	HIS	7.8
1	A	324	VAL	6.9
1	B	418	ASP	6.5
1	B	419	VAL	6.3
1	B	340	LEU	5.9
1	B	341	SER	5.2
1	B	377	GLU	5.2
1	B	415	ILE	5.1
1	B	420	PRO	5.0
1	B	368	ILE	4.9
1	B	322	GLN	4.8
1	B	376	GLY	4.3
1	B	364	VAL	4.0
1	A	322	GLN	4.0
1	A	421	HIS	3.8
1	B	387	LEU	3.6
1	B	324	VAL	3.4
1	B	414	ASP	3.3
1	A	418	ASP	3.3
1	B	424	SER	3.1
1	B	391	TRP	2.8
1	B	378	SER	2.8
1	B	422	SER	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	363	LEU	2.7
1	B	339	LEU	2.7
1	B	416	ASN	2.6
1	A	387	LEU	2.6
1	B	386	LEU	2.6
1	B	349	HIS	2.5
1	B	405	TYR	2.5
1	A	420	PRO	2.4
1	B	428	ARG	2.4
1	B	399	ASP	2.3
1	B	375	THR	2.3
1	B	383	ILE	2.3
1	B	346	GLU	2.2
1	A	368	ILE	2.2
1	A	417	LEU	2.2
1	B	433	CYS	2.2
1	A	367	ALA	2.2
1	A	364	VAL	2.2
1	A	339	LEU	2.1
1	B	381	LYS	2.1
1	A	363	LEU	2.1
1	B	338	TYR	2.1
1	A	323	PRO	2.1
1	B	429	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.