



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 12:09 am GMT

PDB ID : 1ITQ  
Title : HUMAN RENAL DIPEPTIDASE  
Authors : Nitanai, Y.; Satow, Y.; Adachi, H.; Tsujimoto, M.  
Deposited on : 2002-02-02  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	<div> <div></div> <div> <div></div> <div>80%</div> <div>20%</div> <div></div> </div> </div>
1	B	369	<div> <div></div> <div> <div></div> <div>81%</div> <div>19%</div> <div></div> </div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RENAL DIPEPTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	0	0
			2888	1804	516	554	14			
1	B	369	Total	C	N	O	S	0	0	0
			2888	1804	516	554	14			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Zn 2	0	0
3	A	2	Total 2	Zn 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	113	Total 113	O 113	0	0
4	B	100	Total 100	O 100	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: RENAL DIPEPTIDASE



#### • Molecule 1: RENAL DIPEPTIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.95Å 81.21Å 57.16Å 90.00° 96.73° 90.00°	Depositor
Resolution (Å)	10.00 – 2.30 29.90 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.9 (10.00-2.30) 92.4 (29.90-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.93 (at 2.31Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, $R_{free}$	0.181 , 0.258 0.156 , 0.222	Depositor DCC
$R_{free}$ test set	3021 reflections (9.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.5	Xtriage
Anisotropy	0.866	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 55.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6049	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2947	0.64	1/4001 (0.0%)
1	B	0.37	0/2947	0.65	1/4001 (0.0%)
All	All	0.37	0/5894	0.65	2/8002 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	217	PHE	N-CA-C	-5.63	95.80	111.00
1	B	217	PHE	N-CA-C	-5.47	96.23	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2888	0	2815	47	0
1	B	2888	0	2815	40	0
2	A	28	0	26	0	0
2	B	28	0	26	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	113	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	100	0	0	1	0
All	All	6049	0	5682	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:324:GLY:HA2	1:B:328:ASP:HB2	1.60	0.81
1:A:193:LEU:HD23	1:A:337:VAL:HG21	1.68	0.74
1:A:222:ALA:H	1:A:270:HIS:HD2	1.36	0.73
1:B:193:LEU:HD23	1:B:337:VAL:HG21	1.71	0.71
1:A:324:GLY:HA2	1:A:328:ASP:HB2	1.71	0.71
1:A:262:ALA:HA	1:A:266:GLN:HE22	1.58	0.68
1:B:222:ALA:H	1:B:270:HIS:HD2	1.43	0.66
1:A:214:PRO:HB2	1:A:333:VAL:HG13	1.78	0.65
1:B:256:ILE:O	1:B:270:HIS:HE1	1.79	0.65
1:B:354:LEU:HD11	1:B:364:HIS:CD2	2.34	0.62
1:A:221:SER:HB3	1:A:270:HIS:CD2	2.35	0.62
1:B:263:ASN:H	1:B:266:GLN:NE2	1.97	0.61
1:A:263:ASN:H	1:A:266:GLN:HE21	1.46	0.61
1:A:221:SER:H	1:A:231:ASN:ND2	1.99	0.61
1:A:335:GLU:O	1:A:339:GLN:HG2	2.01	0.60
1:A:221:SER:H	1:A:231:ASN:HD22	1.51	0.59
1:A:205:LYS:O	1:A:209:GLN:HG2	2.05	0.56
1:A:256:ILE:O	1:A:270:HIS:HE1	1.88	0.56
1:B:125:GLU:HA	1:B:149:THR:HB	1.88	0.56
1:A:280:ALA:O	1:A:283:VAL:HG12	2.06	0.55
1:B:262:ALA:HA	1:B:266:GLN:HE22	1.70	0.55
1:A:8:GLU:O	1:A:12:ARG:HG3	2.07	0.55
1:A:197:ALA:O	1:A:198:HIS:HB2	2.05	0.55
1:A:354:LEU:HD13	1:A:367:TYR:CD2	2.43	0.54
1:B:8:GLU:O	1:B:12:ARG:HG3	2.08	0.54
1:A:262:ALA:HA	1:A:266:GLN:NE2	2.24	0.53
1:B:196:LEU:O	1:B:199:VAL:HG22	2.09	0.52
1:B:356:GLN:NE2	4:B:1019:HOH:O	2.43	0.52
1:A:103:VAL:HG23	1:A:121:LEU:HG	1.92	0.51
1:B:85:GLN:O	1:B:88:VAL:HG12	2.12	0.50
1:B:222:ALA:N	1:B:270:HIS:HD2	2.08	0.50
1:B:240:VAL:HG13	1:B:245:SER:HB2	1.94	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:THR:O	1:A:86:MET:HG3	2.12	0.49
1:B:263:ASN:H	1:B:266:GLN:HE21	1.59	0.49
1:B:218:SER:O	1:B:286:GLY:HA3	2.12	0.49
1:A:281:ARG:HB2	1:A:332:ARG:NH2	2.28	0.48
1:B:221:SER:HB3	1:B:270:HIS:CD2	2.48	0.48
1:B:82:THR:O	1:B:86:MET:HG3	2.14	0.48
1:A:331:LEU:O	1:A:335:GLU:HG3	2.15	0.47
1:A:67:VAL:HG13	1:A:85:GLN:HB2	1.97	0.47
1:B:214:PRO:HB2	1:B:333:VAL:HG13	1.96	0.47
1:B:126:GLY:H	1:B:149:THR:HB	1.79	0.46
1:A:221:SER:HB2	1:A:231:ASN:ND2	2.31	0.46
1:A:218:SER:O	1:A:286:GLY:HA3	2.14	0.46
1:A:172:GLN:HB3	4:A:1058:HOH:O	2.16	0.46
1:A:73:THR:O	1:A:78:ALA:HB2	2.14	0.46
1:A:319:GLU:HG2	1:A:323:LYS:HD2	1.96	0.46
1:B:10:ILE:HD13	1:B:319:GLU:HG3	1.98	0.46
1:A:229:ARG:HD3	1:A:229:ARG:HA	1.64	0.46
1:B:149:THR:HA	1:B:195:ASP:HB3	1.98	0.45
1:A:354:LEU:HD11	1:A:364:HIS:CD2	2.51	0.45
1:A:281:ARG:NH1	1:A:332:ARG:HD2	2.30	0.45
1:B:251:PHE:O	1:B:296:PRO:HD3	2.16	0.45
1:A:152:HIS:CG	1:A:153:SER:H	2.36	0.44
1:B:70:PRO:HB2	1:B:72:ASP:OD2	2.17	0.44
1:A:296:PRO:HB2	1:A:299:LEU:HB2	2.00	0.44
1:B:354:LEU:HD13	1:B:367:TYR:CD2	2.53	0.43
1:B:346:ALA:HA	1:B:347:PRO:HD3	1.91	0.43
1:B:146:ARG:HD3	1:B:146:ARG:HA	1.71	0.43
1:B:229:ARG:HD3	1:B:229:ARG:HA	1.78	0.43
1:A:221:SER:HB2	1:A:231:ASN:HD21	1.83	0.43
1:A:160:ASP:HB3	1:A:164:VAL:HG21	2.01	0.43
1:B:152:HIS:CG	1:B:153:SER:H	2.37	0.43
1:A:146:ARG:HA	1:A:146:ARG:HD3	1.83	0.42
1:A:256:ILE:HA	1:A:270:HIS:CE1	2.55	0.42
1:A:264:LEU:HD21	1:A:308:LEU:HA	2.01	0.42
1:A:250:ASN:O	1:A:256:ILE:HD11	2.20	0.42
1:A:319:GLU:O	1:A:323:LYS:HG3	2.20	0.42
1:B:189:ARG:O	1:B:341:SER:HB2	2.19	0.42
1:B:367:TYR:O	1:B:368:SER:HB2	2.20	0.42
1:B:249:VAL:HG21	1:B:271:LEU:HD23	2.02	0.41
1:B:263:ASN:N	1:B:266:GLN:NE2	2.66	0.41
1:A:74:GLN:O	1:A:75:ASN:HB2	2.21	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:SER:HB3	1:B:266:GLN:OE1	2.20	0.41
1:B:354:LEU:HD23	1:B:369:SER:HB2	2.01	0.41
1:B:79:VAL:HG22	1:B:129:SER:O	2.20	0.41
1:A:253:ASN:ND2	1:A:294:ARG:HB3	2.35	0.41
1:A:38:GLU:H	1:A:38:GLU:HG2	1.68	0.41
1:B:112:ALA:HB3	1:B:119:ALA:HB2	2.03	0.41
1:B:149:THR:HG22	1:B:151:THR:O	2.19	0.41
1:A:147:TYR:HA	1:A:193:LEU:O	2.20	0.41
1:B:197:ALA:O	1:B:198:HIS:HB2	2.20	0.41
1:B:311:GLU:O	1:B:315:ARG:HG3	2.20	0.41
1:A:346:ALA:HA	1:A:347:PRO:HD3	1.96	0.40
1:A:217:PHE:HB3	1:A:220:SER:HB2	2.03	0.40
1:A:288:ASP:O	1:A:292:VAL:HG22	2.21	0.40
1:A:124:VAL:HG23	1:A:145:MET:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/369 (100%)	345 (94%)	20 (5%)	2 (0%)	32	39
1	B	367/369 (100%)	341 (93%)	26 (7%)	0	100	100
All	All	734/738 (100%)	686 (94%)	46 (6%)	2 (0%)	44	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	368	SER
1	A	70	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/315 (100%)	310 (98%)	5 (2%)	68	82
1	B	315/315 (100%)	306 (97%)	9 (3%)	48	64
All	All	630/630 (100%)	616 (98%)	14 (2%)	57	74

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	147	TYR
1	A	212	ARG
1	A	295	VAL
1	A	334	PHE
1	A	342	ASN
1	B	50	THR
1	B	147	TYR
1	B	161	ASN
1	B	209	GLN
1	B	258	CYS
1	B	261	LYS
1	B	295	VAL
1	B	334	PHE
1	B	342	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	ASN
1	A	242	GLN
1	A	266	GLN
1	A	270	HIS
1	A	342	ASN
1	A	356	GLN
1	A	364	HIS
1	B	36	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	242	GLN
1	B	266	GLN
1	B	270	HIS
1	B	342	ASN
1	B	356	GLN
1	B	364	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	461	1	14,14,15	0.46	0	15,19,21	0.61	0
2	NAG	A	463	1	14,14,15	0.42	0	15,19,21	0.68	0
2	NAG	B	462	1	14,14,15	0.37	0	15,19,21	0.64	0
2	NAG	B	464	1	14,14,15	0.56	0	15,19,21	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	461	1	-	0/6/23/26	0/1/1/1
2	NAG	A	463	1	-	0/6/23/26	0/1/1/1
2	NAG	B	462	1	-	0/6/23/26	0/1/1/1
2	NAG	B	464	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/369 (100%)	-0.52	3 (0%) 86 89	4, 18, 39, 97	0
1	B	369/369 (100%)	-0.42	2 (0%) 90 93	6, 19, 43, 86	0
All	All	738/738 (100%)	-0.47	5 (0%) 87 90	4, 19, 42, 97	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	369	SER	11.2
1	A	369	SER	9.9
1	B	368	SER	6.4
1	A	368	SER	2.4
1	A	343	LEU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	462	14/15	0.83	0.17	0.39	45,52,56,57	0
3	ZN	B	411	1/1	0.99	0.09	-1.22	23,23,23,23	0
3	ZN	A	402	1/1	0.99	0.08	-1.28	19,19,19,19	0
3	ZN	B	412	1/1	1.00	0.08	-1.58	20,20,20,20	0
3	ZN	A	401	1/1	0.99	0.07	-2.10	22,22,22,22	0
2	NAG	A	461	14/15	0.92	0.16	-	38,43,50,55	0
2	NAG	B	464	14/15	0.48	0.52	-	65,73,80,81	0
2	NAG	A	463	14/15	0.91	0.16	-	50,62,69,69	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.