



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:09 am GMT

PDB ID : 2ITP
Title : CRYSTAL STRUCTURE OF EGFR KINASE DOMAIN G719S MUTATION
IN COMPLEX WITH AEE788
Authors : Yun, C.-H.; Boggon, T.J.; Li, Y.; Woo, S.; Greulich, H.; Meyerson, M.; Eck,
M.J.
Deposited on : 2006-05-25
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3342 (2.78-2.70)
Clashscore	112137	3731 (2.78-2.70)
Ramachandran outliers	110173	3670 (2.78-2.70)
Sidechain outliers	110143	3671 (2.78-2.70)
RSRZ outliers	101464	3362 (2.78-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	 9% 57% 31% 5% • 7%


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	65	Total O 65 65	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: EPIDERMAL GROWTH FACTOR RECEPTOR PRECURSOR



4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	144.62Å 144.62Å 144.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.80 – 2.74 24.10 – 2.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (24.80-2.74) 99.9 (24.10-2.74)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.76Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.255 0.193 , 0.252	Depositor DCC
R_{free} test set	940 reflections (7.59%)	DCC
Wilson B-factor (Å ²)	58.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.037 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2546	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AEE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.82	1/2499 (0.0%)	0.85	2/3381 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	818	CYS	CB-SG	-5.36	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	932	ARG	NE-CZ-NH1	-7.99	116.31	120.30
1	A	971	MET	CG-SD-CE	-5.68	91.11	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2448	0	2477	79	0
2	A	33	0	32	1	0
3	A	65	0	0	7	0
All	All	2546	0	2509	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:736:GLU:OE2	1:A:1016:TYR:OH	1.82	0.97
1:A:724:GLY:HA3	1:A:748:ARG:H	1.32	0.92
1:A:759:ILE:CG1	1:A:788:LEU:HD13	2.07	0.84
1:A:766:MET:HE2	3:A:3016:HOH:O	1.76	0.83
1:A:804:GLU:HG3	1:A:804:GLU:O	1.88	0.73
1:A:768:SER:O	1:A:831:ARG:NH1	2.23	0.72
1:A:759:ILE:HG12	1:A:788:LEU:CD1	2.23	0.68
1:A:917:GLY:O	1:A:918:ILE:C	2.33	0.67
1:A:715:ILE:O	1:A:716:LYS:HB3	1.95	0.66
1:A:918:ILE:HG21	1:A:926:ILE:HD12	1.77	0.66
1:A:715:ILE:O	1:A:715:ILE:HG22	1.95	0.66
1:A:766:MET:HB3	1:A:777:LEU:HB2	1.80	0.64
1:A:735:GLY:O	1:A:737:LYS:N	2.30	0.64
1:A:932:ARG:NH1	1:A:951:TRP:O	2.31	0.64
1:A:960:LYS:HD2	3:A:3036:HOH:O	1.97	0.64
1:A:759:ILE:HG12	1:A:788:LEU:HD13	1.80	0.64
1:A:716:LYS:HE3	3:A:3010:HOH:O	1.99	0.63
1:A:724:GLY:CA	1:A:748:ARG:H	2.10	0.63
1:A:945:MET:O	1:A:949:LYS:HG3	2.01	0.61
1:A:773:HIS:CE1	1:A:820:GLN:HG2	2.38	0.59
1:A:723:PHE:O	1:A:748:ARG:HG2	2.02	0.59
1:A:766:MET:HE1	2:A:2019:AEE:HAE	1.84	0.59
1:A:876:VAL:HG12	1:A:877:PRO:HD2	1.85	0.58
1:A:781:CYS:HB3	1:A:787:GLN:HE21	1.69	0.57
1:A:931:GLU:O	1:A:932:ARG:HD3	2.05	0.57
1:A:700:ASN:C	1:A:700:ASN:HD22	2.08	0.56
1:A:759:ILE:CG1	1:A:788:LEU:CD1	2.78	0.56
1:A:733:PRO:O	1:A:734:GLU:C	2.42	0.56
1:A:931:GLU:O	1:A:932:ARG:NE	2.40	0.55
1:A:765:VAL:HG13	1:A:833:LEU:HD11	1.89	0.55
1:A:759:ILE:CD1	1:A:788:LEU:HD13	2.39	0.53
1:A:715:ILE:O	1:A:715:ILE:CG2	2.56	0.52
1:A:931:GLU:O	1:A:932:ARG:CD	2.58	0.52
1:A:813:TYR:HD2	3:A:3030:HOH:O	1.93	0.52
1:A:802:VAL:HA	1:A:809:ILE:CD1	2.39	0.52
1:A:715:ILE:O	1:A:716:LYS:CB	2.58	0.51
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:830:ASP:C	1:A:832:ARG:H	2.12	0.50
1:A:759:ILE:HD11	1:A:788:LEU:HD13	1.93	0.50
1:A:829:GLU:HG3	1:A:893:HIS:CD2	2.46	0.50
1:A:733:PRO:O	1:A:734:GLU:O	2.30	0.50
1:A:766:MET:CE	3:A:3016:HOH:O	2.48	0.48
1:A:879:LYS:HD3	1:A:914:PRO:O	2.14	0.48
1:A:932:ARG:HD3	1:A:932:ARG:HA	1.51	0.48
1:A:805:HIS:O	1:A:806:LYS:C	2.51	0.47
1:A:705:ARG:HG2	1:A:707:LEU:CD2	2.45	0.47
1:A:705:ARG:NH1	1:A:1018:ILE:HG21	2.30	0.47
1:A:759:ILE:HG13	1:A:788:LEU:HD13	1.90	0.47
1:A:829:GLU:HG3	1:A:893:HIS:CG	2.50	0.47
1:A:961:PHE:O	1:A:965:ILE:HG13	2.16	0.46
1:A:824:GLY:HA3	1:A:853:ILE:HD12	1.97	0.46
1:A:983:GLY:O	1:A:984:ASP:C	2.54	0.46
1:A:728:LYS:HZ2	1:A:1002:MET:N	2.13	0.46
1:A:791:GLN:NE2	1:A:1012:ASP:OD2	2.48	0.46
1:A:725:THR:HB	1:A:727:TYR:CE1	2.52	0.45
1:A:886:ILE:HG21	1:A:924:SER:HB3	1.99	0.44
1:A:917:GLY:O	1:A:918:ILE:O	2.35	0.44
1:A:935:GLN:HA	1:A:944:TYR:CE2	2.52	0.44
1:A:714:LYS:C	1:A:715:ILE:HG13	2.37	0.44
1:A:723:PHE:O	1:A:748:ARG:CG	2.67	0.43
1:A:747:LEU:HG	1:A:759:ILE:HD12	2.01	0.43
1:A:815:LEU:HD12	1:A:975:PRO:HB3	2.00	0.43
1:A:960:LYS:HA	3:A:3035:HOH:O	2.19	0.43
1:A:783:THR:OG1	1:A:785:THR:O	2.27	0.42
1:A:815:LEU:O	1:A:819:VAL:HG23	2.20	0.42
1:A:705:ARG:HG2	1:A:707:LEU:HD21	2.01	0.42
1:A:736:GLU:OE2	1:A:1016:TYR:CZ	2.71	0.42
1:A:781:CYS:HB3	1:A:787:GLN:HB2	2.02	0.42
1:A:737:LYS:HA	3:A:3012:HOH:O	2.20	0.42
1:A:802:VAL:HA	1:A:809:ILE:HD11	2.02	0.41
1:A:700:ASN:C	1:A:700:ASN:ND2	2.73	0.41
1:A:876:VAL:HG12	1:A:881:MET:SD	2.61	0.41
1:A:935:GLN:HA	1:A:944:TYR:CD2	2.55	0.41
1:A:793:MET:HG3	1:A:844:LEU:HB2	2.03	0.41
1:A:916:ASP:OD2	1:A:917:GLY:N	2.54	0.41
1:A:714:LYS:HD3	1:A:727:TYR:CD2	2.55	0.41
1:A:738:VAL:O	1:A:738:VAL:HG23	2.21	0.40
1:A:936:PRO:HA	1:A:937:PRO:HD2	1.75	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:736:GLU:O	1:A:737:LYS:CB	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	302/327 (92%)	259 (86%)	24 (8%)	19 (6%)	1 2

All (19) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	722	ALA
1	A	723	PHE
1	A	734	GLU
1	A	736	GLU
1	A	864	ALA
1	A	716	LYS
1	A	865	GLU
1	A	1005	GLU
1	A	857	GLY
1	A	747	LEU
1	A	748	ARG
1	A	806	LYS
1	A	918	ILE
1	A	1006	ASP
1	A	753	PRO
1	A	754	LYS
1	A	872	GLU
1	A	889	ARG
1	A	715	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/288 (94%)	236 (87%)	34 (13%)	5 11

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	697	GLU
1	A	700	ASN
1	A	719	SER
1	A	725	THR
1	A	745	LYS
1	A	748	ARG
1	A	757	LYS
1	A	768	SER
1	A	776	ARG
1	A	784	SER
1	A	797	CYS
1	A	804	GLU
1	A	808	ASN
1	A	832	ARG
1	A	841	ARG
1	A	855	ASP
1	A	858	LEU
1	A	861	LEU
1	A	867	LYS
1	A	888	HIS
1	A	889	ARG
1	A	892	THR
1	A	921	SER
1	A	925	SER
1	A	928	GLU
1	A	932	ARG
1	A	941	ILE
1	A	969	SER
1	A	976	GLN
1	A	982	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	984	ASP
1	A	1004	GLU
1	A	1008	ASP
1	A	1018	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	700	ASN
1	A	756	ASN
1	A	773	HIS
1	A	787	GLN
1	A	893	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AEE	A	2019	-	36,37,37	1.66	3 (8%)	43,51,51	2.51	14 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AEE	A	2019	-	-	0/18/28/28	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2019	AEE	CAY-CBA	-5.85	1.40	1.48
2	A	2019	AEE	C6-C5	-4.20	1.40	1.44
2	A	2019	AEE	CAM-CBA	-3.37	1.33	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2019	AEE	N3-C2-N1	-9.38	120.69	128.86
2	A	2019	AEE	CAX-CAS-NBG	-4.57	104.38	113.17
2	A	2019	AEE	C5-C6-N1	-3.44	119.02	121.37
2	A	2019	AEE	CAM-CBA-CAY	-2.71	125.68	129.44
2	A	2019	AEE	CAZ-CBE-NAV	-2.43	107.79	112.68
2	A	2019	AEE	CAE-CAH-CAZ	-2.20	117.94	120.64
2	A	2019	AEE	CAH-CAZ-CAG	2.00	120.81	118.30
2	A	2019	AEE	CAR-CAP-NBF	2.34	115.37	110.63
2	A	2019	AEE	CAY-CBA-NAW	2.64	125.17	120.81
2	A	2019	AEE	CAP-NBF-CAO	2.69	114.97	108.87
2	A	2019	AEE	CAR-NBG-CAQ	2.72	115.03	108.87
2	A	2019	AEE	CAP-CAR-NBG	3.21	117.14	110.63
2	A	2019	AEE	C2-N3-C4	3.37	121.26	113.33
2	A	2019	AEE	C2-N1-C6	7.14	121.18	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2019	AEE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/327 (93%)	0.15	29 (9%) 9 7	28, 50, 102, 112	8 (2%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	875	LYS	5.1
1	A	863	GLY	4.8
1	A	722	ALA	4.6
1	A	723	PHE	4.5
1	A	864	ALA	4.1
1	A	721	GLY	3.8
1	A	1005	GLU	3.5
1	A	874	GLY	3.4
1	A	865	GLU	3.3
1	A	866	GLU	3.3
1	A	749	GLU	3.2
1	A	922	GLU	3.2
1	A	872	GLU	3.1
1	A	1003	ASP	3.1
1	A	867	LYS	3.0
1	A	747	LEU	3.0
1	A	748	ARG	2.9
1	A	750	ALA	2.8
1	A	1002	MET	2.7
1	A	753	PRO	2.6
1	A	862	LEU	2.6
1	A	868	GLU	2.5
1	A	751	THR	2.5
1	A	1004	GLU	2.4
1	A	754	LYS	2.3
1	A	720	SER	2.3
1	A	870	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	735	GLY	2.1
1	A	871	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	AEE	A	2019	33/33	0.88	0.23	1.28	64,69,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.