



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:24 am GMT

PDB ID : 4ITG  
Title : P113S mutant of E. coli Cystathionine beta-lyase MetC  
Authors : Squire, C.J.; Yosaatmadja, Y.; Soo, V.W.C.; Patrick, W.M.  
Deposited on : 2013-01-18  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2694 (1.76-1.72)
Clashscore	112137	2854 (1.76-1.72)
Ramachandran outliers	110173	2824 (1.76-1.72)
Sidechain outliers	110143	2824 (1.76-1.72)
RSRZ outliers	101464	2705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;">93%</div> <div style="position: absolute; top: 10px; right: 0; width: 10%; text-align: right;">6% ..</div> </div> </div>
1	B	395	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;">92%</div> <div style="position: absolute; top: 10px; right: 0; width: 10%; text-align: right;">7% ..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EPE	A	401	-	-	-	X
2	EPE	B	401	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cystathionine beta-lyase MetC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	P	S	0	5	0
			3065	1943	533	574	1	14			
1	B	392	Total	C	N	O	P	S	0	1	0
			3020	1914	526	565	1	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	SER	PRO	ENGINEERED MUTATION	UNP P06721
B	113	SER	PRO	ENGINEERED MUTATION	UNP P06721

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	193	Total	O	0	0
			193	193		
3	B	184	Total	O	0	0
			184	184		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Cystathionine beta-lyase MetC

Chain A: 



#### • Molecule 1: Cystathionine beta-lyase MetC

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.93Å 152.77Å 150.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.40 – 1.74 19.40 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.40-1.74) 99.5 (19.40-1.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.80 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.160 , 0.198 0.165 , 0.201	Depositor DCC
$R_{free}$ test set	3587 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.1	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6492	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.34 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.8812e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.04	4/3113 (0.1%)	0.96	6/4220 (0.1%)
1	B	1.00	3/3060 (0.1%)	0.95	9/4151 (0.2%)
All	All	1.02	7/6173 (0.1%)	0.95	15/8371 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	GLU	CG-CD	6.74	1.62	1.51
1	A	73	GLU	CD-OE2	6.00	1.32	1.25
1	B	35	VAL	CB-CG1	5.74	1.64	1.52
1	B	211	TYR	CE2-CZ	-5.54	1.31	1.38
1	B	131	TRP	CB-CG	-5.44	1.40	1.50
1	A	81	VAL	CB-CG2	5.36	1.64	1.52
1	A	118	CYS	CB-SG	-5.13	1.73	1.81

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	ARG	NE-CZ-NH1	10.22	125.41	120.30
1	B	285	ARG	NE-CZ-NH2	-9.62	115.49	120.30
1	B	285	ARG	NE-CZ-NH1	9.20	124.90	120.30
1	B	15	ARG	NE-CZ-NH1	8.72	124.66	120.30
1	B	15	ARG	NE-CZ-NH2	-8.69	115.96	120.30
1	A	15	ARG	NE-CZ-NH2	-8.23	116.19	120.30
1	B	254	ARG	NE-CZ-NH2	-6.44	117.08	120.30
1	B	49	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	393	ARG	NE-CZ-NH2	-6.11	117.25	120.30
1	A	393	ARG	NE-CZ-NH1	6.05	123.33	120.30
1	B	254	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	262	ARG	NE-CZ-NH1	5.64	123.12	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	149	LYS	CD-CE-NZ	5.38	124.06	111.70
1	B	217	ASP	CB-CG-OD1	5.24	123.01	118.30
1	A	263	LEU	CB-CG-CD1	-5.01	102.47	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3065	0	3020	11	0
1	B	3020	0	2965	13	0
2	A	15	0	18	0	0
2	B	15	0	18	0	0
3	A	193	0	0	0	0
3	B	184	0	0	1	0
All	All	6492	0	6021	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:355:ILE:HG22	1:A:364:ILE:HD11	1.75	0.68
1:B:9:GLN:NE2	1:B:73:GLU:OE2	2.29	0.61
1:B:211:TYR:HB2	3:B:589:HOH:O	2.01	0.60
1:A:355:ILE:CG2	1:A:364:ILE:HD11	2.33	0.58
1:B:269:SER:O	1:B:273:VAL:HG23	2.12	0.49
1:A:208:ALA:HB3	1:A:220:ILE:HG23	1.94	0.48
1:A:185:ASP:OD2	1:A:210:LLP:N1	2.47	0.47
1:B:208:ALA:HB3	1:B:220:ILE:HG23	1.97	0.46
1:A:28:ILE:CD1	1:B:30:ARG:CZ	2.94	0.46
1:A:15:ARG:HD2	1:A:24:VAL:O	2.16	0.45
1:B:72:CYS:SG	1:B:79:GLY:HA2	2.56	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:ILE:HD13	1:B:169:ILE:HG23	1.97	0.45
1:A:79:GLY:HA3	1:A:230:TRP:CE2	2.52	0.43
1:A:292:PRO:HA	1:A:297:HIS:CD2	2.54	0.42
1:B:370:LEU:C	1:B:370:LEU:HD23	2.39	0.42
1:B:141:VAL:HG13	1:B:172:ALA:HB1	2.01	0.41
1:B:106:MET:O	1:B:131:TRP:HA	2.20	0.41
1:A:98:ILE:HD12	1:A:98:ILE:C	2.41	0.41
1:B:98:ILE:HG23	1:B:150:ILE:HD12	2.02	0.41
1:A:28:ILE:HD11	1:B:30:ARG:NH1	2.35	0.40
1:B:392:ALA:HA	1:B:395:VAL:HG22	2.03	0.40
1:A:392:ALA:HA	1:A:395:VAL:HG23	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/395 (100%)	381 (97%)	11 (3%)	1 (0%)	44	24
1	B	390/395 (99%)	379 (97%)	11 (3%)	0	100	100
All	All	783/790 (99%)	760 (97%)	22 (3%)	1 (0%)	55	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	191	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/321 (100%)	320 (99%)	2 (1%)	89	82
1	B	314/321 (98%)	312 (99%)	2 (1%)	89	82
All	All	636/642 (99%)	632 (99%)	4 (1%)	89	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	364	ILE
1	B	251	ILE
1	B	395	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	LLP	A	210	1	24,24,25	2.71	7 (29%)	28,32,34	2.62	14 (50%)
1	LLP	B	210	1	24,24,25	2.21	6 (25%)	28,32,34	2.86	14 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	210	1	-	0/15/17/19	0/1/1/1
1	LLP	B	210	1	-	0/15/17/19	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	LLP	C2'-C2	-3.28	1.44	1.50
1	A	210	LLP	C3-C2	-3.11	1.38	1.40
1	B	210	LLP	C2'-C2	-2.57	1.45	1.50
1	B	210	LLP	C4-C5	-2.48	1.38	1.42
1	B	210	LLP	O-C	2.03	1.28	1.19
1	B	210	LLP	CB-CA	2.15	1.56	1.53
1	A	210	LLP	CA-C	2.16	1.53	1.50
1	A	210	LLP	CD-CE	3.09	1.62	1.51
1	B	210	LLP	C6-N1	3.38	1.41	1.34
1	A	210	LLP	CB-CA	3.64	1.58	1.53
1	A	210	LLP	C6-N1	4.31	1.43	1.34
1	B	210	LLP	C4'-NZ	7.90	1.50	1.27
1	A	210	LLP	C4'-NZ	9.73	1.55	1.27

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	210	LLP	C3-C2-N1	-4.21	115.22	120.75
1	A	210	LLP	C4-C4'-NZ	-4.16	104.44	124.66
1	B	210	LLP	C4-C4'-NZ	-4.04	105.03	124.66
1	A	210	LLP	C3-C2-N1	-3.71	115.87	120.75
1	A	210	LLP	C3-C4-C4'	-3.53	113.76	120.52
1	A	210	LLP	OP4-P-OP1	-3.52	96.61	106.47
1	B	210	LLP	OP4-P-OP1	-3.45	96.80	106.47
1	B	210	LLP	CD-CE-NZ	-3.42	103.31	110.88
1	B	210	LLP	OP2-P-OP4	-3.38	97.74	106.73
1	B	210	LLP	CG-CD-CE	-2.59	103.65	113.59
1	A	210	LLP	O-C-CA	-2.52	118.07	125.02
1	B	210	LLP	OP3-P-OP4	-2.35	100.49	106.73
1	A	210	LLP	C6-N1-C2	2.04	123.18	119.26
1	A	210	LLP	CB-CA-C	2.12	115.14	111.65
1	A	210	LLP	C2'-C2-C3	2.19	123.57	120.96
1	A	210	LLP	CD-CG-CB	2.28	121.72	113.63
1	B	210	LLP	C3-C4-C5	2.31	120.00	118.24

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	210	LLP	C2'-C2-C3	2.58	124.04	120.96
1	A	210	LLP	C5'-C5-C4	2.77	126.51	121.66
1	A	210	LLP	OP2-P-OP1	2.99	122.19	110.50
1	B	210	LLP	CE-NZ-C4'	3.29	128.58	119.03
1	B	210	LLP	C5'-C5-C4	3.43	127.66	121.66
1	A	210	LLP	CE-NZ-C4'	3.46	129.08	119.03
1	A	210	LLP	C5-C4-C4'	3.52	126.62	121.36
1	B	210	LLP	OP2-P-OP1	4.05	126.35	110.50
1	B	210	LLP	CB-CA-C	5.83	121.26	111.65
1	B	210	LLP	C4-C3-C2	6.43	124.11	120.15
1	A	210	LLP	C4-C3-C2	6.72	124.28	120.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	210	LLP	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EPE	A	401	-	15,15,15	1.11	1 (6%)	18,20,20	3.22	6 (33%)
2	EPE	B	401	-	15,15,15	1.10	1 (6%)	18,20,20	2.09	5 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	A	401	-	-	0/9/19/19	0/1/1/1
2	EPE	B	401	-	-	0/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	EPE	C10-S	2.99	1.81	1.77
2	A	401	EPE	C10-S	3.45	1.82	1.77

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	EPE	O1S-S-C10	-11.42	96.98	106.79
2	B	401	EPE	O1S-S-C10	-2.61	104.55	106.79
2	A	401	EPE	C6-N1-C2	2.18	113.81	108.87
2	A	401	EPE	C2-C3-N4	2.37	115.43	110.63
2	A	401	EPE	O3S-S-O1S	2.97	118.18	111.37
2	A	401	EPE	C5-N4-C3	3.02	115.71	108.87
2	B	401	EPE	C9-N1-C6	3.12	119.25	111.26
2	B	401	EPE	O3S-S-C10	3.19	109.98	106.06
2	A	401	EPE	C6-C5-N4	3.80	118.33	110.63
2	B	401	EPE	C9-N1-C2	4.12	121.82	111.26
2	B	401	EPE	C6-N1-C2	4.28	118.57	108.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	390/395 (98%)	-0.31	3 (0%) 86 90	8, 14, 24, 34	0
1	B	391/395 (98%)	-0.26	2 (0%) 90 93	9, 15, 25, 34	0
All	All	781/790 (98%)	-0.28	5 (0%) 89 92	8, 14, 25, 34	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	395	VAL	4.6
1	B	4	LYS	3.3
1	A	395	VAL	2.7
1	A	364	ILE	2.6
1	A	47[A]	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	B	210	24/25	0.98	0.07	-	9,12,23,24	0
1	LLP	A	210	24/25	0.98	0.07	-	7,11,21,22	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	EPE	B	401	15/15	0.95	0.12	3.57	18,33,45,46	0
2	EPE	A	401	15/15	0.96	0.11	2.96	19,32,41,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.