



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 18, 2017 – 11:39 PM EST

PDB ID : 5IUA
Title : Crystal structure of stabilized A2A adenosine receptor A2AR-StaR2-bRIL in complex with compound 12b at 2.2Å resolution
Authors : Segala, E.; Guo, D.; Cheng, R.K.Y.; Bortolato, A.; Defforian, F.; Dore, A.S.; Errey, J.C.; Heitman, L.H.; Ijzerman, A.P.; Marshall, F.H.; Cooke, R.M.
Deposited on : unknown
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div style="width: 4%; height: 10px; background-color: red;"></div> <div style="width: 85%; height: 10px; background-color: green;"></div> <div style="width: 5%; height: 10px; background-color: yellow;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CLR	A	2402	-	-	-	X
5	OLA	A	2406	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	OLA	A	2407	-	-	-	X
5	OLA	A	2408	-	-	-	X
5	OLA	A	2409	-	-	-	X
5	OLA	A	2410	-	-	-	X
5	OLA	A	2412	-	-	-	X
5	OLA	A	2414	-	-	-	X
5	OLA	A	2416	-	-	-	X
5	OLA	A	2417	-	-	-	X
5	OLA	A	2418	-	-	-	X
5	OLA	A	2420	-	-	-	X
5	OLA	A	2422	-	-	-	X
6	OLB	A	2423	-	-	-	X
6	OLB	A	2425	-	-	-	X
6	OLB	A	2427	-	-	-	X
6	OLB	A	2428	-	-	-	X
6	OLB	A	2429	-	-	-	X
6	OLB	A	2430	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosine receptor A2a,Soluble cytochrome b562,Adenosine receptor A2a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	21	0
			3144	2053	527	541	23			

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	ASP	-	expression tag	UNP P29274
A	-7	TYR	-	expression tag	UNP P29274
A	-6	LYS	-	expression tag	UNP P29274
A	-5	ASP	-	expression tag	UNP P29274
A	-4	ASP	-	expression tag	UNP P29274
A	-3	ASP	-	expression tag	UNP P29274
A	-2	ASP	-	expression tag	UNP P29274
A	-1	GLY	-	expression tag	UNP P29274
A	0	ALA	-	expression tag	UNP P29274
A	1	PRO	-	expression tag	UNP P29274
A	54	LEU	ALA	engineered mutation	UNP P29274
A	88	ALA	THR	engineered mutation	UNP P29274
A	107	ALA	ARG	engineered mutation	UNP P29274
A	122	ALA	LYS	engineered mutation	UNP P29274
A	154	ALA	ASN	engineered mutation	UNP P29274
A	202	ALA	LEU	engineered mutation	UNP P29274
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	LEU	ARG	engineered mutation	UNP P0ABE7
A	235	ALA	LEU	engineered mutation	UNP P29274
A	239	ALA	VAL	engineered mutation	UNP P29274
A	277	ALA	SER	engineered mutation	UNP P29274
A	316	ALA	-	expression tag	UNP P29274
A	317	ALA	-	expression tag	UNP P29274
A	318	ALA	-	expression tag	UNP P29274
A	319	HIS	-	expression tag	UNP P29274

Continued on next page...


Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	320	HIS	-	expression tag	UNP P29274
A	321	HIS	-	expression tag	UNP P29274
A	322	HIS	-	expression tag	UNP P29274
A	323	HIS	-	expression tag	UNP P29274
A	324	HIS	-	expression tag	UNP P29274
A	325	HIS	-	expression tag	UNP P29274
A	326	HIS	-	expression tag	UNP P29274
A	327	HIS	-	expression tag	UNP P29274
A	328	HIS	-	expression tag	UNP P29274

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0


- Molecule 3 is 2-(furan-2-yl)-N 5 -[3-(4-phenylpiperazin-1-yl)propyl][1,2,4]triazolo[1,5-a][1,3,5]triazine-5,7-diamine (three-letter code: 6DX) (formula: C₂₁H₂₅N₉O).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			28	27	1		
4	A	1	Total	C	O	0	0
			28	27	1		
4	A	1	Total	C	O	0	0
			28	27	1		
4	A	1	Total	C	O	0	0
			28	27	1		

- Molecule 5 is OLEIC ACID (three-letter code: OLA) (formula: C₁₈H₃₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			20	18	2		
5	A	1	Total	C	O	0	0
			15	13	2		
5	A	1	Total	C	O	0	0
			9	7	2		
5	A	1	Total	C	O	0	0
			18	16	2		
5	A	1	Total	C	O	0	0
			20	18	2		
5	A	1	Total	C	O	0	0
			15	13	2		
5	A	1	Total	C	O	0	0
			12	10	2		
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	0
			14	12	2		
5	A	1	Total	C	O	0	0
			19	17	2		
5	A	1	Total	C	O	0	0
			11	9	2		
5	A	1	Total	C		0	0
			15	15			
5	A	1	Total	C	O	0	0
			14	12	2		
5	A	1	Total	C	O	0	0
			15	13	2		
5	A	1	Total	C		0	0
			7	7			
5	A	1	Total	C		0	0
			8	8			
5	A	1	Total	C		0	0
			12	12			

- Molecule 6 is (2S)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLB) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			17	13	4		
6	A	1	Total	C	O	0	0
			19	15	4		
6	A	1	Total	C	O	0	0
			16	12	4		
6	A	1	Total	C	O	0	0
			22	18	4		
6	A	1	Total	C	O	0	0
			19	15	4		
6	A	1	Total	C	O	0	0
			25	21	4		
6	A	1	Total	C	O	0	0
			20	16	4		
6	A	1	Total	C	O	0	0
			21	17	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	131	Total	O	0	0
			131	131		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosine receptor A2a,Soluble cytochrome b562,Adenosine receptor A2a



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	39.56Å 179.98Å 139.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.82 – 2.20 33.82 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.7 (33.82-2.20) 94.4 (33.82-2.20)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.200 , 0.233 0.193 , 0.226	Depositor DCC
R_{free} test set	1205 reflections (4.92%)	DCC
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.145	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 61.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3810	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, OLB, NA, CLR, 6DX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3221	0.36	0/4379

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3144	0	3238	17	0
2	A	1	0	0	0	0
3	A	31	0	0	0	0
4	A	112	0	184	3	0
5	A	232	0	330	12	0
6	A	159	0	220	6	0
7	A	131	0	0	3	0
All	All	3810	0	3972	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2417:OLA:H72	6:A:2428:OLB:H19	1.80	0.62
1:A:25:VAL:HG22	5:A:2415:OLA:H132	1.85	0.57
1:A:32:TRP:CE3	5:A:2415:OLA:H71	2.40	0.56
1:A:162:GLY:O	7:A:2501:HOH:O	2.18	0.56
1:A:65:THR:HG21	6:A:2427:OLB:H4A	1.88	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/433 (94%)	403 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/353 (94%)	333 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 31 ligands modelled in this entry, 1 is monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	6DX	A	2401	-	28,35,35	2.85	10 (35%)	31,48,48	2.18	7 (22%)
4	CLR	A	2402	-	31,31,31	0.68	0	48,48,48	1.02	2 (4%)
4	CLR	A	2403	-	31,31,31	0.64	0	48,48,48	1.05	2 (4%)
4	CLR	A	2404	-	31,31,31	0.67	0	48,48,48	0.98	2 (4%)
4	CLR	A	2405	-	31,31,31	0.69	0	48,48,48	0.93	0
5	OLA	A	2406	-	16,19,19	0.41	0	15,19,19	0.28	0
5	OLA	A	2407	-	11,14,19	0.36	0	10,14,19	0.32	0
5	OLA	A	2408	-	5,8,19	0.24	0	4,8,19	0.18	0
5	OLA	A	2409	-	14,17,19	0.40	0	13,17,19	0.24	0
5	OLA	A	2410	-	16,19,19	0.44	0	15,19,19	0.25	0
5	OLA	A	2411	-	11,14,19	0.36	0	10,14,19	0.30	0
5	OLA	A	2412	-	8,11,19	0.45	0	7,11,19	0.45	0
5	OLA	A	2413	-	4,7,19	0.26	0	3,7,19	0.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	OLA	A	2414	-	10,13,19	0.43	0	9,13,19	0.41	0
5	OLA	A	2415	-	15,18,19	0.42	0	14,18,19	0.25	0
5	OLA	A	2416	-	7,10,19	0.24	0	6,10,19	0.33	0
5	OLA	A	2417	-	14,14,19	0.41	0	13,13,19	0.29	0
5	OLA	A	2418	-	10,13,19	0.43	0	9,13,19	0.44	0
5	OLA	A	2419	-	11,14,19	0.37	0	10,14,19	0.35	0
5	OLA	A	2420	-	6,6,19	0.25	0	5,5,19	0.15	0
5	OLA	A	2421	-	7,7,19	0.24	0	6,6,19	0.21	0
5	OLA	A	2422	-	11,11,19	0.41	0	10,10,19	0.31	0
6	OLB	A	2423	-	16,16,24	0.98	2 (12%)	17,17,25	1.05	1 (5%)
6	OLB	A	2424	-	18,18,24	0.91	2 (11%)	19,19,25	1.02	1 (5%)
6	OLB	A	2425	-	15,15,24	0.99	2 (13%)	16,16,25	0.96	1 (6%)
6	OLB	A	2426	-	21,21,24	0.84	2 (9%)	22,22,25	0.97	1 (4%)
6	OLB	A	2427	-	18,18,24	0.93	2 (11%)	19,19,25	1.01	1 (5%)
6	OLB	A	2428	-	24,24,24	0.80	2 (8%)	25,25,25	0.87	1 (4%)
6	OLB	A	2429	-	19,19,24	0.89	2 (10%)	20,20,25	0.92	1 (5%)
6	OLB	A	2430	-	20,20,24	0.88	2 (10%)	21,21,25	0.87	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6DX	A	2401	-	-	0/11/25/25	0/4/5/5
4	CLR	A	2402	-	-	0/10/68/68	0/4/4/4
4	CLR	A	2403	-	-	0/10/68/68	0/4/4/4
4	CLR	A	2404	-	-	0/10/68/68	0/4/4/4
4	CLR	A	2405	-	-	0/10/68/68	0/4/4/4
5	OLA	A	2406	-	-	0/15/17/17	0/0/0/0
5	OLA	A	2407	-	-	0/10/12/17	0/0/0/0
5	OLA	A	2408	-	-	0/4/6/17	0/0/0/0
5	OLA	A	2409	-	-	0/13/15/17	0/0/0/0
5	OLA	A	2410	-	-	0/15/17/17	0/0/0/0
5	OLA	A	2411	-	-	0/10/12/17	0/0/0/0
5	OLA	A	2412	-	-	0/7/9/17	0/0/0/0
5	OLA	A	2413	-	-	0/3/5/17	0/0/0/0
5	OLA	A	2414	-	-	0/9/11/17	0/0/0/0
5	OLA	A	2415	-	-	0/14/16/17	0/0/0/0
5	OLA	A	2416	-	-	0/6/8/17	0/0/0/0
5	OLA	A	2417	-	-	0/12/12/17	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OLA	A	2418	-	-	0/9/11/17	0/0/0/0
5	OLA	A	2419	-	-	0/10/12/17	0/0/0/0
5	OLA	A	2420	-	-	0/4/4/17	0/0/0/0
5	OLA	A	2421	-	-	0/5/5/17	0/0/0/0
5	OLA	A	2422	-	-	0/9/9/17	0/0/0/0
6	OLB	A	2423	-	-	0/16/16/24	0/0/0/0
6	OLB	A	2424	-	-	0/18/18/24	0/0/0/0
6	OLB	A	2425	-	-	0/15/15/24	0/0/0/0
6	OLB	A	2426	-	-	0/21/21/24	0/0/0/0
6	OLB	A	2427	-	-	0/18/18/24	0/0/0/0
6	OLB	A	2428	-	-	0/24/24/24	0/0/0/0
6	OLB	A	2429	-	-	0/19/19/24	0/0/0/0
6	OLB	A	2430	-	-	0/20/20/24	0/0/0/0

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2401	6DX	C08-N09	-8.51	1.27	1.47
3	A	2401	6DX	C11-N12	-5.83	1.37	1.46
3	A	2401	6DX	C13-N12	-5.82	1.37	1.46
3	A	2401	6DX	C14-N09	-2.78	1.39	1.47
3	A	2401	6DX	C10-N09	-2.68	1.39	1.47

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2401	6DX	N25-C24-N23	-9.24	108.46	114.58
3	A	2401	6DX	N21-C04-N03	-3.96	120.23	126.23
4	A	2403	CLR	C18-C13-C17	-2.62	106.78	111.73
3	A	2401	6DX	C06-N05-C04	-2.21	119.65	123.68
3	A	2401	6DX	N03-C02-N26	-2.00	119.65	124.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2402	CLR	2	0
4	A	2403	CLR	1	0
4	A	2405	CLR	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	2410	OLA	5	0
5	A	2414	OLA	2	0
5	A	2415	OLA	2	0
5	A	2417	OLA	3	0
5	A	2419	OLA	1	0
5	A	2421	OLA	1	0
6	A	2425	OLB	2	0
6	A	2427	OLB	1	0
6	A	2428	OLB	2	0
6	A	2430	OLB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/433 (89%)	-0.10	19 (4%) 30 29	11, 26, 66, 103	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1021	ASP	4.3
1	A	-1	GLY	4.1
1	A	1061	PHE	4.1
1	A	1062	ARG	3.8
1	A	306	HIS	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CLR	A	2402	28/28	0.57	0.35	18.49	92,96,100,101	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	OLB	A	2430	21/25	0.79	0.24	10.10	40,50,58,60	0
5	OLA	A	2408	9/20	0.86	0.23	8.65	29,38,53,60	0
5	OLA	A	2410	20/20	0.62	0.25	7.93	43,53,63,65	0
5	OLA	A	2416	11/20	0.79	0.20	7.65	44,56,81,81	0
5	OLA	A	2417	15/20	0.85	0.17	6.98	45,49,56,56	0
6	OLB	A	2425	16/25	0.76	0.20	6.94	44,70,82,82	0
5	OLA	A	2422	12/20	0.84	0.24	6.42	40,44,48,50	0
5	OLA	A	2418	14/20	0.78	0.19	6.34	41,52,65,67	0
6	OLB	A	2429	20/25	0.71	0.28	6.27	39,52,89,91	0
5	OLA	A	2420	7/20	0.54	0.20	5.28	52,54,55,55	0
5	OLA	A	2407	15/20	0.81	0.23	4.77	45,51,66,67	0
5	OLA	A	2406	20/20	0.90	0.15	4.77	26,44,60,61	0
6	OLB	A	2423	17/25	0.80	0.20	4.10	55,60,64,66	0
6	OLB	A	2428	25/25	0.69	0.25	4.07	38,49,72,74	0
5	OLA	A	2414	14/20	0.63	0.23	4.00	44,57,60,60	0
5	OLA	A	2412	12/20	0.78	0.28	3.18	35,48,61,62	0
6	OLB	A	2427	19/25	0.86	0.17	2.90	33,37,65,67	0
5	OLA	A	2409	18/20	0.82	0.17	2.86	40,51,62,66	0
5	OLA	A	2419	15/20	0.78	0.20	1.98	37,41,64,67	0
5	OLA	A	2415	19/20	0.73	0.22	1.93	46,52,68,69	0
6	OLB	A	2426	22/25	0.85	0.18	1.87	37,43,66,69	0
5	OLA	A	2421	8/20	0.76	0.14	1.58	45,46,48,48	0
3	6DX	A	2401	31/31	0.94	0.18	1.46	12,21,60,62	0
4	CLR	A	2405	28/28	0.92	0.11	0.66	19,26,43,45	0
4	CLR	A	2404	28/28	0.94	0.11	0.64	21,26,39,46	0
4	CLR	A	2403	28/28	0.94	0.10	0.08	19,25,41,42	0
2	NA	A	2400	1/1	0.96	0.13	-0.96	31,31,31,31	0
5	OLA	A	2413	8/20	0.61	0.18	-	52,56,61,63	0
5	OLA	A	2411	15/20	0.76	0.22	-	62,63,73,74	0
6	OLB	A	2424	19/25	0.87	0.17	-	40,49,63,64	0

6.5 Other polymers ⓘ

There are no such residues in this entry.