



Full wwPDB X-ray Structure Validation Report i

Feb 12, 2017 – 09:56 pm GMT

PDB ID : 1IW6
Title : Crystal Structure of the Ground State of Bacteriorhodopsin
Authors : Kouyama, T.; Okumura, H.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2002-04-22
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbitiy : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriaage (Phenix) : NOT EXECUTED
EDS : NOT EXECUTED
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	400	X	-	-	-

2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 2095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called bacteriorhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	3	Total	C	O	0	0


- Molecule 3 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0


- Molecule 4 is 2,3-DI-O-PHYTANLY-3-SN-GLYCERO-1-PHOSPHORYL-3'-SN-GLYCER

OL-1'-PHOSPHATE (three-letter code: L3P) (formula: C₄₆H₉₄O₁₁P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			59	46	11	2		
4	A	1	Total	C	O	P	0	0
			40	33	6	1		
4	A	1	Total	C	O	P	0	0
			50	43	6	1		
4	A	1	Total	C	O	P	0	0
			50	43	6	1		

- Molecule 5 is 2,3-DI-PHYTANYL-GLYCEROL (three-letter code: L2P) (formula: C₄₃H₈₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total 46 43 3	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	41	Total 41 41	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: bacteriorhodopsin



4 Data and refinement statistics [\(i\)](#)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value			Source
Space group	P 6 2 2			Depositor
Cell constants a, b, c, α , β , γ	102.40 Å 90.00°	102.40 Å 90.00°	112.10 Å 120.00°	Depositor
Resolution (Å)	28.40 – 2.30			Depositor
% Data completeness (in resolution range)	98.0 (28.40-2.30)			Depositor
R_{merge}	0.08			Depositor
R_{sym}	0.08			Depositor
Refinement program	CNS 1.0			Depositor
R , R_{free}	0.251 , 0.275			Depositor
Estimated twinning fraction	No twinning to report.			Xtriage
Total number of atoms	2095			wwPDB-VP
Average B, all atoms (Å ²)	56.0			wwPDB-VP

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, RET, L2P, GAL, L3P, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1804	0.60	0/2464

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	400	GLC	C1

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1813	49	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	33	0	28	2	0
3	A	20	0	27	3	0
4	A	199	0	331	6	0
5	A	46	0	86	2	0
6	A	41	0	0	4	0
All	All	2095	0	2285	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:THR:HG22	1:A:141:SER:HB2	1.52	0.90
1:A:15:LEU:HD23	1:A:209:MET:HE1	1.67	0.77
1:A:44:ALA:HA	4:A:260:L3P:H111	1.65	0.77
1:A:107:THR:HG21	6:A:629:HOH:O	1.88	0.72
1:A:117:ILE:O	1:A:121:THR:HB	1.90	0.72
1:A:174:LEU:O	1:A:178:THR:HG23	1.90	0.72
1:A:40:LYS:HZ1	4:A:260:L3P:H42	1.55	0.71
1:A:227:ARG:HB3	1:A:227:ARG:HH11	1.60	0.66
1:A:29:VAL:O	1:A:29:VAL:HG12	1.97	0.65
1:A:121:THR:CG2	1:A:141:SER:HB2	2.29	0.61
1:A:49:VAL:HG21	1:A:93:LEU:HD13	1.84	0.59
1:A:104:ASP:HB2	1:A:107:THR:HG22	1.85	0.59
1:A:48:LEU:HG	4:A:260:L3P:H143	1.86	0.57
1:A:101:VAL:HG22	1:A:159:LYS:HB3	1.86	0.57
1:A:6:GLY:HA2	1:A:9:GLU:OE1	2.03	0.57
1:A:107:THR:O	1:A:111:LEU:HD23	2.05	0.57
1:A:38:ASP:HB3	6:A:628:HOH:O	2.04	0.57
1:A:70:PRO:HA	1:A:74:GLU:O	2.06	0.56
1:A:85:ASP:O	1:A:89:THR:HB	2.06	0.55
1:A:44:ALA:HA	4:A:260:L3P:H142	1.91	0.53
1:A:141:SER:HB3	3:A:250:RET:H41	1.90	0.52
1:A:156:PHE:HB3	1:A:171:PHE:CZ	2.44	0.52
1:A:186:PRO:HB3	3:A:250:RET:H183	1.93	0.51
1:A:25:LEU:O	1:A:29:VAL:HG23	2.11	0.50
1:A:90:THR:OG1	1:A:91:PRO:HD3	2.13	0.49
1:A:53:ALA:HB2	1:A:216:LYS:HE2	1.94	0.48
1:A:166:GLU:HG3	6:A:630:HOH:O	2.12	0.48
1:A:129:LYS:HZ1	2:A:410:MAN:H61	1.79	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:TRP:O	1:A:209:MET:HE2	2.15	0.46
1:A:111:LEU:HD21	1:A:151:VAL:HG11	1.96	0.46
1:A:168:ALA:O	1:A:172:LYS:HG2	2.15	0.46
1:A:7:ARG:H	1:A:8:PRO:HD2	1.81	0.46
1:A:152:LEU:O	1:A:156:PHE:HB2	2.16	0.46
1:A:15:LEU:HB3	1:A:209:MET:HE2	1.98	0.46
1:A:90:THR:N	1:A:91:PRO:CD	2.79	0.45
1:A:198:ILE:HG12	4:A:290:L3P:H441	1.99	0.45
3:A:250:RET:H7	3:A:250:RET:H181	1.85	0.44
1:A:87:LEU:HD21	5:A:270:L2P:H262	2.00	0.44
1:A:101:VAL:CG2	1:A:159:LYS:HB3	2.47	0.43
1:A:5:THR:O	1:A:201:LEU:HD23	2.17	0.43
1:A:165:PRO:HG2	6:A:630:HOH:O	2.18	0.42
1:A:15:LEU:HB3	1:A:209:MET:CE	2.49	0.42
1:A:46:THR:O	1:A:50:PRO:HD2	2.19	0.42
1:A:196:ALA:O	1:A:198:ILE:N	2.46	0.42
1:A:12:TRP:CD2	1:A:206:LEU:HD23	2.55	0.42
1:A:7:ARG:N	1:A:8:PRO:HD2	2.35	0.42
1:A:104:ASP:O	1:A:107:THR:HG23	2.21	0.41
1:A:120:GLY:HA3	5:A:270:L2P:H491	2.02	0.41
1:A:129:LYS:NZ	2:A:410:MAN:H61	2.35	0.41
1:A:28:LEU:HD13	4:A:260:L3P:H522	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	225/248 (91%)	214 (95%)	7 (3%)	4 (2%)	10 9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	73	GLY
1	A	197	GLY
1	A	29	VAL

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	182/194 (94%)	169 (93%)	13 (7%)	17 22

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	38	ASP
1	A	66	LEU
1	A	89	THR
1	A	93	LEU
1	A	94	LEU
1	A	97	LEU
1	A	101	VAL
1	A	107	THR
1	A	121	THR
1	A	161	GLU
1	A	211	LEU
1	A	227	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	A	400	2,5	11,11,12	2.35	4 (36%)	13,15,17	1.14	0
2	MAN	A	410	2	11,11,12	2.18	6 (54%)	13,15,17	1.49	2 (15%)
2	GAL	A	420	2	11,11,12	2.62	6 (54%)	13,15,17	1.23	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	400	2,5	1/1/4/5	0/2/19/22	0/1/1/1
2	MAN	A	410	2	-	0/2/19/22	0/1/1/1
2	GAL	A	420	2	-	0/2/19/22	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	410	MAN	C4-C5	2.04	1.57	1.53
2	A	410	MAN	O5-C5	2.16	1.48	1.43
2	A	410	MAN	C1-C2	2.42	1.57	1.52
2	A	410	MAN	O5-C1	2.44	1.47	1.43
2	A	420	GAL	C1-C2	2.78	1.58	1.52
2	A	420	GAL	C2-C3	2.84	1.56	1.52
2	A	400	GLC	C2-C3	3.23	1.56	1.52
2	A	420	GAL	C4-C3	3.26	1.60	1.52
2	A	410	MAN	O2-C2	3.29	1.50	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	GLC	C1-C2	3.30	1.60	1.52
2	A	420	GAL	O5-C1	3.49	1.49	1.43
2	A	420	GAL	C4-C5	3.69	1.60	1.53
2	A	400	GLC	O5-C1	3.76	1.49	1.43
2	A	410	MAN	C2-C3	4.18	1.58	1.52
2	A	400	GLC	O5-C5	4.26	1.52	1.43
2	A	420	GAL	O5-C5	4.41	1.52	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	420	GAL	C3-C4-C5	-2.29	106.18	110.22
2	A	410	MAN	C1-O5-C5	2.61	115.77	112.17
2	A	410	MAN	O6-C6-C5	3.35	122.62	111.34

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	400	GLC	C1

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	410	MAN	2	0

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	RET	A	250	1	19,20,21	2.12	4 (21%)	27,27,28	1.59	7 (25%)
4	L3P	A	260	-	58,58,58	1.10	6 (10%)	64,73,73	1.20	6 (9%)
5	L2P	A	270	2	45,45,45	1.41	7 (15%)	49,53,53	1.92	14 (28%)
4	L3P	A	280	-	39,39,58	1.93	12 (30%)	44,48,73	0.97	1 (2%)
4	L3P	A	290	-	49,49,58	1.76	15 (30%)	56,60,73	1.23	4 (7%)
4	L3P	A	300	-	49,49,58	1.62	11 (22%)	56,60,73	1.13	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RET	A	250	1	-	0/13/30/31	0/1/1/1
4	L3P	A	260	-	-	0/67/67/67	0/0/0/0
5	L2P	A	270	2	-	0/51/51/51	0/0/0/0
4	L3P	A	280	-	-	0/43/43/67	0/0/0/0
4	L3P	A	290	-	-	0/55/55/67	0/0/0/0
4	L3P	A	300	-	-	0/55/55/67	0/0/0/0

All (55) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	250	RET	C2-C3	-2.29	1.46	1.52
4	A	280	L3P	C12-C11	2.02	1.58	1.50
4	A	260	L3P	P2-O6	2.08	1.66	1.60
4	A	280	L3P	C56-C57	2.08	1.61	1.52
4	A	280	L3P	C52-C53	2.09	1.64	1.52
5	A	270	L2P	C1-C2	2.11	1.56	1.50
4	A	290	L3P	O2-C41	2.12	1.48	1.43
4	A	260	L3P	C6-C5	2.15	1.59	1.51
4	A	260	L3P	C42-C41	2.19	1.58	1.50
4	A	280	L3P	C42-C41	2.20	1.58	1.50
4	A	300	L3P	C14-C13	2.20	1.60	1.52
5	A	270	L2P	O2-C2	2.21	1.49	1.43
4	A	290	L3P	C56-C55	2.21	1.62	1.52
4	A	300	L3P	C55-C53	2.22	1.64	1.52
4	A	300	L3P	C19-C18	2.24	1.60	1.52
4	A	290	L3P	C26-C25	2.27	1.62	1.52
5	A	270	L2P	C47-C48	2.28	1.65	1.52
4	A	300	L3P	C17-C18	2.28	1.65	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	290	L3P	C17-C18	2.31	1.65	1.52
4	A	290	L3P	C55-C53	2.33	1.65	1.52
4	A	300	L3P	C26-C27	2.35	1.62	1.52
4	A	280	L3P	O1-C1	2.35	1.48	1.42
5	A	270	L2P	C25-C23	2.38	1.65	1.52
4	A	300	L3P	O1-C11	2.38	1.52	1.42
4	A	300	L3P	C15-C13	2.40	1.65	1.52
4	A	280	L3P	P1-O3	2.43	1.68	1.60
4	A	290	L3P	C19-C18	2.43	1.60	1.52
4	A	290	L3P	C25-C23	2.44	1.66	1.52
4	A	290	L3P	C14-C13	2.45	1.60	1.52
4	A	280	L3P	C56-C55	2.47	1.63	1.52
4	A	290	L3P	C26-C27	2.48	1.63	1.52
5	A	270	L2P	C42-C41	2.51	1.59	1.50
4	A	260	L3P	O2-C2	2.51	1.50	1.43
4	A	290	L3P	O1-C11	2.54	1.52	1.42
4	A	260	L3P	P1-O3	2.62	1.70	1.59
3	A	250	RET	C7-C6	2.62	1.55	1.45
4	A	290	L3P	C15-C13	2.64	1.67	1.52
4	A	290	L3P	P1-O2P	2.71	1.66	1.54
4	A	300	L3P	P1-O2P	2.71	1.66	1.54
4	A	260	L3P	O2-C41	2.75	1.50	1.43
5	A	270	L2P	O2-C41	2.78	1.50	1.43
4	A	300	L3P	C12-C11	2.95	1.61	1.50
4	A	280	L3P	O2-C41	2.98	1.51	1.43
4	A	290	L3P	C12-C11	2.98	1.61	1.50
4	A	300	L3P	C3-C2	3.16	1.59	1.50
5	A	270	L2P	C26-C25	3.34	1.67	1.52
4	A	290	L3P	C3-C2	3.46	1.60	1.50
4	A	280	L3P	C1-C2	3.77	1.61	1.50
4	A	280	L3P	P1-O2P	3.80	1.70	1.54
4	A	280	L3P	C3-C2	4.13	1.62	1.50
4	A	300	L3P	P1-O3	4.16	1.73	1.60
4	A	290	L3P	P1-O3	4.39	1.74	1.60
3	A	250	RET	C5-C6	4.80	1.42	1.34
4	A	280	L3P	O2-C2	5.39	1.57	1.43
3	A	250	RET	C1-C6	6.26	1.62	1.53

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	270	L2P	C54-C53-C52	-3.15	99.88	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	260	L3P	C46-C45-C43	-2.74	106.73	115.73
4	A	260	L3P	C26-C25-C23	-2.74	106.75	115.73
5	A	270	L2P	C16-C15-C13	-2.67	106.96	115.73
5	A	270	L2P	C16-C17-C18	-2.66	107.01	115.73
4	A	260	L3P	C51-C50-C48	-2.65	107.02	115.73
4	A	300	L3P	C47-C46-C45	-2.51	101.13	113.25
5	A	270	L2P	C30-C28-C29	-2.45	98.90	110.50
5	A	270	L2P	C52-C51-C50	-2.42	101.56	113.25
5	A	270	L2P	C14-C13-C12	-2.40	102.61	111.36
4	A	290	L3P	C47-C46-C45	-2.38	101.75	113.25
3	A	250	RET	C8-C9-C10	-2.35	115.34	118.94
3	A	250	RET	C1-C6-C5	-2.29	119.38	122.59
3	A	250	RET	C7-C6-C5	-2.14	116.43	121.54
5	A	270	L2P	C19-C18-C20	-2.11	103.67	111.36
5	A	270	L2P	C22-C21-C20	-2.08	103.18	113.25
3	A	250	RET	C16-C1-C6	2.05	113.63	110.31
3	A	250	RET	C2-C1-C6	2.06	113.70	110.48
4	A	260	L3P	P2-O6-C6	2.17	124.28	118.30
4	A	300	L3P	C16-C17-C18	2.19	122.94	115.73
5	A	270	L2P	C41-O2-C2	2.28	120.71	115.38
4	A	300	L3P	C21-C20-C18	2.29	123.26	115.73
4	A	300	L3P	O4-P1-O3	2.32	112.91	106.73
4	A	290	L3P	O4-P1-O3	2.42	113.18	106.73
3	A	250	RET	C2-C3-C4	2.48	117.27	111.34
4	A	300	L3P	O1-C1-C2	2.77	115.84	109.36
5	A	270	L2P	C51-C52-C53	2.80	124.92	115.73
4	A	300	L3P	O2P-P1-O3	2.80	114.18	106.73
4	A	290	L3P	O2P-P1-O3	2.91	114.48	106.73
5	A	270	L2P	O3-C3-C2	2.94	119.89	111.76
3	A	250	RET	C1-C6-C7	3.01	124.18	115.73
4	A	260	L3P	C11-O1-C1	3.12	124.81	113.46
5	A	270	L2P	C26-C27-C28	3.13	130.88	115.96
4	A	280	L3P	C51-C50-C48	3.59	127.52	115.73
4	A	260	L3P	C41-O2-C2	4.51	125.93	115.38
4	A	290	L3P	O1-C1-C2	5.10	121.28	109.36
5	A	270	L2P	C29-C28-C27	5.39	146.32	111.50
5	A	270	L2P	O1-C1-C2	6.48	124.50	109.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	250	RET	3	0
4	A	260	L3P	5	0
5	A	270	L2P	2	0
4	A	290	L3P	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

EDS was not executed - this section is therefore empty.