



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 10:09 pm GMT

PDB ID : 2IXQ
Title : THE SOLUTION STRUCTURE OF THE INVASIVE TIP COMPLEX
FROM AFA-DR FIBRILS
Authors : Cota, E.; Jones, C.; Simpson, P.; Altroff, H.; Anderson, K.L.; Du Merle, L.;
Guignot, J.; Servin, A.; Le Bouguenec, C.; Mardon, H.; Matthews, S.
Deposited on : 2006-07-10

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	142	
2	B	143	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4181 atoms, of which 2018 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN AFAD.

Mol	Chain	Residues	Atoms						Trace
1	A	142	Total	C	H	N	O	S	0
			2063	653	988	205	213	4	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	ALA	CONFLICT	UNP Q47038


- Molecule 2 is a protein called AFIMBRIAL ADHESIN AFA-III.

Mol	Chain	Residues	Atoms						Trace
2	B	143	Total	C	H	N	O	S	0
			2118	677	1030	188	220	3	


4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: PROTEIN AFAD



• Molecule 2: AFIMBRIAL ADHESIN AFA-III



5 Refinement protocol and experimental data overview ⓘ

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	8.04	320/1094 (29.3%)	3.63	119/1471 (8.1%)
2	B	8.41	220/1109 (19.8%)	4.31	113/1510 (7.5%)
All	All	8.23	540/2203 (24.5%)	3.99	232/2981 (7.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	4	0
2	B	1	0
All	All	5	0

5 of 540 bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLU	CG-CD	-77.45	0.35	1.51
2	B	2	GLU	CD-OE2	-68.63	0.50	1.25
2	B	50	GLU	CD-OE2	-60.67	0.58	1.25
2	B	1	GLU	CB-CG	-59.42	0.39	1.52
2	B	50	GLU	CD-OE1	-58.85	0.60	1.25

5 of 232 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	50	GLU	OE1-CD-OE2	-61.96	48.95	123.30
2	B	63	ASP	CB-CG-OD1	38.10	152.59	118.30
2	B	66	TYR	CD1-CG-CD2	-34.97	79.44	117.90
2	B	66	TYR	CB-CG-CD2	32.18	140.31	121.00
2	B	66	TYR	CB-CG-CD1	32.09	140.25	121.00

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	67	THR	CB
1	A	140	THR	CB
1	A	142	THR	CB,CA
2	B	1	GLU	CA

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1075	988	989	97
2	B	1088	1030	1023	68
All	All	2163	2018	2012	160

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

5 of 160 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
2:B:19:LEU:CB	2:B:19:LEU:CD1	1.58	1.80
2:B:4:GLN:NE2	2:B:4:GLN:CG	1.48	1.69
1:A:141:VAL:CA	1:A:141:VAL:CG1	1.41	1.97
1:A:138:LYS:NZ	1:A:138:LYS:CD	1.39	1.82
1:A:141:VAL:CG2	1:A:141:VAL:CA	1.38	1.98

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	140/142 (99%)	98 (70%)	34 (24%)	8 (6%)	4 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	141/143 (99%)	112 (79%)	21 (15%)	8 (6%)	4	22
All	All	281/285 (99%)	210 (75%)	55 (20%)	16 (6%)	4	22

5 of 16 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
2	B	103	GLN
1	A	124	ASN
1	A	120	PRO
2	B	19	LEU
1	A	102	GLN

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/111 (100%)	80 (72%)	31 (28%)	2	20
2	B	117/117 (100%)	89 (76%)	28 (24%)	3	27
All	All	228/228 (100%)	169 (74%)	59 (26%)	2	24

5 of 59 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
2	B	137	THR
1	A	142	THR
2	B	115	THR
1	A	16	LEU
2	B	58	SER

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided