



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 03:18 am GMT

PDB ID : 2J0S
Title : THE CRYSTAL STRUCTURE OF THE EXON JUNCTION COMPLEX AT 2.2 Å RESOLUTION
Authors : Bono, F.; Ebert, J.; Lorentzen, E.; Conti, E.
Deposited on : 2006-08-04
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4744 (2.24-2.20)
Clashscore	112137	5509 (2.24-2.20)
Ramachandran outliers	110173	5427 (2.24-2.20)
Sidechain outliers	110143	5428 (2.24-2.20)
RSRZ outliers	101464	4776 (2.24-2.20)
RNA backbone	2435	1022 (2.76-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	
2	C	146	
3	D	89	
4	E	15	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	T	150	<div><div><div></div><div></div><div></div><div></div></div><div>3%23%6%71%</div></div>

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-DEPENDENT RNA HELICASE DDX48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	S	0	8	0
			3138	1986	540	591	21			

- Molecule 2 is a protein called PROTEIN MAGO NASHI HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	143	Total	C	N	O	S	0	2	0
			1192	773	199	217	3			

- Molecule 3 is a protein called RNA-BINDING PROTEIN 8A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	89	Total	C	N	O	S	0	1	0
			710	453	116	138	3			

- Molecule 4 is a RNA chain called 5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*UP*UP*UP*UP*UP*U)-3'.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	6	Total	C	N	O	P	0	0	0
			121	54	12	49	6			

- Molecule 5 is a protein called PROTEIN CASC3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	T	44	Total	C	N	O	0	0	0
			369	228	69	72			

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

- Molecule 7 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $\text{C}_{10}\text{H}_{17}\text{N}_6\text{O}_{12}\text{P}_3$).



- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 8 | A | 168 | Total O
168 168 | 0 | 0 |
| 8 | C | 117 | Total O
117 117 | 0 | 0 |
| 8 | D | 46 | Total O
46 46 | 0 | 0 |
| 8 | E | 4 | Total O
4 4 | 0 | 0 |
| 8 | T | 9 | Total O
9 9 | 0 | 0 |

- Molecule 1: ATP-DEPENDENT RNA HELICASE DDX48



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	169.44Å 169.44Å 71.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.73 – 2.21 43.72 – 2.21	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.73-2.21) 99.7 (43.72-2.21)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.185 , 0.216 0.235 , 0.263	Depositor DCC
R_{free} test set	2918 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	35.6	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.030 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5906	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.62	0/3211	0.75	2/4340 (0.0%)
2	C	0.80	1/1228 (0.1%)	0.84	3/1655 (0.2%)
3	D	0.65	0/730	0.67	0/989
4	E	1.33	1/132 (0.8%)	1.77	3/200 (1.5%)
5	T	0.66	0/379	0.72	0/510
All	All	0.69	2/5680 (0.0%)	0.80	8/7694 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	1	U	OP3-P	-9.43	1.49	1.61
2	C	131	CYS	CB-SG	-6.91	1.70	1.82

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	27	ARG	NE-CZ-NH1	7.26	123.93	120.30
1	A	48	ASP	CB-CG-OD1	6.39	124.05	118.30
4	E	1	U	O4'-C1'-N1	6.33	113.26	108.20
2	C	27	ARG	NE-CZ-NH2	-5.86	117.37	120.30
2	C	82	ARG	NE-CZ-NH2	5.72	123.16	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	177	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3138	0	3166	35	0
2	C	1192	0	1173	8	0
3	D	710	0	662	6	0
4	E	121	0	61	0	0
5	T	369	0	308	4	0
6	A	1	0	0	0	0
7	A	31	0	13	0	0
8	A	168	0	0	0	0
8	C	117	0	0	2	0
8	D	46	0	0	0	0
8	E	4	0	0	0	0
8	T	9	0	0	0	0
All	All	5906	0	5383	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:GLN:HE22	1:A:342:ASP:H	1.29	0.80
1:A:121:GLN:HE21	1:A:324:ARG:HH22	1.35	0.72
1:A:275:LEU:HD23	1:A:348[B]:LEU:CD2	2.18	0.72
2:C:102[A]:SER:OG	8:C:2087:HOH:O	2.08	0.70
1:A:275:LEU:HD23	1:A:348[B]:LEU:HD21	1.74	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/410 (97%)	388 (98%)	6 (2%)	3 (1%)	22	20
2	C	143/146 (98%)	142 (99%)	1 (1%)	0	100	100
3	D	88/89 (99%)	86 (98%)	1 (1%)	1 (1%)	17	13
5	T	40/150 (27%)	36 (90%)	3 (8%)	1 (2%)	6	3
All	All	668/795 (84%)	652 (98%)	11 (2%)	5 (1%)	22	24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	408	ALA
1	A	409	ASP
3	D	139	LEU
1	A	23	MET
5	T	216	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/360 (96%)	324 (94%)	21 (6%)	22	24
2	C	130/134 (97%)	122 (94%)	8 (6%)	21	23
3	D	72/75 (96%)	67 (93%)	5 (7%)	18	18
5	T	35/133 (26%)	33 (94%)	2 (6%)	24	27
All	All	582/702 (83%)	546 (94%)	36 (6%)	22	23

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	337	TRP
1	A	407	VAL
3	D	138	ASP
1	A	348[B]	LEU
2	C	7	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	356	ASN
2	C	13	HIS
3	D	103	HIS
1	A	351	ASN
3	D	90	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	E	5/15 (33%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ANP	A	1413	6	29,33,33	2.29	9 (31%)	28,52,52	2.14	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ANP	A	1413	6	-	0/13/38/38	0/3/3/3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1413	ANP	PG-O2G	-3.36	1.47	1.56
7	A	1413	ANP	PB-O2B	-2.15	1.50	1.56
7	A	1413	ANP	C2-N3	2.12	1.35	1.32
7	A	1413	ANP	C5-C4	2.77	1.46	1.40
7	A	1413	ANP	PG-N3B	2.91	1.71	1.63

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1413	ANP	N3-C2-N1	-7.52	122.31	128.86
7	A	1413	ANP	O1G-PG-N3B	-3.06	107.21	111.79
7	A	1413	ANP	C4-C5-N7	-2.34	107.15	109.41
7	A	1413	ANP	C1'-N9-C4	-2.33	122.61	126.64
7	A	1413	ANP	O3A-PB-N3B	-2.07	100.85	106.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/410 (95%)	0.35	29 (7%) 15 14	41, 47, 57, 85	0
2	C	143/146 (97%)	0.28	1 (0%) 87 86	41, 46, 53, 60	0
3	D	89/89 (100%)	0.46	14 (15%) 2 2	39, 46, 54, 58	0
4	E	6/15 (40%)	-0.60	0 100 100	45, 47, 60, 82	0
5	T	44/150 (29%)	0.39	5 (11%) 6 5	36, 61, 76, 77	0
All	All	673/810 (83%)	0.34	49 (7%) 16 14	36, 47, 62, 85	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	139	LEU	5.8
1	A	275	LEU	4.6
1	A	408	ALA	4.4
3	D	140	MET	4.1
5	T	169	HIS	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	ANP	A	1413	31/31	0.98	0.11	-1.59	28,32,35,37	0
6	MG	A	1412	1/1	0.99	0.07	-	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.