



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 10:01 am GMT

PDB ID : 4J16  
Title : Crystal structure of Thermus thermophilus transhydrogenase heterotrimeric complex of the Alpha1 subunit dimer with the NADP binding domain (domain III) of the Beta subunit  
Authors : Yamaguchi, M.; Leung, J.; Schurig Briccio, L.A.; Gennis, R.B.; Stout, C.D.  
Deposited on : 2013-02-01  
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3709 (2.44-2.40)
Clashscore	112137	4241 (2.44-2.40)
Ramachandran outliers	110173	4178 (2.44-2.40)
Sidechain outliers	110143	4179 (2.44-2.40)
RSRZ outliers	101464	3740 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	<div> <div>9%</div> <div>83%</div> <div>12%</div> <div>...</div> </div>
1	B	381	<div> <div>7%</div> <div>80%</div> <div>14%</div> <div>..</div> </div>
2	C	185	<div> <div>%</div> <div>83%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAD	A	500	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7186 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD/NADP transhydrogenase alpha subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2794	1772	498	512	12			
1	B	369	Total	C	N	O	S	0	0	0
			2769	1756	495	506	12			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-4	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-3	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-2	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-1	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	0	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-5	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-4	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-3	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-2	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-1	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	0	HIS	-	EXPRESSION TAG	UNP Q72GR8

- Molecule 2 is a protein called NAD(P) transhydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	177	Total	C	N	O	S	0	0	0
			1343	860	229	248	6			

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	52	Total O 52 52	0	0
8	B	35	Total O 35 35	0	0
8	C	34	Total O 34 34	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: NAD/NADP transhydrogenase alpha subunit 1



#### • Molecule 1: NAD/NADP transhydrogenase alpha subunit 1



#### • Molecule 2: NAD(P) transhydrogenase subunit beta





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.62Å 75.04Å 198.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.98 – 2.41 39.98 – 2.41	Depositor EDS
% Data completeness (in resolution range)	96.3 (39.98-2.41) 96.3 (39.98-2.41)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.201 , 0.253 0.199 , 0.254	Depositor DCC
$R_{free}$ test set	1918 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.4	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7186	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, PGE, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2840	0.66	2/3852 (0.1%)
1	B	0.35	0/2813	0.59	0/3814
2	C	0.39	0/1366	0.61	0/1848
All	All	0.37	0/7019	0.62	2/9514 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	274	ARG	NE-CZ-NH1	7.42	124.01	120.30
1	A	353	ARG	NE-CZ-NH2	6.36	123.48	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2794	0	2911	28	0
1	B	2769	0	2892	35	0
2	C	1343	0	1378	11	0
3	A	44	0	26	2	0
3	B	44	0	26	2	0
4	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	10	0	14	2	0
6	B	6	0	8	2	0
6	C	6	0	8	0	0
7	C	48	0	25	0	0
8	A	52	0	0	0	0
8	B	35	0	0	0	0
8	C	34	0	0	1	0
All	All	7186	0	7288	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:LEU:HB3	1:B:217:PRO:HA	1.53	0.89
2:C:374:ILE:CD1	2:C:436:ALA:HB1	2.22	0.69
1:B:227:ALA:O	1:B:231:THR:HG23	1.96	0.66
1:B:175:VAL:HG23	1:B:197:ASP:HB2	1.79	0.63
1:B:216:LEU:CB	1:B:217:PRO:HA	2.28	0.63
2:C:374:ILE:HD11	2:C:436:ALA:HB1	1.80	0.61
1:A:353:ARG:HG3	1:A:353:ARG:HH21	1.66	0.60
1:B:114:VAL:HG12	1:B:358:MET:HG2	1.83	0.59
1:A:114:VAL:HG13	1:A:358:MET:HB2	1.85	0.57
1:A:272:VAL:HG11	1:A:294:LEU:HD11	1.86	0.56
1:B:143:ALA:HB2	1:B:284:LEU:HD11	1.87	0.56
1:A:274:ARG:HH11	1:A:274:ARG:HG3	1.69	0.56
1:B:215:GLU:HA	1:B:216:LEU:HD12	1.88	0.56
6:B:404:GOL:H2	2:C:347:GLU:HA	1.86	0.56
1:A:352:VAL:O	1:A:356:LEU:HB2	2.07	0.54
1:A:65:LEU:HD11	1:A:71:LEU:HG	1.90	0.54
1:A:176:GLY:HA3	3:A:500:NAD:H51N	1.90	0.53
1:B:187:LYS:HE2	1:B:210:GLY:O	2.08	0.53
1:A:273:GLU:HG2	1:A:304:ARG:HH22	1.74	0.53
2:C:278:LYS:NZ	2:C:426:ALA:O	2.40	0.53
1:B:285:ALA:O	1:B:289:GLY:O	2.28	0.52
1:A:113:THR:HG23	1:A:359:LYS:HA	1.93	0.51
1:A:353:ARG:CG	1:A:353:ARG:HH21	2.24	0.51
1:A:294:LEU:HD13	1:A:303:VAL:HG13	1.93	0.50
2:C:282:VAL:HG13	8:C:618:HOH:O	2.12	0.49
2:C:280:ILE:HG12	2:C:284:ASP:HB2	1.93	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:VAL:HG12	1:B:358:MET:CG	2.43	0.49
1:B:242:LEU:O	1:B:246:VAL:HB	2.13	0.49
1:B:111:LYS:HA	1:B:361:GLY:CA	2.43	0.49
2:C:358:LEU:HA	2:C:361:ILE:HG22	1.95	0.49
1:B:216:LEU:HB3	1:B:217:PRO:CA	2.35	0.49
1:A:2:VAL:HG23	1:A:69:HIS:HB2	1.95	0.48
1:B:111:LYS:HA	1:B:361:GLY:HA3	1.96	0.47
1:B:216:LEU:CB	1:B:217:PRO:CA	2.91	0.47
1:B:227:ALA:HB1	1:B:231:THR:CG2	2.44	0.47
1:B:238:GLN:O	1:B:242:LEU:HG	2.14	0.47
1:A:143:ALA:HB2	1:A:284:LEU:HD11	1.97	0.46
1:B:219:SER:HB3	1:B:229:GLU:OE1	2.16	0.46
1:B:157:MET:CE	2:C:355:LEU:HD23	2.45	0.46
1:A:274:ARG:HH11	1:A:274:ARG:CG	2.28	0.46
1:B:257:GLN:H	3:B:401:NAD:H52A	1.80	0.46
1:B:314:LEU:N	1:B:315:PRO:CD	2.78	0.46
1:A:155:PHE:N	1:A:156:PRO:HD2	2.30	0.46
1:B:65:LEU:HD21	1:B:71:LEU:HG	1.97	0.45
2:C:380:VAL:HB	2:C:399:VAL:HG11	1.98	0.45
1:B:227:ALA:HB1	1:B:231:THR:HG23	1.98	0.44
1:A:231:THR:O	1:A:235:LYS:HB2	2.17	0.44
2:C:362:ASN:N	2:C:363:PRO:HD2	2.33	0.44
1:B:35:ARG:NH1	1:B:63:GLU:O	2.51	0.44
5:B:403:PGE:H42	5:B:403:PGE:H2	1.72	0.44
1:B:215:GLU:C	1:B:216:LEU:HD12	2.39	0.44
1:A:188:ARG:HG2	6:B:404:GOL:H32	2.00	0.43
1:A:2:VAL:HG22	1:A:3:THR:N	2.33	0.43
1:B:33:ARG:NH2	1:B:57:GLU:OE1	2.52	0.43
1:A:294:LEU:HD12	1:A:308:VAL:HG21	2.02	0.42
1:B:296:LYS:HG3	1:B:301:VAL:HG11	2.01	0.42
1:A:256:ALA:HB1	3:A:500:NAD:H52A	2.02	0.42
1:B:234:GLU:O	1:B:236:ARG:N	2.53	0.42
1:A:6:VAL:CG1	1:A:36:VAL:HG22	2.49	0.41
1:A:188:ARG:HB3	1:B:188:ARG:O	2.20	0.41
1:A:183:ILE:HG23	1:A:193:VAL:HG11	2.02	0.41
1:B:267:LEU:HB2	1:B:292:CYS:HA	2.02	0.41
1:B:28:VAL:HA	1:B:32:ALA:O	2.20	0.41
3:B:401:NAD:H51N	5:B:403:PGE:H22	2.03	0.41
1:A:184:ALA:O	1:A:188:ARG:HG3	2.21	0.41
1:B:239:HIS:HA	1:B:271:MET:CE	2.50	0.41
1:A:218:ILE:O	1:A:238:GLN:NE2	2.53	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:358:MET:HA	1:A:362:GLU:O	2.20	0.41
1:A:188:ARG:HD3	1:B:188:ARG:O	2.21	0.41
1:B:200:LYS:CE	1:B:216:LEU:HD23	2.50	0.41
1:A:276:LYS:HB3	1:A:276:LYS:HE2	1.91	0.40
1:B:142:LEU:HB3	1:B:314:LEU:HG	2.03	0.40
2:C:303:MET:HB2	2:C:374:ILE:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/381 (97%)	352 (95%)	18 (5%)	0	100	100
1	B	365/381 (96%)	345 (94%)	15 (4%)	5 (1%)	13	17
2	C	175/185 (95%)	172 (98%)	3 (2%)	0	100	100
All	All	910/947 (96%)	869 (96%)	36 (4%)	5 (0%)	32	45

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	216	LEU
1	B	228	ARG
1	B	235	LYS
1	B	217	PRO
1	B	215	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/293 (98%)	257 (90%)	30 (10%)	8	11
1	B	285/293 (97%)	259 (91%)	26 (9%)	11	15
2	C	139/146 (95%)	130 (94%)	9 (6%)	20	30
All	All	711/732 (97%)	646 (91%)	65 (9%)	11	15

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	4	VAL
1	A	60	GLU
1	A	71	LEU
1	A	79	GLU
1	A	101	LEU
1	A	114	VAL
1	A	115	ILE
1	A	123	ILE
1	A	124	THR
1	A	149	ARG
1	A	158	LEU
1	A	200	LYS
1	A	203	LEU
1	A	223	GLU
1	A	229	GLU
1	A	230	LEU
1	A	232	GLU
1	A	235	LYS
1	A	261	ARG
1	A	274	ARG
1	A	276	LYS
1	A	303	VAL
1	A	304	ARG
1	A	314	LEU
1	A	333	LEU
1	A	336	LEU
1	A	348	GLU
1	A	353	ARG
1	A	362	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1	MET
1	B	10	ARG
1	B	49	GLU
1	B	71	LEU
1	B	86	GLU
1	B	98	HIS
1	B	99	LYS
1	B	105	ARG
1	B	123	ILE
1	B	125	ARG
1	B	142	LEU
1	B	149	ARG
1	B	175	VAL
1	B	203	LEU
1	B	212	LYS
1	B	228	ARG
1	B	230	LEU
1	B	231	THR
1	B	242	LEU
1	B	246	VAL
1	B	262	ARG
1	B	314	LEU
1	B	333	LEU
1	B	336	LEU
1	B	350	GLU
1	B	364	LEU
2	C	277	LEU
2	C	325	GLU
2	C	355	LEU
2	C	374	ILE
2	C	397	LEU
2	C	427	GLU
2	C	432	LEU
2	C	437	GLN
2	C	447	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	108	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAD	A	500	-	41,48,48	1.31	5 (12%)	43,73,73	1.60	2 (4%)
3	NAD	B	401	-	41,48,48	1.16	3 (7%)	43,73,73	1.64	6 (13%)
5	PGE	B	403	-	9,9,9	0.65	0	8,8,8	0.46	0
6	GOL	B	404	-	5,5,5	0.43	0	5,5,5	0.88	0
7	NAP	C	501	-	44,52,52	1.39	6 (13%)	51,80,80	2.46	12 (23%)
6	GOL	C	502	-	5,5,5	0.39	0	5,5,5	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	500	-	-	0/22/62/62	0/5/5/5
3	NAD	B	401	-	-	0/22/62/62	0/5/5/5
5	PGE	B	403	-	-	0/7/7/7	0/0/0/0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	404	-	-	0/4/4/4	0/0/0/0
7	NAP	C	501	-	-	0/27/67/67	0/5/5/5
6	GOL	C	502	-	-	0/4/4/4	0/0/0/0

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	NAD	C2A-N3A	2.02	1.35	1.32
3	A	500	NAD	PA-O2A	2.05	1.65	1.55
7	C	501	NAP	P2B-O2X	2.22	1.63	1.54
3	A	500	NAD	O4D-C1D	2.28	1.44	1.41
3	A	500	NAD	O4B-C1B	2.35	1.44	1.41
7	C	501	NAP	P2B-O2B	2.37	1.63	1.59
7	C	501	NAP	P2B-O3X	2.39	1.64	1.54
7	C	501	NAP	PA-O1A	3.10	1.62	1.50
3	B	401	NAD	PN-O1N	3.66	1.64	1.50
3	B	401	NAD	PA-O1A	3.70	1.64	1.50
7	C	501	NAP	PN-O1N	3.80	1.65	1.50
3	A	500	NAD	PN-O1N	3.99	1.65	1.50
3	A	500	NAD	PA-O1A	4.02	1.66	1.50
7	C	501	NAP	P2B-O1X	4.16	1.64	1.50

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	501	NAP	N3A-C2A-N1A	-10.27	119.92	128.86
3	A	500	NAD	N3A-C2A-N1A	-8.11	121.79	128.86
3	B	401	NAD	N3A-C2A-N1A	-7.60	122.24	128.86
7	C	501	NAP	O3B-C3B-C4B	-4.07	99.21	111.09
7	C	501	NAP	O7N-C7N-C3N	-3.90	115.06	119.62
3	A	500	NAD	C4A-C5A-N7A	-3.30	106.22	109.41
7	C	501	NAP	O4B-C4B-C3B	-3.24	98.72	105.17
7	C	501	NAP	O3D-C3D-C4D	-3.21	101.71	111.09
3	B	401	NAD	C4A-C5A-N7A	-3.19	106.33	109.41
7	C	501	NAP	C4A-C5A-N7A	-2.85	106.66	109.41
7	C	501	NAP	O4B-C1B-C2B	-2.85	101.61	106.59
7	C	501	NAP	O7N-C7N-N7N	-2.82	118.57	122.58
3	B	401	NAD	C4D-O4D-C1D	-2.72	106.87	109.77
3	B	401	NAD	O4B-C4B-C5B	-2.35	101.46	109.40
7	C	501	NAP	O2D-C2D-C3D	-2.33	104.37	111.83
3	B	401	NAD	C5D-C4D-C3D	-2.19	106.95	115.29
3	B	401	NAD	C5B-C4B-C3B	-2.09	107.31	115.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	501	NAP	C2B-C3B-C4B	2.74	108.18	101.95
7	C	501	NAP	C4D-O4D-C1D	4.67	114.74	109.77
7	C	501	NAP	C3N-C7N-N7N	7.52	126.36	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	NAD	2	0
3	B	401	NAD	2	0
5	B	403	PGE	2	0
6	B	404	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	372/381 (97%)	0.50	36 (9%) <b>8</b> <b>7</b>	39, 70, 103, 120	0
1	B	369/381 (96%)	0.36	26 (7%) <b>17</b> <b>15</b>	42, 77, 114, 146	0
2	C	177/185 (95%)	-0.16	1 (0%) <b>89</b> <b>87</b>	40, 59, 79, 96	0
All	All	918/947 (96%)	0.31	63 (6%) <b>18</b> <b>15</b>	39, 69, 107, 146	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	ARG	6.0
1	B	230	LEU	5.4
1	A	112	ALA	3.7
1	B	373	GLY	3.6
1	A	371	LEU	3.5
1	A	62	GLY	3.4
1	A	370	ALA	3.2
1	B	99	LYS	3.2
1	A	185	THR	3.2
1	A	304	ARG	3.2
1	B	189	LEU	3.1
1	B	242	LEU	3.1
1	A	189	LEU	3.1
1	B	215	GLU	3.0
1	A	145	ILE	2.9
1	A	138	VAL	2.8
1	A	363	VAL	2.8
1	A	186	ALA	2.8
1	A	188	ARG	2.8
1	A	261	ARG	2.8
1	A	158	LEU	2.8
1	A	141	TYR	2.7
1	A	184	ALA	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	224	GLY	2.7
1	A	159	THR	2.7
1	A	360	GLU	2.6
1	A	1	MET	2.5
1	A	364	LEU	2.5
2	C	337	MET	2.5
1	B	219	SER	2.5
1	A	348	GLU	2.5
1	A	157	MET	2.4
1	B	262	ARG	2.4
1	A	137	THR	2.4
1	B	144	ALA	2.4
1	A	193	VAL	2.3
1	B	141	TYR	2.3
1	A	38	ARG	2.3
1	B	234	GLU	2.3
1	A	101	LEU	2.3
1	B	265	ILE	2.2
1	B	186	ALA	2.2
1	B	216	LEU	2.2
1	A	109	ALA	2.2
1	B	360	GLU	2.2
1	B	289	GLY	2.1
1	A	190	GLY	2.1
1	B	322	ALA	2.1
1	A	269	GLU	2.1
1	B	363	VAL	2.1
1	B	105	ARG	2.1
1	B	195	ALA	2.1
1	B	228	ARG	2.1
1	B	218	ILE	2.1
1	A	177	VAL	2.0
1	B	142	LEU	2.0
1	A	140	GLY	2.0
1	A	181	MET	2.0
1	A	156	PRO	2.0
1	B	55	GLY	2.0
1	B	145	ILE	2.0
1	A	154	PHE	2.0
1	B	340	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAD	A	500	44/44	0.66	0.40	4.51	103,120,160,164	0
6	GOL	C	502	6/6	0.87	0.19	0.74	83,89,90,92	0
3	NAD	B	401	44/44	0.94	0.18	0.06	67,76,91,94	0
5	PGE	B	403	10/10	0.88	0.15	0.00	65,72,75,78	0
7	NAP	C	501	48/48	0.97	0.15	-0.10	43,47,60,63	0
6	GOL	B	404	6/6	0.85	0.24	-0.13	53,57,60,61	0
4	CL	B	402	1/1	0.94	0.27	-	83,83,83,83	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.