



wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Mar 2, 2017 – 11:29 am GMT

PDB ID : 3J2U
EMDB ID: : EMD-5565
Title : Kinesin-13 KLP10A HD in complex with CS-tubulin and a microtubule
Authors : Asenjo, A.B.; Chatterjee, C.; Tan, D.; DePaoli, V.; Rice, W.J.; Diaz-Avalos, R.; Silvestry, M.; Sosa, H.
Deposited on : 2013-01-10
Resolution : 10.80 Å(reported)
Based on PDB ID : 1JFF

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report
for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : recalc29047


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 10.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 125131 | 1336 |
| Ramachandran outliers | 121729 | 1120 |
| Sidechain outliers | 121581 | 1026 |

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | K | 374 | <div> <div>27%</div> <div>52%</div> <div>•</div> <div>17%</div> </div> |
| 2 | A | 451 | <div> <div>18%</div> <div>55%</div> <div>17%</div> <div>•</div> <div>9%</div> </div> |
| 2 | C | 451 | <div> <div>17%</div> <div>56%</div> <div>17%</div> <div>•</div> <div>9%</div> </div> |
| 3 | B | 445 | <div> <div>21%</div> <div>55%</div> <div>18%</div> <div>•</div> <div>•</div> </div> |
| 3 | D | 445 | <div> <div>21%</div> <div>55%</div> <div>18%</div> <div>•</div> <div>•</div> </div> |

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin-like protein Klp10A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1 | K | 309 | Total | C | N | O | S | 0 | 0 |
| | | | 2419 | 1525 | 432 | 446 | 16 | | |

There are 37 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| K | 242 | MET | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 243 | ARG | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 244 | GLY | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 245 | SER | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 246 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 247 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 248 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 249 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 250 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 251 | HIS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 252 | GLY | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 253 | MET | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 254 | ALA | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 255 | SER | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 256 | MET | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 257 | THR | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 258 | GLY | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 259 | GLY | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 260 | GLN | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 261 | GLN | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 262 | MET | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 263 | GLY | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 264 | ARG | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 265 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 266 | LEU | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 267 | TYR | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 268 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 269 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| K | 270 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 271 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 272 | LYS | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 273 | ASP | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 274 | PRO | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 275 | SER | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 276 | SER | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 277 | ARG | - | EXPRESSION TAG | UNP Q960Z0 |
| K | 278 | SER | - | EXPRESSION TAG | UNP Q960Z0 |

- Molecule 2 is a protein called Tubulin alpha-1A chain.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 2 | A | 412 | Total | C | N | O | S | 0 | 0 |
| | | | 3227 | 2043 | 551 | 613 | 20 | | |
| 2 | C | 412 | Total | C | N | O | S | 0 | 0 |
| | | | 3227 | 2043 | 551 | 613 | 20 | | |

There are 10 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| A | 136 | SER | LEU | CONFLICT | UNP P81947 |
| A | 232 | GLY | SER | CONFLICT | UNP P81947 |
| A | 265 | GLY | ILE | CONFLICT | UNP P81947 |
| A | 340 | THR | SER | CONFLICT | UNP P81947 |
| A | 358 | GLU | GLN | CONFLICT | UNP P81947 |
| C | 136 | SER | LEU | CONFLICT | UNP P81947 |
| C | 232 | GLY | SER | CONFLICT | UNP P81947 |
| C | 265 | GLY | ILE | CONFLICT | UNP P81947 |
| C | 340 | THR | SER | CONFLICT | UNP P81947 |
| C | 358 | GLU | GLN | CONFLICT | UNP P81947 |

- Molecule 3 is a protein called Tubulin beta-2B chain.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 3 | B | 426 | Total | C | N | O | S | 0 | 0 |
| | | | 3351 | 2105 | 575 | 646 | 25 | | |
| 3 | D | 426 | Total | C | N | O | S | 0 | 0 |
| | | | 3351 | 2105 | 575 | 646 | 25 | | |


There are 12 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| B | 57 | ALA | THR | CONFLICT | UNP Q6B856 |
| B | 172 | VAL | MET | CONFLICT | UNP Q6B856 |
| B | 298 | ALA | SER | CONFLICT | UNP Q6B856 |
| B | 318 | VAL | ILE | CONFLICT | UNP Q6B856 |
| B | 450 | GLY | GLU | CONFLICT | UNP Q6B856 |
| B | 451 | GLU | GLY | CONFLICT | UNP Q6B856 |
| D | 57 | ALA | THR | CONFLICT | UNP Q6B856 |
| D | 172 | VAL | MET | CONFLICT | UNP Q6B856 |
| D | 298 | ALA | SER | CONFLICT | UNP Q6B856 |
| D | 318 | VAL | ILE | CONFLICT | UNP Q6B856 |
| D | 450 | GLY | GLU | CONFLICT | UNP Q6B856 |
| D | 451 | GLU | GLY | CONFLICT | UNP Q6B856 |


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kinesin-like protein Klp10A




• Molecule 2: Tubulin alpha-1A chain






• Molecule 2: Tubulin alpha-1A chain


Chain C: 17% 56% 17% 9%



• Molecule 3: Tubulin beta-2B chain

Chain B: 21% 55% 18%





4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| Reconstruction method | HELICAL | Depositor |
| Imposed symmetry | POINT, Not provided | Depositor |
| Number of segments used | 54584 | Depositor |
| Resolution determination method | FSC calculated from two independently re-fined maps at 0.143 cut-off. | Depositor |
| CTF correction method | each particle | Depositor |
| Microscope | JEOL 3200FSC, FEI TECNAI F20 | Depositor |
| Voltage (kV) | 300, 200 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | Not provided, Not provided | Depositor |
| Minimum defocus (nm) | Not provided, Not provided | Depositor |
| Maximum defocus (nm) | Not provided, Not provided | Depositor |
| Magnification | Not provided, Not provided | Depositor |
| Image detector | Not provided | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-------------|-------------|----------------|
| | | RMSZ | $\# Z > 2$ | RMSZ | $\# Z > 2$ |
| 1 | K | 0.93 | 0/2453 | 1.10 | 1/3290 (0.0%) |
| 2 | A | 0.51 | 0/3300 | 0.73 | 0/4482 |
| 2 | C | 0.51 | 0/3300 | 0.73 | 0/4482 |
| 3 | B | 0.51 | 0/3426 | 0.76 | 2/4642 (0.0%) |
| 3 | D | 0.51 | 0/3426 | 0.76 | 2/4642 (0.0%) |
| All | All | 0.59 | 0/15905 | 0.81 | 5/21538 (0.0%) |

There are no bond length outliers.

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 3 | D | 235 | MET | CG-SD-CE | 6.08 | 109.92 | 100.20 |
| 3 | B | 235 | MET | CG-SD-CE | 6.07 | 109.92 | 100.20 |
| 1 | K | 384 | GLY | N-CA-C | -5.47 | 99.43 | 113.10 |
| 3 | D | 217 | LEU | N-CA-C | -5.36 | 96.54 | 111.00 |
| 3 | B | 217 | LEU | N-CA-C | -5.35 | 96.56 | 111.00 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | K | 2419 | 0 | 2487 | 389 | 0 |
| 2 | A | 3227 | 0 | 3143 | 520 | 0 |
| 2 | C | 3227 | 0 | 3143 | 569 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 3 | B | 3351 | 0 | 3229 | 563 | 0 |
| 3 | D | 3351 | 0 | 3229 | 568 | 0 |
| All | All | 15575 | 0 | 15231 | 2506 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 81.

The worst 5 of 2506 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|-----------------|--------------------------|-------------------|
| 1:K:444:ASP:HA | 3:B:419:THR:CG2 | 1.37 | 1.48 |
| 1:K:444:ASP:N | 3:B:416:MET:HE1 | 1.26 | 1.43 |
| 1:K:444:ASP:N | 3:B:416:MET:CE | 1.87 | 1.37 |
| 1:K:444:ASP:CA | 3:B:419:THR:CG2 | 2.01 | 1.35 |
| 2:C:407:TRP:CZ2 | 3:D:257:VAL:HA | 1.60 | 1.34 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-----------|-------------|----|
| 1 | K | 303/374 (81%) | 285 (94%) | 12 (4%) | 6 (2%) | 9 | 46 |
| 2 | A | 408/451 (90%) | 265 (65%) | 84 (21%) | 59 (14%) | 0 | 5 |
| 2 | C | 408/451 (90%) | 266 (65%) | 83 (20%) | 59 (14%) | 0 | 5 |
| 3 | B | 424/445 (95%) | 274 (65%) | 94 (22%) | 56 (13%) | 0 | 6 |
| 3 | D | 424/445 (95%) | 274 (65%) | 94 (22%) | 56 (13%) | 0 | 6 |
| All | All | 1967/2166 (91%) | 1364 (69%) | 367 (19%) | 236 (12%) | 1 | 7 |

5 of 236 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | K | 501 | GLN |
| 1 | K | 503 | SER |
| 1 | K | 506 | ILE |
| 2 | A | 96 | LYS |
| 2 | A | 97 | GLU |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 1 | K | 269/323 (83%) | 261 (97%) | 8 (3%) | 46 | 72 |
| 2 | A | 347/377 (92%) | 298 (86%) | 49 (14%) | 4 | 22 |
| 2 | C | 347/377 (92%) | 298 (86%) | 49 (14%) | 4 | 22 |
| 3 | B | 367/381 (96%) | 308 (84%) | 59 (16%) | 3 | 17 |
| 3 | D | 367/381 (96%) | 307 (84%) | 60 (16%) | 3 | 17 |
| All | All | 1697/1839 (92%) | 1472 (87%) | 225 (13%) | 8 | 24 |

5 of 225 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | B | 324 | SER |
| 2 | C | 125 | LEU |
| 3 | D | 306 | ASP |
| 3 | B | 344 | VAL |
| 3 | B | 432 | TYR |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | B | 331 | GLN |
| 2 | C | 11 | GLN |
| 3 | D | 337 | ASN |
| 3 | B | 334 | ASN |
| 3 | B | 349 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.