



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 06:15 pm GMT

PDB ID : 1J3Z
Title : Direct observation of photolysis-induced tertiary structural changes in human haemoglobin; Crystal structure of alpha(Fe-CO)-beta(Ni) hemoglobin (laser unphotolysed)
Authors : Adachi, S.; Park, S.-Y.; Tame, J.R.H.; Shiro, Y.; Shibayama, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2003-02-21
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.








Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	141	 95% . .
1	C	141	 96% .
1	E	141	 90% 7% . .
1	G	141	 96% .
2	B	146	 92% 6% .
2	D	146	 89% 11%
2	F	146	 91% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	146	 <div>90%10%</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CMO	A	143	-	-	X	-
4	CMO	G	143	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10467 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemoglobin alpha Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			
1	C	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			
1	E	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			
1	G	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			

- Molecule 2 is a protein called Hemoglobin beta Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1123	724	195	201	3			
2	D	146	Total	C	N	O	S	0	0	0
			1123	724	195	201	3			
2	F	146	Total	C	N	O	S	0	0	0
			1123	724	195	201	3			
2	H	146	Total	C	N	O	S	0	0	0
			1123	724	195	201	3			

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	G	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 2	C 1	O 1	0	0
4	C	1	Total 2	C 1	O 1	0	0
4	E	1	Total 2	C 1	O 1	0	0
4	G	1	Total 2	C 1	O 1	0	0

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING NI(II) (three-letter code: HNI) (formula: $\text{C}_{34}\text{H}_{32}\text{N}_4\text{NiO}_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
5	D	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
5	F	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
5	H	1	Total 43	C 34	N 4	Ni 1	O 4	0	0

- Molecule 6 is BUT-2-ENEDIAL (three-letter code: 2FU) (formula: $C_4H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	4	2		
6	F	1	Total	C	O	0	0
			6	4	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	190	Total	O	0	0
			190	190		
7	B	149	Total	O	0	0
			149	149		
7	C	171	Total	O	0	0
			171	171		
7	D	142	Total	O	0	0
			142	142		
7	E	176	Total	O	0	0
			176	176		
7	F	204	Total	O	0	0
			204	204		
7	G	160	Total	O	0	0
			160	160		
7	H	143	Total	O	0	0
			143	143		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Hemoglobin alpha Chain

Chain A: 





- Molecule 1: Hemoglobin alpha Chain

Chain C: 



- Molecule 1: Hemoglobin alpha Chain

Chain E: 



- Molecule 1: Hemoglobin alpha Chain

Chain G: 




- Molecule 2: Hemoglobin beta Chain


Chain B: 




- Molecule 2: Hemoglobin beta Chain



● Molecule 2: Hemoglobin beta Chain



● Molecule 2: Hemoglobin beta Chain



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.06Å 93.70Å 99.21Å 90.00° 101.14° 90.00°	Depositor
Resolution (Å)	20.00 – 1.60	Depositor
% Data completeness (in resolution range)	94.4 (20.00-1.60)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.159 , 0.196	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10467	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HNI, 2FU, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/1097	0.70	1/1491 (0.1%)
1	C	0.48	0/1097	0.70	1/1491 (0.1%)
1	E	0.49	0/1097	0.75	3/1491 (0.2%)
1	G	0.53	0/1097	0.71	0/1491
2	B	0.53	0/1153	0.75	3/1566 (0.2%)
2	D	0.49	0/1153	0.69	2/1566 (0.1%)
2	F	0.55	0/1153	0.73	1/1566 (0.1%)
2	H	0.53	0/1153	0.75	1/1566 (0.1%)
All	All	0.51	0/9000	0.72	12/12228 (0.1%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	134	VAL	CG1-CB-CG2	7.34	122.65	110.90
2	H	52	ASP	CB-CG-OD2	6.90	124.51	118.30
1	E	105	LEU	CA-CB-CG	6.65	130.59	115.30
1	C	74	ASP	CB-CG-OD2	5.85	123.56	118.30
2	B	68	LEU	CB-CG-CD2	5.53	120.39	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	5	0
1	C	1069	0	1073	3	0
1	E	1069	0	1073	7	0
1	G	1069	0	1073	8	0
2	B	1123	0	1116	10	0
2	D	1123	0	1116	10	0
2	F	1123	0	1116	9	0
2	H	1123	0	1116	11	0
3	A	43	0	30	4	0
3	C	43	0	30	0	0
3	E	43	0	30	1	0
3	G	43	0	30	7	0
4	A	2	0	0	4	0
4	C	2	0	0	0	0
4	E	2	0	0	0	0
4	G	2	0	0	4	0
5	B	43	0	30	0	0
5	D	43	0	30	1	0
5	F	43	0	30	1	0
5	H	43	0	30	0	0
6	B	6	0	2	0	0
6	F	6	0	2	0	0
7	A	190	0	0	1	0
7	B	149	0	0	5	0
7	C	171	0	0	1	0
7	D	142	0	0	2	0
7	E	176	0	0	1	0
7	F	204	0	0	3	0
7	G	160	0	0	1	0
7	H	143	0	0	3	0
All	All	10467	0	9000	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:146:HIS:HA	7:F:2098:HOH:O	1.62	1.00
1:E:113:LEU:HB3	1:E:116:GLU:HG2	1.44	1.00
2:D:107:GLY:HA3	2:D:134:VAL:HG11	1.41	0.99
2:D:107:GLY:CA	2:D:134:VAL:HG11	1.95	0.97

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:107:GLY:HA3	2:D:134:VAL:CG1	2.03	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
1	C	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
1	E	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
1	G	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
2	B	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
2	D	144/146 (99%)	143 (99%)	1 (1%)	0	100	100
2	F	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
2	H	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
All	All	1132/1148 (99%)	1115 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	112 (99%)	1 (1%)	82	69
1	C	113/113 (100%)	112 (99%)	1 (1%)	82	69
1	E	113/113 (100%)	107 (95%)	6 (5%)	26	6
1	G	113/113 (100%)	113 (100%)	0	100	100
2	B	118/118 (100%)	114 (97%)	4 (3%)	42	15
2	D	118/118 (100%)	113 (96%)	5 (4%)	34	11
2	F	118/118 (100%)	113 (96%)	5 (4%)	34	11
2	H	118/118 (100%)	112 (95%)	6 (5%)	28	7
All	All	924/924 (100%)	896 (97%)	28 (3%)	46	18

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	83	LEU
1	E	105	LEU
2	H	78	LEU
1	E	90	LYS
1	E	99	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	9	ASN
1	E	97	ASN
2	H	117	HIS
1	E	72	HIS
2	F	77	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	A	142	1,4	28,50,50	2.03	5 (17%)	17,82,82	1.64	3 (17%)
4	CMO	A	143	3	0,1,1	0.00	-	0,0,0	0.00	-
5	HNI	B	147	2	28,50,50	3.84	14 (50%)	20,82,82	5.79	10 (50%)
6	2FU	B	2001	2	5,5,5	0.59	0	4,4,4	1.87	2 (50%)
3	HEM	C	142	1,4	28,50,50	2.05	6 (21%)	17,82,82	1.81	6 (35%)
4	CMO	C	143	3	0,1,1	0.00	-	0,0,0	0.00	-
5	HNI	D	147	2	28,50,50	3.89	15 (53%)	20,82,82	5.56	10 (50%)
3	HEM	E	142	1,4	28,50,50	2.13	7 (25%)	17,82,82	2.13	5 (29%)
4	CMO	E	143	3	0,1,1	0.00	-	0,0,0	0.00	-
5	HNI	F	147	2	28,50,50	3.91	15 (53%)	20,82,82	5.71	9 (45%)
6	2FU	F	2002	2	5,5,5	0.71	0	4,4,4	1.52	1 (25%)
3	HEM	G	142	1,4	28,50,50	2.14	8 (28%)	17,82,82	1.71	4 (23%)
4	CMO	G	143	3	0,1,1	0.00	-	0,0,0	0.00	-
5	HNI	H	147	2	28,50,50	3.92	16 (57%)	20,82,82	5.80	11 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	142	1,4	-	0/6/54/54	0/0/8/8
4	CMO	A	143	3	-	0/0/0/0	0/0/0/0
5	HNI	B	147	2	-	0/8/94/94	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	2FU	B	2001	2	-	2/3/3/3	0/0/0/0
3	HEM	C	142	1,4	-	0/6/54/54	0/0/8/8
4	CMO	C	143	3	-	0/0/0/0	0/0/0/0
5	HNI	D	147	2	-	0/8/94/94	0/0/8/8
3	HEM	E	142	1,4	-	0/6/54/54	0/0/8/8
4	CMO	E	143	3	-	0/0/0/0	0/0/0/0
5	HNI	F	147	2	-	0/8/94/94	0/0/8/8
6	2FU	F	2002	2	-	2/3/3/3	0/0/0/0
3	HEM	G	142	1,4	-	0/6/54/54	0/0/8/8
4	CMO	G	143	3	-	0/0/0/0	0/0/0/0
5	HNI	H	147	2	-	0/8/94/94	0/0/8/8

The worst 5 of 86 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	147	HNI	CHD-C4C	-8.17	1.37	1.53
5	B	147	HNI	CHC-C1C	-8.06	1.37	1.53
5	F	147	HNI	CHC-C1C	-7.98	1.38	1.53
5	H	147	HNI	CHD-C4C	-7.88	1.38	1.53
5	H	147	HNI	CHC-C1C	-7.85	1.38	1.53

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	142	HEM	CAA-CBA-CGA	-5.91	102.57	112.66
5	D	147	HNI	CHC-C4B-C3B	-4.51	123.49	129.55
5	F	147	HNI	CHC-C4B-C3B	-4.14	123.98	129.55
5	H	147	HNI	CHD-C1D-C2D	-3.98	122.56	129.45
3	G	142	HEM	CBD-CAD-C3D	-3.86	105.10	112.47

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	2001	2FU	O8-C7-C5-C1
6	F	2002	2FU	O3-C2-C1-C5
6	F	2002	2FU	O8-C7-C5-C1
6	B	2001	2FU	O3-C2-C1-C5

There are no ring outliers.

7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	142	HEM	4	0
4	A	143	CMO	4	0
5	D	147	HNI	1	0
3	E	142	HEM	1	0
5	F	147	HNI	1	0
3	G	142	HEM	7	0
4	G	143	CMO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.