



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2018 – 09:23 AM EST

PDB ID : 5J42
Title : Crystal structure of m2hTDP2-CAT in complex with a small molecule inhibitor
Authors : Hornyak, P.; Pearl, L.H.; Caldecott, K.W.; Oliver, A.W.
Deposited on : 2016-03-31
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div> <div>5%</div> <div>82%</div> <div>13%</div> <div>• •</div> </div>
1	B	256	<div> <div>7%</div> <div>78%</div> <div>18%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	404	-	-	X	X
4	GOL	B	406	-	-	-	X
5	EDO	A	405	-	-	-	X
5	EDO	B	407	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4803 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosyl-DNA phosphodiesterase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	12	0
			1987	1270	339	357	21			
1	B	249	Total	C	N	O	S	0	9	0
			1981	1268	344	351	18			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	115	SER	-	expression tag	UNP Q9JJX7
A	116	ASN	-	expression tag	UNP Q9JJX7
A	117	ALA	-	expression tag	UNP Q9JJX7
A	242	GLY	GLU	engineered mutation	UNP Q9JJX7
A	278	ARG	GLN	engineered mutation	UNP Q9JJX7
A	321	CYS	TYR	engineered mutation	UNP Q9JJX7
A	323	LEU	HIS	engineered mutation	UNP Q9JJX7
B	115	SER	-	expression tag	UNP Q9JJX7
B	116	ASN	-	expression tag	UNP Q9JJX7
B	117	ALA	-	expression tag	UNP Q9JJX7
B	242	GLY	GLU	engineered mutation	UNP Q9JJX7
B	278	ARG	GLN	engineered mutation	UNP Q9JJX7
B	321	CYS	TYR	engineered mutation	UNP Q9JJX7
B	323	LEU	HIS	engineered mutation	UNP Q9JJX7

- Molecule 2 is 10-(4-hydroxyphenyl)-2,4-dioxo-2,3,4,10-tetrahydropyrimido[4,5-b]quinoline-8-carbonitrile (three-letter code: 6FV) (formula: C₁₈H₁₀N₄O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			25	18	4	3		
2	B	1	Total	C	N	O	0	0
			25	18	4	3		

- Molecule 3 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		
3	A	1	Total	C	O	0	0
			10	4	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	1
			20	8	12		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mn	0	0
			1	1		
6	A	1	Total	Mn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	356	Total	O	0	0
			356	356		
7	B	337	Total	O	0	0
			337	337		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosyl-DNA phosphodiesterase 2



• Molecule 1: Tyrosyl-DNA phosphodiesterase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.05Å 42.87Å 109.05Å 90.00° 94.05° 90.00°	Depositor
Resolution (Å)	42.07 – 1.70 42.07 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.1 (42.07-1.70) 99.1 (42.07-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 1.70Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.160 , 0.197 0.154 , 0.195	Depositor DCC
R_{free} test set	3137 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4803	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TLA, MN, EDO, 6FV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	3/2065 (0.1%)	0.96	4/2795 (0.1%)
1	B	0.93	2/2055 (0.1%)	0.93	5/2783 (0.2%)
All	All	0.96	5/4120 (0.1%)	0.95	9/5578 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	165	PRO	N-CD	-15.12	1.26	1.47
1	B	168[A]	CYS	CB-SG	-11.63	1.62	1.82
1	B	168[B]	CYS	CB-SG	-11.63	1.62	1.82
1	A	170	TYR	CB-CG	-7.72	1.40	1.51
1	A	168	CYS	CB-SG	-6.16	1.71	1.82

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	316	ARG	NE-CZ-NH1	-8.79	115.91	120.30
1	B	214	MET	CG-SD-CE	7.65	112.44	100.20
1	A	160	LEU	CB-CG-CD1	-6.75	99.53	111.00
1	A	278	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	B	249	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	276	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	B	308	ASP	CB-CG-OD1	5.56	123.30	118.30
1	A	218	LEU	CB-CG-CD2	-5.18	102.19	111.00
1	B	249	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1987	0	2010	32	0
1	B	1981	0	1996	43	0
2	A	25	0	0	1	0
2	B	25	0	0	1	0
3	A	20	0	7	1	0
3	B	50	0	19	1	0
4	A	6	0	8	9	0
4	B	6	0	8	1	0
5	A	4	0	6	1	0
5	B	4	0	6	2	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	356	0	0	13	0
7	B	337	0	0	9	2
All	All	4803	0	4060	78	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:MET:HE3	1:B:239:SER:H	1.13	1.07
4:A:404:GOL:O1	7:A:501:HOH:O	1.82	0.98
1:A:144:ARG:HG3	1:A:170:TYR:OH	1.61	0.97
1:B:214:MET:HE2	1:B:238:GLU:HG3	1.51	0.92
1:B:214:MET:CE	1:B:239:SER:H	1.82	0.91
1:B:214:MET:HE3	1:B:239:SER:N	1.92	0.84
1:B:340:GLN:HE21	1:B:367:ASN:HD21	1.27	0.83
1:A:220[B]:CYS:SG	7:A:520:HOH:O	2.38	0.79
4:A:404:GOL:O3	7:A:502:HOH:O	2.06	0.73
1:A:252:LYS:HE3	4:A:404:GOL:O1	1.90	0.71
1:B:214:MET:CE	1:B:239:SER:N	2.53	0.69
1:A:278:ARG:HH11	1:A:278:ARG:HG3	1.59	0.68
1:B:204:GLN:NE2	7:B:502:HOH:O	2.26	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:GLN:OE1	4:A:404:GOL:H32	1.94	0.68
1:B:196:LYS:NZ	7:B:503:HOH:O	2.28	0.66
1:B:214:MET:CE	1:B:238:GLU:HG3	2.24	0.63
4:A:404:GOL:O3	7:A:503:HOH:O	2.14	0.63
1:B:340:GLN:NE2	1:B:367:ASN:HD21	1.98	0.61
1:B:305:TYR:HB2	5:B:407:EDO:H22	1.84	0.59
1:B:310:LYS:NZ	7:B:507:HOH:O	2.37	0.57
1:B:344:LEU:HD11	5:B:407:EDO:H11	1.85	0.57
1:A:340:GLN:CD	1:A:367:ASN:HD21	2.09	0.54
1:B:337:LEU:CD1	1:B:368:VAL:HG22	2.38	0.53
1:A:287:PRO:HG3	4:A:404:GOL:H11	1.89	0.53
1:A:301:LYS:HD3	7:A:676:HOH:O	2.08	0.53
1:B:261:ALA:HB2	7:B:564:HOH:O	2.10	0.52
1:A:337:LEU:HA	1:A:367:ASN:O	2.10	0.52
1:A:125[A]:SER:HB3	1:A:365[A]:THR:HG22	1.93	0.52
1:A:340:GLN:CD	1:A:367:ASN:ND2	2.63	0.51
1:B:170:TYR:CZ	1:B:174:ARG:HG3	2.45	0.51
1:B:214:MET:HE1	1:B:239:SER:O	2.10	0.51
1:B:241:ARG:HH22	3:B:405[A]:TLA:C1	2.23	0.51
1:A:351:ASP:HB3	7:A:757:HOH:O	2.09	0.51
1:A:255:LEU:HB2	4:A:404:GOL:H12	1.93	0.50
1:B:124:ILE:HD12	1:B:225:LEU:HD23	1.93	0.50
2:A:401:6FV:N4	1:B:318:PRO:HD2	2.27	0.49
1:A:298:GLY:N	7:A:507:HOH:O	2.45	0.49
1:A:200:LYS:HE2	7:A:504:HOH:O	2.13	0.48
1:B:243[A]:HIS:HD2	7:B:724:HOH:O	1.95	0.48
1:B:207:ILE:HD11	1:B:257:LYS:HE2	1.94	0.48
1:B:336:HIS:HA	1:B:369:VAL:HG12	1.96	0.48
1:A:243:HIS:HD2	7:A:781:HOH:O	1.96	0.48
1:B:294:TRP:CZ2	1:B:304:GLN:HG3	2.49	0.48
1:B:347:LEU:HD22	1:B:360:TRP:CE3	2.48	0.48
1:B:339:PRO:HA	1:B:366:LEU:HD23	1.96	0.48
1:A:259:GLN:OE1	4:A:404:GOL:C3	2.62	0.47
1:A:222:ASN:ND2	7:A:520:HOH:O	2.48	0.47
1:B:202:LYS:HB3	1:B:202:LYS:HE2	1.74	0.47
1:B:154:SER:HB2	7:B:690:HOH:O	2.15	0.46
1:B:323:LEU:HA	7:B:504:HOH:O	2.15	0.46
1:A:276:ARG:HA	5:A:405:EDO:H12	1.96	0.46
1:A:233:MET:HE1	1:A:257:LYS:HG3	1.97	0.46
1:B:218:LEU:HD11	1:B:233:MET:HE2	1.97	0.46
1:A:291:PHE:HE2	1:A:332:ALA:HB2	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179[B]:THR:OG1	1:B:196:LYS:HE3	2.16	0.45
1:A:278:ARG:NH1	1:A:278:ARG:HG3	2.29	0.45
1:B:337:LEU:HD12	1:B:337:LEU:HA	1.82	0.45
1:A:318:PRO:HD2	2:B:401:6FV:N4	2.31	0.45
1:B:224:SER:HB3	1:B:229:GLU:OE1	2.17	0.45
1:A:127:ILE:HA	1:A:362:LEU:O	2.17	0.44
1:A:164:ILE:HB	1:A:165:PRO:HD2	1.98	0.44
1:B:252:LYS:HD2	7:B:710:HOH:O	2.17	0.44
1:B:209:PHE:HA	4:B:406:GOL:H32	2.00	0.43
1:B:339:PRO:HA	1:B:366:LEU:CD2	2.49	0.43
1:B:337:LEU:HD21	1:B:366:LEU:HD13	2.00	0.43
1:A:365[A]:THR:HG23	7:A:606:HOH:O	2.17	0.43
1:A:162:GLU:OE1	3:A:403:TLA:O3	2.37	0.42
1:B:230:PHE:HZ	1:B:368:VAL:HG21	1.83	0.42
1:A:207:ILE:HB	1:A:218:LEU:HB3	2.00	0.42
1:A:252:LYS:HD2	1:A:252:LYS:HA	1.80	0.42
1:A:215[B]:MET:HE2	7:A:511:HOH:O	2.19	0.42
1:B:207:ILE:HD11	1:B:257:LYS:CE	2.50	0.42
1:B:305:TYR:HE1	7:B:504:HOH:O	2.03	0.41
1:A:336:HIS:HA	7:A:525:HOH:O	2.20	0.41
1:B:165:PRO:HB2	1:B:166:PRO:HD3	2.02	0.41
1:B:195:LYS:HD3	1:B:195:LYS:HA	1.94	0.41
1:A:252:LYS:HE2	4:A:404:GOL:O2	2.22	0.40
1:B:330:PHE:CD1	1:B:337:LEU:HD23	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:557:HOH:O	7:B:694:HOH:O[2_555]	2.02	0.18
7:B:551:HOH:O	7:B:689:HOH:O[2_445]	2.13	0.07

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/256 (101%)	251 (97%)	7 (3%)	0	100	100
1	B	256/256 (100%)	252 (98%)	4 (2%)	0	100	100
All	All	514/512 (100%)	503 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/223 (100%)	220 (99%)	3 (1%)	73	60
1	B	219/223 (98%)	217 (99%)	2 (1%)	82	74
All	All	442/446 (99%)	437 (99%)	5 (1%)	80	68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	276	ARG
1	A	283[A]	CYS
1	A	283[B]	CYS
1	B	224	SER
1	B	310	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	336	HIS
1	B	340	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	6FV	A	401	-	24,28,28	0.83	2 (8%)	28,41,41	3.06	5 (17%)
3	TLA	A	402	-	3,9,9	0.36	0	6,12,12	1.81	2 (33%)
3	TLA	A	403	6	3,9,9	1.23	0	6,12,12	1.65	1 (16%)
4	GOL	A	404	-	5,5,5	0.78	0	5,5,5	2.36	3 (60%)
5	EDO	A	405	-	3,3,3	0.39	0	2,2,2	0.49	0
2	6FV	B	401	-	24,28,28	0.86	1 (4%)	28,41,41	2.56	7 (25%)
3	TLA	B	402	-	3,9,9	0.90	0	6,12,12	1.63	2 (33%)
3	TLA	B	403	6	3,9,9	1.51	1 (33%)	6,12,12	2.13	2 (33%)
3	TLA	B	404	-	3,9,9	0.70	0	6,12,12	1.36	1 (16%)
3	TLA	B	405[A]	-	3,9,9	0.98	0	6,12,12	2.34	3 (50%)
3	TLA	B	405[B]	-	3,9,9	1.16	0	6,12,12	1.66	1 (16%)
4	GOL	B	406	-	5,5,5	0.78	0	5,5,5	0.68	0
5	EDO	B	407	-	3,3,3	1.05	0	2,2,2	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6FV	A	401	-	-	0/6/6/6	0/4/4/4
3	TLA	A	402	-	-	0/4/12/12	0/0/0/0
3	TLA	A	403	6	-	0/4/12/12	0/0/0/0
4	GOL	A	404	-	-	0/4/4/4	0/0/0/0
5	EDO	A	405	-	-	0/1/1/1	0/0/0/0
2	6FV	B	401	-	-	0/6/6/6	0/4/4/4
3	TLA	B	402	-	-	0/4/12/12	0/0/0/0
3	TLA	B	403	6	-	0/4/12/12	0/0/0/0
3	TLA	B	404	-	-	0/4/12/12	0/0/0/0
3	TLA	B	405[A]	-	-	0/4/12/12	0/0/0/0
3	TLA	B	405[B]	-	-	0/4/12/12	0/0/0/0
4	GOL	B	406	-	-	0/4/4/4	0/0/0/0
5	EDO	B	407	-	-	0/1/1/1	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	TLA	O3-C3	2.21	1.47	1.42
2	A	401	6FV	C4-C3	2.52	1.43	1.40
2	A	401	6FV	C1-N1	2.66	1.37	1.33
2	B	401	6FV	C1-N1	2.96	1.38	1.33

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	6FV	C4-C1-N1	-11.67	116.22	124.45
2	B	401	6FV	C4-C1-N1	-9.73	117.58	124.45
3	B	405[A]	TLA	C4-C3-C2	-4.19	104.09	113.11
3	B	405[B]	TLA	C4-C3-C2	-3.85	104.81	113.11
3	B	403	TLA	C4-C3-C2	-3.64	105.28	113.11
2	B	401	6FV	C10-C12-N4	-3.33	169.27	177.94
3	A	402	TLA	C1-C2-C3	-3.18	106.27	113.11
3	B	405[A]	TLA	C1-C2-C3	-3.09	106.45	113.11
3	B	404	TLA	C4-C3-C2	-3.08	106.48	113.11
4	A	404	GOL	C3-C2-C1	-3.08	99.29	111.52
3	B	402	TLA	C4-C3-C2	-2.93	106.79	113.11
3	B	402	TLA	C1-C2-C3	-2.59	107.54	113.11
3	A	403	TLA	C4-C3-C2	-2.37	108.00	113.11
3	A	402	TLA	O3-C3-C4	-2.35	105.42	111.13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	6FV	C4-C5-C7	-2.10	119.48	122.30
2	B	401	6FV	C1-C4-C3	-2.09	117.75	120.01
2	B	401	6FV	C4-C5-C7	-2.00	119.61	122.30
3	B	405[A]	TLA	O3-C3-C4	2.00	116.01	111.13
4	A	404	GOL	O1-C1-C2	2.23	121.31	110.07
2	B	401	6FV	C7-C6-N3	2.23	120.75	118.53
3	B	403	TLA	O2-C2-C3	2.32	115.88	108.88
2	A	401	6FV	C7-C6-N3	2.78	121.29	118.53
2	B	401	6FV	C5-C4-C3	3.38	121.01	117.01
4	A	404	GOL	O2-C2-C1	3.44	125.08	108.84
2	A	401	6FV	C5-C4-C3	3.79	121.50	117.01
2	B	401	6FV	C1-N1-C2	6.81	121.11	115.16
2	A	401	6FV	C1-N1-C2	9.29	123.28	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	6FV	1	0
3	A	403	TLA	1	0
4	A	404	GOL	9	0
5	A	405	EDO	1	0
2	B	401	6FV	1	0
3	B	405[A]	TLA	1	0
4	B	406	GOL	1	0
5	B	407	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/256 (96%)	0.15	13 (5%) 28 31	15, 23, 40, 73	0
1	B	249/256 (97%)	0.26	17 (6%) 18 21	18, 25, 49, 73	0
All	All	497/512 (97%)	0.20	30 (6%) 23 25	15, 24, 46, 73	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	370	LEU	9.8
1	A	370	LEU	5.9
1	B	369	VAL	5.8
1	B	335	GLY	4.8
1	A	336	HIS	4.2
1	B	227	GLY	4.0
1	A	368	VAL	4.0
1	A	334	GLU	3.6
1	B	264	SER	3.5
1	B	336	HIS	3.5
1	A	332	ALA	3.4
1	A	335	GLY	3.4
1	B	320	ALA	3.3
1	A	337	LEU	3.0
1	B	334	GLU	3.0
1	B	263	ASP	3.0
1	A	263	ASP	2.8
1	A	264	SER	2.7
1	B	333	GLU	2.6
1	B	226	GLY	2.4
1	B	170	TYR	2.4
1	B	368	VAL	2.3
1	A	227	GLY	2.2
1	A	352	CYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	278	ARG	2.2
1	B	351	ASP	2.1
1	B	262	PRO	2.1
1	B	337	LEU	2.1
1	A	278	ARG	2.0
1	A	170	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	A	404	6/6	0.74	0.31	10.49	50,53,55,57	0
5	EDO	A	405	4/4	0.80	0.33	7.93	54,55,59,61	0
4	GOL	B	406	6/6	0.93	0.12	3.37	25,30,34,34	0
5	EDO	B	407	4/4	0.90	0.14	3.24	36,40,41,41	0
3	TLA	B	404	10/10	0.82	0.18	1.31	35,43,46,48	0
2	6FV	A	401	25/25	0.94	0.13	0.78	20,26,31,43	0
3	TLA	B	403	10/10	0.94	0.10	0.36	20,23,25,25	0
3	TLA	B	402	10/10	0.93	0.11	0.17	23,27,31,32	0
3	TLA	A	403	10/10	0.95	0.09	-0.18	17,22,25,26	0
3	TLA	A	402	10/10	0.92	0.12	-0.21	24,26,28,29	0
2	6FV	B	401	25/25	0.92	0.10	-0.45	20,25,31,33	0
6	MN	B	408	1/1	0.99	0.05	-1.89	20,20,20,20	0
6	MN	A	406	1/1	1.00	0.04	-2.41	19,19,19,19	0
3	TLA	B	405[B]	10/10	0.80	0.31	-	38,41,41,42	10
3	TLA	B	405[A]	10/10	0.80	0.31	-	37,41,43,43	10

6.5 Other polymers [i](#)

There are no such residues in this entry.