



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:37 pm GMT

PDB ID : 4J7Q  
Title : Crystal structure of *Saccharomyces cerevisiae* Sfh3 complexed with phosphatidylinositol  
Authors : Yang, H.; Im, Y.J.  
Deposited on : 2013-02-14  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	
1	B	333	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	B7N	A	401	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol transfer protein PDR16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2652	1719	449	478	6			
1	B	322	Total	C	N	O	S	0	0	0
			2647	1715	448	478	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	GLY	-	EXPRESSION TAG	UNP P53860
A	14	SER	-	EXPRESSION TAG	UNP P53860
B	13	GLY	-	EXPRESSION TAG	UNP P53860
B	14	SER	-	EXPRESSION TAG	UNP P53860

- Molecule 2 is (1R)-2-{[(S)-HYDROXY{[(1S,2R,3R,4S,5S,6R)-2,3,4,5,6-PENTAHYDROXY CYCLOHEXYL]OXY}PHOSPHORYL]OXY}-1-[(OCTADECANOYLOXY)METHYL]ETHYL (9Z)-OCTADEC-9-ENOATE (three-letter code: B7N) (formula: C<sub>45</sub>H<sub>85</sub>O<sub>13</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			59	45	13	1		
2	B	1	Total	C	O	P	0	0
			59	45	13	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	141	Total	O	0	0
			141	141		
3	B	178	Total	O	0	0
			178	178		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phosphatidylinositol transfer protein PDR16



#### • Molecule 1: Phosphatidylinositol transfer protein PDR16



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.89Å 144.15Å 55.66Å 90.00° 111.28° 90.00°	Depositor
Resolution (Å)	34.40 – 1.55 34.40 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.9 (34.40-1.55) 98.9 (34.40-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.90 (at 1.55Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.221 , 0.239 0.219 , 0.238	Depositor DCC
$R_{free}$ test set	5571 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.6	Xtriage
Anisotropy	0.348	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5736	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B7N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/2720	0.55	0/3680
1	B	0.29	0/2715	0.56	0/3674
All	All	0.29	0/5435	0.56	0/7354

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2652	0	2678	75	0
1	B	2647	0	2668	62	0
2	A	59	0	84	15	0
2	B	59	0	84	9	0
3	A	141	0	0	1	0
3	B	178	0	0	7	0
All	All	5736	0	5514	126	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (126) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:GLN:HB3	2:A:401:B7N:H2A	1.44	0.95
1:B:157:ILE:HD13	1:B:196:ALA:HB3	1.53	0.89
1:B:210:VAL:H	1:B:211:PRO:HA	1.40	0.87
1:A:120:ILE:HG23	1:A:131:ILE:HD12	1.57	0.86
1:A:165:GLN:HE22	2:A:401:B7N:H1A	1.43	0.83
1:B:329:SER:O	1:B:333:LEU:HD13	1.79	0.82
1:B:140:ASN:HD22	1:B:176:HIS:HE1	1.30	0.80
1:B:140:ASN:HD22	1:B:176:HIS:CE1	1.99	0.79
1:A:165:GLN:NE2	2:A:401:B7N:H1A	1.98	0.77
1:A:165:GLN:HE22	2:A:401:B7N:C1	1.96	0.76
1:A:140:ASN:HD22	1:A:176:HIS:CE1	2.05	0.74
1:B:210:VAL:HB	1:B:212:GLY:N	2.03	0.73
1:A:222:GLY:HA2	1:B:247:TRP:CZ3	2.23	0.73
1:A:120:ILE:CG2	1:A:131:ILE:HD12	2.22	0.69
1:A:255:LEU:HD13	2:B:401:B7N:H39A	1.75	0.68
1:A:118:PHE:CD2	1:A:131:ILE:HD11	2.29	0.67
1:A:256:ILE:HD12	1:A:257:HIS:N	2.09	0.67
1:B:62:VAL:HG13	1:B:109:THR:HG23	1.76	0.67
1:A:209:LYS:HB2	1:A:209:LYS:NZ	2.10	0.67
1:B:192:GLN:HE21	1:B:193:ASP:H	1.42	0.66
1:B:328:LEU:HB3	1:B:333:LEU:HD11	1.78	0.65
1:B:157:ILE:HG23	1:B:198:LEU:HD13	1.77	0.65
1:B:248:LEU:HB3	3:B:669:HOH:O	1.96	0.65
1:A:140:ASN:HD22	1:A:176:HIS:HE1	1.43	0.64
1:A:165:GLN:NE2	2:A:401:B7N:C1	2.59	0.64
1:B:209:LYS:HB2	1:B:209:LYS:NZ	2.13	0.63
1:B:246:PRO:HB2	3:B:669:HOH:O	1.99	0.62
1:A:52:LEU:HD23	1:A:345:LYS:HE2	1.81	0.61
1:B:63:TYR:CZ	1:B:69:LYS:HE3	2.36	0.61
1:A:259:PHE:HB3	1:B:261:ASP:OD2	2.02	0.60
1:A:192:GLN:HE21	1:A:193:ASP:H	1.48	0.59
1:B:210:VAL:N	1:B:211:PRO:HA	2.14	0.59
1:B:280:LYS:HE2	1:B:290:ASP:OD1	2.02	0.59
1:A:301:TRP:HB3	1:A:302:PRO:HD3	1.85	0.59
1:A:176:HIS:HD2	3:A:557:HOH:O	1.86	0.56
1:B:54:HIS:HE1	3:B:527:HOH:O	1.87	0.56
1:A:222:GLY:HA2	1:B:247:TRP:HZ3	1.67	0.56
1:B:229:LEU:HD13	2:B:401:B7N:H40A	1.88	0.56
1:B:237:LEU:HB2	3:B:657:HOH:O	2.06	0.55
1:B:301:TRP:HB3	1:B:302:PRO:HD3	1.89	0.55
1:A:247:TRP:HE1	1:B:222:GLY:H	1.56	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ILE:HG23	1:A:198:LEU:HD13	1.90	0.54
1:B:160:LEU:HD13	2:B:401:B7N:H10	1.90	0.54
2:A:401:B7N:H5A	1:B:247:TRP:HH2	1.74	0.53
1:A:171:HIS:HE1	1:A:224:GLU:OE2	1.92	0.53
1:A:41:THR:H	1:A:44:GLN:NE2	2.06	0.53
1:B:334:ARG:NH1	3:B:601:HOH:O	2.42	0.52
2:A:401:B7N:H5A	1:B:247:TRP:CH2	2.45	0.52
1:B:266:GLU:HG2	3:B:677:HOH:O	2.10	0.52
1:A:138:VAL:O	1:A:141:GLU:HG3	2.11	0.51
1:A:111:THR:OG1	1:A:232:HIS:HD2	1.94	0.51
1:A:280:LYS:HE2	1:A:290:ASP:OD1	2.09	0.51
1:A:181:LEU:HD22	1:A:233:TYR:CE1	2.46	0.51
1:B:41:THR:H	1:B:44:GLN:NE2	2.09	0.50
2:B:401:B7N:H24A	3:B:657:HOH:O	2.11	0.50
1:B:192:GLN:HE21	1:B:193:ASP:N	2.10	0.50
1:B:173:GLN:C	2:B:401:B7N:H1	2.32	0.49
1:A:138:VAL:HA	1:A:141:GLU:HG3	1.94	0.49
1:B:111:THR:OG1	1:B:232:HIS:HD2	1.93	0.49
1:B:160:LEU:CD1	2:B:401:B7N:H10	2.43	0.49
1:B:221:VAL:O	1:B:221:VAL:HG22	2.12	0.49
1:A:197:LEU:HD11	2:A:401:B7N:H14A	1.95	0.49
1:B:140:ASN:ND2	1:B:176:HIS:HE1	2.04	0.49
1:B:209:LYS:HB2	1:B:209:LYS:HZ3	1.77	0.48
1:A:140:ASN:ND2	1:A:176:HIS:HE1	2.11	0.48
1:B:174:VAL:N	2:B:401:B7N:H1	2.28	0.48
1:A:62:VAL:HG13	1:A:109:THR:HG23	1.96	0.48
1:A:221:VAL:HG13	1:A:222:GLY:N	2.29	0.48
1:A:169:THR:HA	1:A:173:GLN:OE1	2.14	0.47
1:A:16:LYS:C	1:A:18:LEU:H	2.17	0.47
1:A:222:GLY:CA	1:B:247:TRP:CZ3	2.97	0.47
1:A:255:LEU:HD22	2:B:401:B7N:O7	2.15	0.46
1:A:247:TRP:CD1	1:B:220:GLY:HA3	2.49	0.46
1:A:192:GLN:HE21	1:A:193:ASP:N	2.13	0.46
1:B:262:PRO:HG3	1:B:265:ARG:HH21	1.81	0.46
1:A:256:ILE:HD12	1:A:256:ILE:C	2.37	0.45
1:A:229:LEU:HD22	2:A:401:B7N:H23	1.98	0.45
1:A:209:LYS:HB2	1:A:209:LYS:HZ3	1.81	0.45
1:A:306:GLU:O	1:A:310:GLU:HG3	2.17	0.45
1:A:225:VAL:HG21	2:A:401:B7N:H5	1.96	0.45
1:A:41:THR:H	1:A:44:GLN:HE21	1.65	0.45
1:B:125:GLU:HA	1:B:129:ASP:HB2	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:LEU:HD12	1:A:256:ILE:HD11	1.98	0.45
1:B:171:HIS:HE1	1:B:224:GLU:OE2	1.99	0.45
1:A:241:LEU:HB3	1:A:274:PHE:CZ	2.52	0.44
1:A:247:TRP:NE1	1:B:220:GLY:CA	2.80	0.44
1:B:210:VAL:HB	1:B:211:PRO:C	2.37	0.44
1:A:181:LEU:HD23	1:A:181:LEU:O	2.17	0.44
1:A:139:GLU:O	1:A:144:LYS:HE2	2.17	0.44
1:A:318:ARG:HD2	1:A:335:GLY:O	2.18	0.44
1:A:247:TRP:CD1	1:B:220:GLY:CA	3.01	0.44
1:B:62:VAL:HG12	1:B:63:TYR:N	2.32	0.44
1:A:181:LEU:O	1:A:184:VAL:HG12	2.17	0.44
1:A:247:TRP:CZ2	1:B:222:GLY:CA	3.00	0.44
2:A:401:B7N:H39A	1:B:255:LEU:HD13	2.00	0.43
1:A:247:TRP:CZ2	1:B:222:GLY:N	2.86	0.43
1:A:181:LEU:HD11	2:A:401:B7N:H25	2.00	0.43
1:A:192:GLN:HE21	1:A:194:SER:H	1.66	0.43
1:A:249:ALA:HB1	2:A:401:B7N:H29	2.00	0.43
1:B:337:HIS:HD2	1:B:339:LYS:O	2.02	0.43
1:A:118:PHE:CE2	1:A:131:ILE:HD11	2.54	0.43
1:A:247:TRP:HE1	1:B:220:GLY:C	2.22	0.43
1:A:209:LYS:HB2	1:A:209:LYS:HZ2	1.83	0.43
1:A:262:PRO:O	1:A:266:GLU:HG2	2.17	0.43
1:A:198:LEU:HD11	1:A:241:LEU:HD22	2.01	0.43
1:B:276:LYS:HD3	1:B:276:LYS:O	2.19	0.43
1:A:256:ILE:HD13	1:A:260:ILE:HD11	2.01	0.42
1:B:179:PHE:O	1:B:183:ARG:HG2	2.19	0.42
1:A:247:TRP:HZ2	1:B:222:GLY:N	2.18	0.42
1:B:241:LEU:HB3	1:B:274:PHE:CZ	2.54	0.42
1:A:229:LEU:HD13	2:A:401:B7N:H13	2.02	0.42
1:B:111:THR:HA	1:B:232:HIS:CD2	2.54	0.42
1:A:62:VAL:HG12	1:A:63:TYR:N	2.35	0.41
2:B:401:B7N:H40	2:B:401:B7N:H14A	2.01	0.41
1:A:195:LEU:HD12	1:A:195:LEU:C	2.40	0.41
1:A:179:PHE:O	1:A:183:ARG:HG2	2.21	0.41
1:A:165:GLN:HE22	2:A:401:B7N:H1	1.80	0.41
1:A:197:LEU:C	1:A:198:LEU:HD12	2.41	0.41
1:A:247:TRP:CZ2	1:B:222:GLY:HA2	2.56	0.41
1:B:41:THR:H	1:B:44:GLN:HE21	1.68	0.41
1:A:118:PHE:CD2	1:A:131:ILE:CD1	3.00	0.41
1:A:111:THR:HA	1:A:232:HIS:CD2	2.56	0.41
1:A:52:LEU:CD2	1:A:345:LYS:HE2	2.50	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:LEU:HD12	1:B:195:LEU:C	2.42	0.40
1:B:241:LEU:N	1:B:241:LEU:HD12	2.37	0.40
1:B:263:LEU:HD22	1:B:263:LEU:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/333 (96%)	315 (99%)	1 (0%)	2 (1%)	28	6
1	B	318/333 (96%)	313 (98%)	3 (1%)	2 (1%)	28	6
All	All	636/666 (96%)	628 (99%)	4 (1%)	4 (1%)	28	6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	209	LYS
1	A	206	ASP
1	B	210	VAL
1	A	17	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/300 (97%)	288 (99%)	3 (1%)	80	61
1	B	290/300 (97%)	286 (99%)	4 (1%)	71	45
All	All	581/600 (97%)	574 (99%)	7 (1%)	75	53

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	151	GLU
1	A	209	LYS
1	B	140	ASN
1	B	151	GLU
1	B	181	LEU
1	B	247	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	68	ASN
1	A	140	ASN
1	A	171	HIS
1	A	176	HIS
1	A	192	GLN
1	A	230	GLN
1	A	232	HIS
1	A	244	ASN
1	B	44	GLN
1	B	54	HIS
1	B	68	ASN
1	B	140	ASN
1	B	171	HIS
1	B	176	HIS
1	B	192	GLN
1	B	232	HIS
1	B	244	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	B7N	A	401	-	59,59,59	0.82	1 (1%)	67,71,71	1.03	5 (7%)
2	B7N	B	401	-	59,59,59	0.74	0	67,71,71	1.06	4 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B7N	A	401	-	-	0/54/78/78	0/1/1/1
2	B7N	B	401	-	-	0/54/78/78	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	B7N	C11-C10	-2.26	1.38	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	B7N	C24-C25-C27	-2.83	103.24	113.58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	B7N	C24-C25-C27	-2.49	104.51	113.58
2	B	401	B7N	C12-C13-C14	-2.47	101.72	114.45
2	A	401	B7N	C6-C7-C8	-2.46	104.29	113.74
2	B	401	B7N	C15-O8-C27	-2.34	112.36	117.88
2	A	401	B7N	C6-C5-C4	-2.16	103.31	114.45
2	A	401	B7N	O9-C42-C40	2.49	119.15	111.90
2	A	401	B7N	O8-C27-C25	3.45	118.72	111.55
2	B	401	B7N	O8-C27-C25	3.90	119.65	111.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	B7N	15	0
2	B	401	B7N	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/333 (96%)	0.89	41 (12%) 4 4	14, 21, 38, 48	0
1	B	322/333 (96%)	0.64	29 (9%) 10 12	11, 18, 32, 48	0
All	All	644/666 (96%)	0.76	70 (10%) 6 7	11, 20, 37, 48	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	210	VAL	13.6
1	A	247	TRP	12.2
1	B	210	VAL	10.5
1	A	16	LYS	9.5
1	A	211	PRO	9.5
1	B	17	ASN	9.4
1	A	205	PRO	9.1
1	B	211	PRO	8.9
1	B	16	LYS	8.4
1	B	247	TRP	8.1
1	A	209	LYS	7.9
1	A	17	ASN	7.0
1	B	220	GLY	6.2
1	A	220	GLY	6.2
1	B	221	VAL	5.9
1	B	212	GLY	5.8
1	A	206	ASP	5.8
1	A	221	VAL	5.8
1	B	209	LYS	5.7
1	B	18	LEU	5.4
1	B	38	LYS	5.3
1	A	207	VAL	5.1
1	A	345	LYS	4.6
1	A	36	LYS	4.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	18	LEU	4.2
1	A	168	LYS	4.2
1	B	36	LYS	3.5
1	A	263	LEU	3.4
1	B	168	LYS	3.4
1	A	256	ILE	3.4
1	A	169	THR	3.2
1	A	181	LEU	3.1
1	B	252	PHE	3.1
1	A	38	LYS	3.0
1	A	246	PRO	3.0
1	A	261	ASP	3.0
1	A	266	GLU	3.0
1	A	208	PRO	3.0
1	B	181	LEU	2.9
1	A	262	PRO	2.9
1	B	185	ILE	2.8
1	A	37	GLU	2.7
1	B	158	LEU	2.7
1	A	338	GLU	2.7
1	A	43	GLU	2.7
1	A	198	LEU	2.7
1	A	185	ILE	2.6
1	A	184	VAL	2.6
1	A	19	ILE	2.6
1	A	158	LEU	2.6
1	B	39	PRO	2.5
1	A	339	LYS	2.5
1	B	286	LEU	2.5
1	B	169	THR	2.5
1	B	37	GLU	2.4
1	B	157	ILE	2.4
1	B	266	GLU	2.4
1	A	286	LEU	2.4
1	B	263	LEU	2.4
1	A	204	TYR	2.3
1	A	203	ASP	2.3
1	A	248	LEU	2.2
1	B	280	LYS	2.2
1	A	249	ALA	2.1
1	A	40	LEU	2.1
1	A	197	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	198	LEU	2.1
1	B	338	GLU	2.1
1	B	147	ILE	2.0
1	B	261	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	B7N	A	401	59/59	0.83	0.25	2.74	22,44,47,51	0
2	B7N	B	401	59/59	0.86	0.21	1.16	18,40,45,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.