



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2018 – 10:41 PM EST

PDB ID : 2JAI
Title : DDAH1 complexed with citrulline
Authors : Murray-Rust, J.; O'Hara, B.P.; Rossiter, S.; Leiper, J.M.; Vallance, P.; McDonald, N.Q.
Deposited on : 2006-11-29
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>83%</div> <div>10% 6%</div> </div>
1	B	289	<div> <div>21%</div> <div>81% 12% 6%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NG, NG-DIMETHYLARGININE DIMETHYLAMINOHYDROLYASE 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	Se	0	3	0
			2042	1288	347	391	7	9			
1	B	271	Total	C	N	O	S	Se	0	0	0
			1877	1192	326	345	6	8			

- Molecule 2 is CITRULLINE (three-letter code: CIR) (formula: $C_6H_{13}N_3O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	114	Total	O	0	0
			114	114		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	11	Total	O	0	0
			11	11		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.83 Å 46.80 Å 147.79 Å 90.00° 94.64° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 44.68 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.8 (50.00-2.30) 92.8 (44.68-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.80 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.198 , 0.258 0.201 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.442	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4056	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/2071	0.80	1/2794 (0.0%)
1	B	0.89	11/1895 (0.6%)	0.72	1/2570 (0.0%)
All	All	0.84	11/3966 (0.3%)	0.76	2/5364 (0.0%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	146	ASN	CG-OD1	9.77	1.45	1.24
1	B	146	ASN	CG-ND2	9.76	1.57	1.32
1	B	122	ASN	CG-ND2	8.94	1.55	1.32
1	B	122	ASN	CG-OD1	8.66	1.43	1.24
1	B	122	ASN	C-N	6.97	1.50	1.34
1	B	207	ARG	N-CA	6.21	1.58	1.46
1	B	158	LYS	C-O	6.03	1.34	1.23
1	B	176	PHE	CG-CD1	5.53	1.47	1.38
1	B	122	ASN	N-CA	5.23	1.56	1.46
1	B	207	ARG	CA-CB	5.04	1.65	1.53
1	B	149	GLY	C-O	5.03	1.31	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	146	ASN	CB-CG-OD1	-5.10	111.40	121.60
1	A	165	VAL	CB-CA-C	-5.04	101.82	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2042	0	2007	20	0
1	B	1877	0	1750	16	0
2	A	12	0	12	2	0
3	A	114	0	0	1	0
3	B	11	0	0	0	0
All	All	4056	0	3769	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ASN:ND2	1:A:206[A]:HIS:NE2	2.33	0.75
1:A:270:LEU:O	1:A:273[B]:CYS:SG	2.47	0.73
1:B:178:SER:HB2	1:B:219:ALA:O	1.88	0.71
1:B:163:SER:HB3	1:B:203:MSE:SE	2.41	0.70
1:A:236:ARG:NH2	1:A:265:GLU:OE1	2.29	0.63
1:A:203:MSE:SE	1:B:-4:GLY:HA2	2.50	0.60
1:A:154:ALA:HA	1:A:162:VAL:HG21	1.84	0.58
1:A:98:ARG:O	1:A:101:VAL:HB	2.06	0.56
1:A:206[B]:HIS:HE1	3:A:2083:HOH:O	1.89	0.55
1:A:273[B]:CYS:HB3	2:A:1281:CIR:C7	2.38	0.53
1:A:43:GLU:OE2	1:A:47:GLN:NE2	2.33	0.51
1:A:197:LEU:HA	1:A:200:MSE:HE3	1.92	0.51
1:A:221:CYS:SG	1:A:233:LEU:HD11	2.51	0.50
1:A:270:LEU:N	1:A:273[B]:CYS:SG	2.85	0.50
1:A:183:ASN:HD21	1:A:206[A]:HIS:CD2	2.28	0.50
1:B:75:PHE:CE2	1:B:269:GLY:HA3	2.48	0.49
1:B:64:LEU:HD13	1:B:103:MSE:HG2	1.95	0.48
1:A:227:PRO:O	1:A:228:ASN:HB2	2.13	0.48
1:B:221:CYS:HB2	1:B:234:LEU:O	2.14	0.48
1:B:76:VAL:HB	1:B:104:MSE:CE	2.44	0.47
1:A:129:ASP:OD1	1:A:172:HIS:HA	2.16	0.46
1:B:197:LEU:HA	1:B:200:MSE:HE3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:VAL:O	1:B:164:THR:HA	2.17	0.44
1:B:89:ILE:CG2	1:B:101:VAL:HG13	2.48	0.43
1:B:129:ASP:OD1	1:B:172:HIS:HA	2.18	0.43
1:B:92:PRO:HD2	1:B:98:ARG:HG2	2.01	0.43
1:A:237:THR:OG1	1:A:240:GLU:HG3	2.19	0.43
1:B:45:GLN:HE22	1:B:262:SER:HB2	1.83	0.43
1:A:273[B]:CYS:HB3	2:A:1281:CIR:O7	2.19	0.42
1:A:226:ILE:HB	1:A:229:LYS:HB2	2.02	0.41
1:A:105:LYS:O	1:A:109:GLU:HG3	2.20	0.41
1:B:18:ARG:HD3	1:B:73:CYS:HB2	2.03	0.41
1:B:51:GLY:O	1:B:55:SER:HB3	2.20	0.41
1:B:79:VAL:HG23	1:B:104:MSE:HG2	2.03	0.40
1:A:77:GLU:HB3	1:A:273[B]:CYS:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/289 (93%)	260 (96%)	10 (4%)	0	100	100
1	B	257/289 (89%)	248 (96%)	9 (4%)	0	100	100
All	All	527/578 (91%)	508 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/232 (93%)	214 (99%)	2 (1%)	82	91
1	B	173/232 (75%)	169 (98%)	4 (2%)	56	73
All	All	389/464 (84%)	383 (98%)	6 (2%)	68	83

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	VAL
1	A	260	SER
1	B	44	ARG
1	B	103	MSE
1	B	178	SER
1	B	262	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	183	ASN
1	B	235	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CIR	A	1281	1	6,11,11	0.77	0	6,13,13	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIR	A	1281	1	-	0/7/11/11	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1281	CIR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/289 (91%)	0.63	3 (1%) 80 84	31, 35, 42, 52	0
1	B	262/289 (90%)	1.33	61 (23%) 1 1	28, 36, 44, 55	0
All	All	525/578 (90%)	0.98	64 (12%) 5 7	28, 36, 42, 55	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	SER	5.4
1	B	115	ILE	4.9
1	B	190	SER	4.8
1	B	161	ALA	4.7
1	B	140	GLY	4.5
1	B	3	LEU	4.3
1	B	165	VAL	4.3
1	B	154	ALA	4.3
1	B	197	LEU	3.8
1	B	213	VAL	3.8
1	B	123	ALA	3.8
1	B	186	ALA	3.8
1	B	193	ALA	3.7
1	B	156	THR	3.7
1	B	1	ALA	3.7
1	B	150	ALA	3.3
1	B	149	GLY	3.2
1	B	67	ASP	3.2
1	B	167	VAL	3.2
1	B	70	LEU	3.1
1	B	141	LEU	3.0
1	B	138	PHE	3.0
1	B	114	ASN	3.0
1	B	215	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	146	ASN	3.0
1	B	184	LEU	2.9
1	B	191	GLU	2.9
1	B	153	LEU	2.9
1	B	55	SER	2.9
1	B	187	ILE	2.8
1	B	199	ILE	2.8
1	B	259	VAL	2.8
1	B	134	GLY	2.7
1	B	248	TYR	2.7
1	B	194	GLN	2.6
1	B	4	GLY	2.6
1	B	235	HIS	2.6
1	B	247	VAL	2.5
1	B	230	GLY	2.5
1	B	145	THR	2.5
1	B	198	LYS	2.5
1	B	192	SER	2.4
1	B	2	GLY	2.4
1	B	79	VAL	2.4
1	A	206[A]	HIS	2.4
1	B	185	ILE	2.4
1	B	196	ALA	2.3
1	B	211	LEU	2.3
1	B	152	ILE	2.3
1	B	93	GLY	2.2
1	B	177	CYS	2.2
1	B	122	ASN	2.2
1	B	151	GLU	2.2
1	B	214	PRO	2.2
1	B	-1	GLY	2.2
1	B	86	THR	2.2
1	B	250	LYS	2.1
1	B	109	GLU	2.1
1	A	273[A]	CYS	2.1
1	B	157	PHE	2.1
1	B	42	ALA	2.0
1	A	228	ASN	2.0
1	B	209	ASP	2.0
1	B	260	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CIR	A	1281	12/12	0.93	0.21	1.18	11,13,17,17	12

6.5 Other polymers [i](#)

There are no such residues in this entry.