



wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Mar 2, 2017 – 11:43 am GMT

PDB ID : 3JB1
EMDB ID: : EMD-6375
Title : Atomic model of cytoplasmic polyhedrosis virus with SAM
Authors : Yu, X.K.; Jiang, J.S.; Sun, J.C.; Zhou, Z.H.
Deposited on : 2015-07-06
Resolution : 3.10 Å(reported)

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report
for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : recalc29047


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	125131	1336
Ramachandran outliers	121729	1120
Sidechain outliers	121581	1026

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	1058	64% 31% 5%
2	B	1333	59% 28% • 11%
2	C	1333	64% 26% • 6%
3	D	448	43% 19% • 35%
3	E	448	47% 16% • 35%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 32271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Structural protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1057	Total	C	N	O	S	0	0
			8434	5345	1457	1587	45		


- Molecule 2 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1191	Total	C	N	O	S	0	0
			9397	5937	1634	1789	37		
2	C	1250	Total	C	N	O	S	0	0
			9851	6219	1712	1882	38		

- Molecule 3 is a protein called Viral structural protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	292	Total	C	N	O	S	0	0
			2281	1449	399	425	8		
3	E	292	Total	C	N	O	S	0	0
			2281	1449	399	425	8		

- Molecule 4 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).




Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	N	O	S	0
			27	15	6	5	1	

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Structural protein VP3



• Molecule 2: Capsid protein VP1









• Molecule 3: Viral structural protein 5

Chain D: 43% 19% 35%





4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	44908	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Not provided	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	Not provided	Depositor
Maximum defocus (nm)	Not provided	Depositor
Magnification	60535	Depositor
Image detector	KODAK SO-163 FILM	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 2$	RMSZ	$\# Z > 2$
1	A	0.28	0/8619	0.52	3/11737 (0.0%)
2	B	0.33	0/9590	0.54	0/13056
2	C	0.34	0/10052	0.56	0/13687
3	D	0.33	0/2327	0.54	0/3163
3	E	0.30	0/2327	0.52	0/3163
All	All	0.32	0/32915	0.54	3/44806 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	ALA	C-N-CD	-8.20	102.57	120.60
1	A	186	ALA	C-N-CA	5.69	145.91	122.00
1	A	890	LEU	CA-CB-CG	5.39	127.70	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8434	0	8399	219	0
2	B	9397	0	9315	255	0
2	C	9851	0	9762	224	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	2281	0	2282	62	0
3	E	2281	0	2282	47	0
4	A	27	0	22	5	0
All	All	32271	0	32062	785	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 785 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:709:MET:O	2:C:715:ASN:ND2	2.09	0.85
2:C:873:TYR:HB3	2:C:898:GLN:HB2	1.60	0.83
3:D:44:LEU:HG	3:D:174:VAL:HG22	1.64	0.79
1:A:752:VAL:HG12	1:A:781:VAL:HG23	1.65	0.78
2:C:462:LEU:HD13	2:C:680:THR:HG22	1.64	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1055/1058 (100%)	1004 (95%)	45 (4%)	6 (1%)	28	67
2	B	1187/1333 (89%)	1135 (96%)	49 (4%)	3 (0%)	44	79
2	C	1246/1333 (94%)	1181 (95%)	60 (5%)	5 (0%)	38	75
3	D	290/448 (65%)	281 (97%)	8 (3%)	1 (0%)	44	79
3	E	290/448 (65%)	281 (97%)	9 (3%)	0	100	100
All	All	4068/4620 (88%)	3882 (95%)	171 (4%)	15 (0%)	42	75

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ALA
2	C	1145	ALA
2	C	1251	VAL
2	B	1123	PRO
3	D	244	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	942/943 (100%)	829 (88%)	113 (12%)	6	24
2	B	1038/1153 (90%)	963 (93%)	75 (7%)	17	51
2	C	1089/1153 (94%)	980 (90%)	109 (10%)	9	33
3	D	240/379 (63%)	215 (90%)	25 (10%)	8	31
3	E	240/379 (63%)	222 (92%)	18 (8%)	16	49
All	All	3549/4007 (89%)	3209 (90%)	340 (10%)	14	36

5 of 340 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	797	SER
2	C	98	ASN
3	D	160	LEU
2	B	851	THR
2	B	1092	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1138	HIS
2	C	156	GLN
2	C	981	HIS
2	B	858	HIS
2	C	1138	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SAM	A	1101	-	21,29,29	1.15	2 (9%)	17,42,42	2.48	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAM	A	1101	-	-	0/8/33/33	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1101	SAM	C2-N1	2.60	1.38	1.33
4	A	1101	SAM	C2-N3	3.68	1.38	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	1101	SAM	N3-C2-N1	-9.59	120.51	128.86
4	A	1101	SAM	C4'-O4'-C1'	2.01	111.91	109.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1101	SAM	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.