



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:26 am GMT

PDB ID : 4JKF  
Title : Open and closed forms of T1791P+R1865A human PRP8 RNase H-like domain with bound Mg ion  
Authors : Schellenberg, M.J.; Wu, T.; Ritchie, D.B.; Atta, K.A.; MacMillan, A.M.  
Deposited on : 2013-03-09  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>7% ••</div> </div> </div>
1	B	222	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>12% • 5%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-processing-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	1	0
			1794	1177	298	316	3			
1	B	210	Total	C	N	O	S	0	3	0
			1725	1136	286	299	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1789	PRO	THR	ENGINEERED MUTATION	UNP Q6P2Q9
A	1865	ALA	ARG	ENGINEERED MUTATION	UNP Q6P2Q9
B	1789	PRO	THR	ENGINEERED MUTATION	UNP Q6P2Q9
B	1865	ALA	ARG	ENGINEERED MUTATION	UNP Q6P2Q9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	257	Total	O	0	2
			259	259		
3	B	177	Total	O	0	1
			178	178		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-processing-splicing factor 8



- Molecule 1: Pre-mRNA-processing-splicing factor 8



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.17Å 77.25Å 93.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.03 – 1.95 40.03 – 1.95	Depositor EDS
% Data completeness (in resolution range)	94.0 (40.03-1.95) 94.0 (40.03-1.95)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.186 , 0.226 0.192 , 0.233	Depositor DCC
$R_{free}$ test set	2016 reflections (5.55%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.8	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3957	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	1/1839 (0.1%)	0.71	0/2500
1	B	0.64	1/1771 (0.1%)	0.66	0/2408
All	All	0.65	2/3610 (0.1%)	0.69	0/4908

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1827	TRP	CD2-CE2	5.31	1.47	1.41
1	B	1839	TRP	CD2-CE2	5.29	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1794	0	1863	11	0
1	B	1725	0	1809	21	0
2	B	1	0	0	0	0
3	A	259	0	0	4	0
3	B	178	0	0	4	0
All	All	3957	0	3672	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1850[B]:ARG:NH1	3:B:2190[B]:HOH:O	1.77	1.15
1:B:1820:LYS:HG3	1:B:1914[A]:MET:HE3	1.73	0.69
1:A:1773:SER:HG	1:A:1775:GLN:N	1.92	0.68
1:B:1985:ASP:HB3	3:B:2207:HOH:O	1.96	0.65
1:B:1820:LYS:HG3	1:B:1914[A]:MET:CE	2.31	0.61
1:A:1813:ARG:HD3	3:A:2167:HOH:O	2.01	0.59
1:B:1789:PRO:O	1:B:1790:ILE:HG22	2.03	0.58
1:B:1957:ASP:OD1	1:B:1958:LYS:N	2.37	0.57
1:B:1977:ILE:O	1:B:1981:VAL:HG23	2.07	0.54
1:A:1813:ARG:CD	3:A:2167:HOH:O	2.55	0.53
1:B:1819:LEU:N	1:B:1914[A]:MET:HE2	2.24	0.52
3:A:2244:HOH:O	1:B:1914[B]:MET:HE2	2.09	0.52
1:B:1869:LEU:HD23	1:B:1884:ILE:HG22	1.92	0.51
1:B:1970:THR:O	1:B:1971:LEU:HD23	2.09	0.51
1:B:1824:THR:HB	1:B:1840:LYS:HE2	1.94	0.49
1:B:1789:PRO:O	1:B:1790:ILE:CB	2.60	0.48
1:A:1827:TRP:O	1:A:1828:ALA:C	2.48	0.48
1:B:1859:LYS:HB2	3:B:2202:HOH:O	2.15	0.47
1:A:1978:LYS:NZ	1:A:1978:LYS:HB3	2.30	0.46
1:A:1968:TRP:HB3	1:A:1969:PRO:HD2	1.98	0.45
1:B:1870:ASP:HB3	3:B:2228:HOH:O	2.16	0.44
1:B:1789:PRO:O	1:B:1790:ILE:HB	2.18	0.44
1:A:1831:LYS:O	3:A:2257:HOH:O	2.21	0.44
1:B:1818:PHE:HB3	1:B:1914[A]:MET:CE	2.49	0.43
1:B:1775:GLN:HE22	1:B:1860:GLN:HE21	1.66	0.42
1:A:1879:PHE:CB	1:A:1882:ILE:HD12	2.50	0.42
1:A:1855:GLU:OE1	1:B:1813:ARG:NH2	2.44	0.42
1:B:1973:ASP:O	1:B:1977:ILE:HG13	2.21	0.41
1:B:1789:PRO:O	1:B:1790:ILE:CG2	2.67	0.41
1:A:1961:ILE:H	1:A:1961:ILE:HD13	1.85	0.41
1:A:1920:TYR:O	1:A:1921:ASP:C	2.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/222 (98%)	211 (97%)	6 (3%)	0	100	100
1	B	209/222 (94%)	204 (98%)	5 (2%)	0	100	100
All	All	426/444 (96%)	415 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/200 (100%)	197 (98%)	3 (2%)	70	65
1	B	193/200 (96%)	187 (97%)	6 (3%)	45	32
All	All	393/400 (98%)	384 (98%)	9 (2%)	59	47

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1881	ASN
1	A	1907	LEU
1	A	1961	ILE
1	B	1800	THR
1	B	1870	ASP
1	B	1908	LYS
1	B	1914[A]	MET
1	B	1914[B]	MET

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1984	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1966	HIS
1	B	1775	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	220/222 (99%)	0.09	8 (3%) 43 54	16, 26, 51, 72	0
1	B	210/222 (94%)	0.28	14 (6%) 19 28	16, 31, 61, 73	0
All	All	430/444 (96%)	0.18	22 (5%) 29 39	16, 28, 58, 73	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1959	THR	6.3
1	A	1829	GLY	5.7
1	B	1774	ASN	4.5
1	B	1957	ASP	4.1
1	B	1790	ILE	3.9
1	A	1772	PHE	3.8
1	A	1830	GLN	3.4
1	B	1958	LYS	3.1
1	A	1828	ALA	3.1
1	B	1789	PRO	3.0
1	B	1961	ILE	2.9
1	B	1960	THR	2.9
1	B	1989	ALA	2.7
1	B	1956	PRO	2.7
1	A	1958	LYS	2.6
1	A	1773	SER	2.6
1	A	1770	GLU	2.5
1	B	1988	LEU	2.5
1	B	1771	LEU	2.4
1	A	1771	LEU	2.4
1	B	1952	VAL	2.4
1	B	1949	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	2001	1/1	0.99	0.07	-1.36	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.