



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 03:28 am GMT

PDB ID : 1JM6
Title : Pyruvate dehydrogenase kinase, isozyme 2, containing ADP
Authors : Steussy, C.N.; Popov, K.M.; Bowker-Kinley, M.M.; Sloan, R.B.; Harris, R.A.;
Hamilton, J.A.
Deposited on : 2001-07-17
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	<div> <div>2%</div> <div> <div></div> <div>34%</div> <div>42%</div> <div>6%</div> <div>17%</div> </div> </div>
1	B	407	<div> <div>2%</div> <div> <div></div> <div>37%</div> <div>39%</div> <div>6%</div> <div>17%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ADP	B	3510	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyruvate dehydrogenase kinase, isozyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	0	0
			2709	1743	444	505	17			
1	B	336	Total	C	N	O	S	0	0	0
			2684	1728	441	498	17			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	119	Total	O	0	0
			119	119		
4	B	109	Total	O	0	0
			109	109		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate dehydrogenase kinase, isozyme 2



• Molecule 1: Pyruvate dehydrogenase kinase, isozyme 2



K1206	E1293	V1365
Y1207	R1294	TYR
M1208	L1295	ASN
M1209	F1296	LYS
	S1297	SER
P1212	Y1298	ALA
		TRP
E1215	T1302	ARG
I1216	A1303	HIS
Q1217	P1304	TYR
E1218	T1305	GLN
V1219	P1306	THR
N1220	GLN	ILE
	PRO	GLN
N1223	GLY	GLU
	THR	ALA
I1228	GLY	ALA
	GLY	GLY
	ASP	GLY
Y1232	THR	TRP
V1233	P1314	CYS
P1234	L1315	VAL
S1235	A1316	PRO
H1236	G1317	SER
L1237	F1318	THR
Y1238	G1319	GLU
	Y1320	PRO
F1245	G1321	LYS
K1246	L1322	ASN
N1247	P1323	THR
A1248	I1324	SER
M1249	S1325	THR
R1250	R1326	TYR
	L1327	ARG
V1253		VAL
S1258	K1330	SER
	Y1331	
T1261	F1332	
L1262	Q1333	
P1263	G1334	
P1264	D1335	
I1265	L1338	
K1266	F1339	
I1267	S1340	
M1268	M1341	
V1269	E1342	
A1270	G1343	
L1271	F1344	
	G1345	
L1276	T1346	
	D1347	
K1279		
R1283	T1357	
	D1358	
V1287	S1359	
	V1360	
R1290	E1361	
K1291	R1362	
I1292	L1363	
	P1364	

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.42Å 109.87Å 71.41Å 90.00° 104.52° 90.00°	Depositor
Resolution (Å)	6.00 – 2.50 29.26 – 2.40	Depositor EDS
% Data completeness (in resolution range)	84.2 (6.00-2.50) 85.4 (29.26-2.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 2.39Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.202 , 0.262 0.188 , 0.241	Depositor DCC
R_{free} test set	1539 reflections (4.91%)	DCC
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.478 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5677	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	6/2771 (0.2%)	1.14	16/3752 (0.4%)
1	B	0.65	5/2745 (0.2%)	0.94	6/3715 (0.2%)
All	All	0.72	11/5516 (0.2%)	1.05	22/7467 (0.3%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1090	PHE	C-N	-8.56	1.14	1.34
1	A	1178	LYS	C-N	-7.66	1.16	1.34
1	B	1306	PRO	CA-C	7.37	1.67	1.52
1	A	1215	GLU	CB-CG	7.13	1.65	1.52
1	B	1215	GLU	CD-OE2	6.47	1.32	1.25

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1090	PHE	O-C-N	-12.48	102.73	122.70
1	A	1137	PRO	C-N-CA	10.68	148.39	121.70
1	A	1138	VAL	N-CA-C	10.08	138.22	111.00
1	A	1006	ALA	C-N-CA	-9.54	102.27	122.30
1	A	1090	PHE	CA-C-N	9.25	137.55	117.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2709	0	2692	254	0
1	B	2684	0	2679	192	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	27	0	12	1	0
3	B	27	0	12	9	0
4	A	119	0	0	17	0
4	B	109	0	0	16	0
All	All	5677	0	5395	442	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

The worst 5 of 442 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1042:PHE:CE2	1:A:1095:PRO:HG3	1.64	1.33
1:A:1138:VAL:CG1	1:A:1138:VAL:O	1.75	1.29
1:A:1223:ASN:HD22	1:A:1226:GLN:CB	1.58	1.16
1:A:1031:PHE:HA	1:A:1036:ALA:N	1.60	1.15
1:A:1177:PRO:HD2	1:A:1179:HIS:NE2	1.62	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/407 (81%)	290 (88%)	32 (10%)	7 (2%)	8	13
1	B	326/407 (80%)	306 (94%)	18 (6%)	2 (1%)	28	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	655/814 (80%)	596 (91%)	50 (8%)	9 (1%)	13	23

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1138	VAL
1	A	1224	ALA
1	A	1295	LEU
1	A	1039	LYS
1	B	1199	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/363 (84%)	277 (91%)	28 (9%)	11	20
1	B	302/363 (83%)	276 (91%)	26 (9%)	12	23
All	All	607/726 (84%)	553 (91%)	54 (9%)	11	22

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1353	LYS
1	B	1016	SER
1	B	1340	SER
1	A	1357	THR
1	A	1362	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1236	HIS
1	A	1239	HIS
1	B	1082	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1217	GLN
1	A	1223	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ADP	A	3500	2	25,29,29	1.32	3 (12%)	24,45,45	2.52	2 (8%)
3	ADP	B	3510	2	25,29,29	1.32	3 (12%)	24,45,45	2.52	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	3500	2	-	0/12/32/32	0/3/3/3
3	ADP	B	3510	2	-	0/12/32/32	0/3/3/3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3500	ADP	PB-O3A	-2.97	1.55	1.60
3	B	3510	ADP	PB-O3A	-2.93	1.55	1.60
3	A	3500	ADP	PB-O2B	2.50	1.65	1.54
3	B	3510	ADP	PB-O2B	2.50	1.65	1.54
3	A	3500	ADP	O4'-C1'	3.12	1.45	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	3500	ADP	N3-C2-N1	-11.01	119.27	128.86
3	B	3510	ADP	N3-C2-N1	-11.01	119.27	128.86
3	A	3500	ADP	C4-C5-N7	-4.84	104.73	109.41
3	B	3510	ADP	C4-C5-N7	-4.83	104.74	109.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3500	ADP	1	0
3	B	3510	ADP	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	339/407 (83%)	-0.05	9 (2%) 55 58	19, 41, 80, 117	0
1	B	336/407 (82%)	-0.07	7 (2%) 64 66	20, 40, 70, 107	0
All	All	675/814 (82%)	-0.06	16 (2%) 59 61	19, 41, 74, 117	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1366	TYR	6.8
1	A	1177	PRO	6.0
1	A	1036	ALA	5.4
1	B	1002	ASN	5.2
1	A	1003	ALA	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ADP	A	3500	27/27	0.98	0.13	-0.38	16,26,35,38	0
3	ADP	B	3510	27/27	0.97	0.14	-0.40	16,26,35,38	0
2	MG	B	4611	1/1	0.96	0.10	-1.59	47,47,47,47	0
2	MG	A	4601	1/1	0.97	0.09	-	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.