



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 06:33 pm GMT

PDB ID : 1JOO  
Title : Averaged structure for unligated Staphylococcal nuclease-H124L  
Authors : Wang, J.; Truckses, D.M.; Abildgaard, F.; Dzakula, Z.; Zolnai, Z.; Markley, J.L.  
Deposited on : 2001-07-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	149	 54% 34% 12% .

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2397 atoms, of which 1216 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called staphylococcal nuclease.

Mol	Chain	Residues	Atoms						Trace
1	A	149	Total	C	H	N	O	S	0
			2397	747	1216	206	224	4	


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	LEU	HIS	ENGINEERED	UNP P00644

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: staphylococcal nuclease



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *Distance, hydrogen bond and torsion angle constraints*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.1
INSIGHTii	structure solution	2.3
HBPLUS	structure solution	
X-PLOR	refinement	3.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4053, BMRB entry 4052
Number of chemical shift lists	2
Total number of shifts	3323
Number of shifts mapped to atoms	3323
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	5
All	All	0	5

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	87	ARG	Sidechain
1	A	105	ARG	Sidechain
1	A	81	ARG	Sidechain
1	A	35	ARG	Sidechain
1	A	126	ARG	Sidechain

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1181	1216	1216	42
All	All	1181	1216	1216	42

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:24:LYS:O	1:A:25:LEU:HD23	0.86	1.70
1:A:100:ASN:O	1:A:104:VAL:HG23	0.76	1.81
1:A:62:THR:O	1:A:66:VAL:HG13	0.69	1.88
1:A:36:LEU:HB3	1:A:39:VAL:HG21	0.63	1.68
1:A:46:HIS:N	1:A:47:PRO:CD	0.61	2.64
1:A:37:LEU:O	1:A:39:VAL:HG23	0.60	1.97
1:A:24:LYS:C	1:A:25:LEU:HD23	0.58	2.18
1:A:93:TYR:CZ	1:A:98:MET:SD	0.57	2.98
1:A:53:LYS:O	1:A:54:TYR:CD1	0.57	2.57
1:A:62:THR:O	1:A:66:VAL:HG22	0.57	2.00
1:A:76:PHE:CD1	1:A:90:ALA:HB2	0.49	2.42
1:A:133:LYS:HG2	1:A:140:TRP:CE2	0.48	2.43
1:A:62:THR:O	1:A:66:VAL:CG1	0.47	2.60
1:A:4:THR:O	1:A:5:LYS:CB	0.47	2.62
1:A:51:VAL:HG11	1:A:55:GLY:N	0.47	2.24
1:A:37:LEU:O	1:A:39:VAL:CG2	0.47	2.63
1:A:46:HIS:CB	1:A:47:PRO:HD3	0.47	2.39
1:A:100:ASN:OD1	1:A:100:ASN:N	0.47	2.47
1:A:133:LYS:HD3	1:A:140:TRP:CZ3	0.46	2.45
1:A:15:ILE:HG13	1:A:26:MET:N	0.46	2.26
1:A:133:LYS:HD3	1:A:140:TRP:CE3	0.45	2.47
1:A:69:ALA:HB2	1:A:94:ALA:HB1	0.45	1.89
1:A:16:LYS:O	1:A:16:LYS:HG3	0.44	2.12
1:A:27:TYR:CD2	1:A:28:LYS:HG3	0.44	2.48
1:A:68:ASN:O	1:A:70:LYS:N	0.44	2.50
1:A:39:VAL:O	1:A:112:ALA:HB2	0.44	2.12
1:A:77:ASP:OD1	1:A:120:THR:OG1	0.43	2.34
1:A:101:GLU:OE1	1:A:124:LEU:HD21	0.43	2.13
1:A:16:LYS:O	1:A:16:LYS:CG	0.43	2.67
1:A:27:TYR:CE2	1:A:28:LYS:HG3	0.43	2.49
1:A:51:VAL:HG11	1:A:55:GLY:CA	0.43	2.44
1:A:112:ALA:O	1:A:113:TYR:C	0.42	2.58
1:A:68:ASN:ND2	1:A:68:ASN:N	0.42	2.66
1:A:3:SER:OG	1:A:4:THR:N	0.41	2.53
1:A:133:LYS:HG2	1:A:140:TRP:CD2	0.41	2.51
1:A:35:ARG:O	1:A:36:LEU:C	0.41	2.59
1:A:43:GLU:OE1	1:A:43:GLU:N	0.41	2.54
1:A:145:ALA:O	1:A:146:ASP:C	0.41	2.57
1:A:1:ALA:O	1:A:2:THR:O	0.41	2.38
1:A:36:LEU:HB3	1:A:39:VAL:CG2	0.41	2.44
1:A:94:ALA:O	1:A:95:ASP:HB2	0.40	2.16
1:A:62:THR:O	1:A:66:VAL:CG2	0.40	2.69



## 6.3 Torsion angles

### 6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/149 (99%)	120 (82%)	22 (15%)	5 (3%)	7	38
All	All	147/149 (99%)	120 (82%)	22 (15%)	5 (3%)	7	38

All 5 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	2	THR
1	A	80	GLN
1	A	83	ASP
1	A	118	ASN
1	A	69	ALA

### 6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/125 (100%)	92 (74%)	33 (26%)	2	23
All	All	125/125 (100%)	92 (74%)	33 (26%)	2	23

All 33 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	28	LYS
1	A	39	VAL
1	A	133	LYS
1	A	126	ARG
1	A	113	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	26	MET
1	A	33	THR
1	A	57	GLU
1	A	105	ARG
1	A	41	THR
1	A	127	LYS
1	A	65	MET
1	A	83	ASP
1	A	4	THR
1	A	13	THR
1	A	49	LYS
1	A	59	SER
1	A	131	GLN
1	A	6	LYS
1	A	110	LYS
1	A	147	SER
1	A	37	LEU
1	A	46	HIS
1	A	136	LYS
1	A	48	LYS
1	A	5	LYS
1	A	143	ASP
1	A	2	THR
1	A	43	GLU
1	A	24	LYS
1	A	78	LYS
1	A	16	LYS
1	A	124	LEU

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.6 Ligand geometry

There are no ligands in this entry.

## 6.7 Other polymers

There are no such molecules in this entry.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 80% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 4052

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1655
Number of shifts mapped to atoms	1655
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	136	$1.27 \pm 0.10$	Should be applied
$^{13}\text{C}_\beta$	112	$1.64 \pm 0.15$	Should be applied
$^{13}\text{C}'$	130	$1.87 \pm 0.12$	Should be applied
$^{15}\text{N}$	133	$-1.38 \pm 0.47$	Should be applied

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 1458 atoms were assigned a chemical shift out of a possible 1877. 18 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	663/733 (90%)	264/292 (90%)	266/298 (89%)	133/143 (93%)
Sidechain	688/1025 (67%)	502/603 (83%)	181/372 (49%)	5/50 (10%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	107/119 (90%)	54/61 (89%)	49/51 (96%)	4/7 (57%)
Overall	1458/1877 (78%)	820/956 (86%)	496/721 (69%)	142/200 (71%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 78%, i.e. 1458 atoms were assigned a chemical shift out of a possible 1877. 18 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	663/733 (90%)	264/292 (90%)	266/298 (89%)	133/143 (93%)
Sidechain	688/1025 (67%)	502/603 (83%)	181/372 (49%)	5/50 (10%)
Aromatic	107/119 (90%)	54/61 (89%)	49/51 (96%)	4/7 (57%)
Overall	1458/1877 (78%)	820/956 (86%)	496/721 (69%)	142/200 (71%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	121	HIS	HD2	4.33	9.28 – 4.78	-6.0
1	A	9	LYS	HE2	1.80	3.87 – 1.97	-5.9
1	A	9	LYS	HE3	1.80	3.86 – 1.96	-5.8

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: BMRB entry 4053

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1668
Number of shifts mapped to atoms	1668
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	137	$1.36 \pm 0.15$	Should be applied
$^{13}\text{C}_\beta$	120	$1.64 \pm 0.12$	Should be applied
$^{13}\text{C}'$	132	$1.77 \pm 0.10$	Should be applied
$^{15}\text{N}$	127	$-1.15 \pm 0.47$	Should be applied

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 1467 atoms were assigned a chemical shift out of a possible 1877. 21 out of 21 assigned methyl groups (LEU and VAL) were assigned

stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	660/733 (90%)	264/292 (90%)	269/298 (90%)	127/143 (89%)
Sidechain	701/1025 (68%)	503/603 (83%)	193/372 (52%)	5/50 (10%)
Aromatic	106/119 (89%)	54/61 (89%)	47/51 (92%)	5/7 (71%)
Overall	1467/1877 (78%)	821/956 (86%)	509/721 (71%)	137/200 (68%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 78%, i.e. 1467 atoms were assigned a chemical shift out of a possible 1877. 21 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	660/733 (90%)	264/292 (90%)	269/298 (90%)	127/143 (89%)
Sidechain	701/1025 (68%)	503/603 (83%)	193/372 (52%)	5/50 (10%)
Aromatic	106/119 (89%)	54/61 (89%)	47/51 (92%)	5/7 (71%)
Overall	1467/1877 (78%)	821/956 (86%)	509/721 (71%)	137/200 (68%)

## 7.2.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	80	GLN	CG	125.57	39.38 – 28.18	82.0
1	A	117	PRO	CD	58.57	55.31 – 45.41	8.3
1	A	121	HIS	HD2	4.32	9.28 – 4.78	-6.0
1	A	9	LYS	HE2	1.82	3.87 – 1.97	-5.8
1	A	9	LYS	HE3	1.82	3.86 – 1.96	-5.7

## 7.2.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

