



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 2, 2017 – 09:31 PM EST

PDB ID : 5JRI
Title : Structure of an oxidoreductase SeMet-labelled from *Synechocystis* sp. PCC6803
Authors : Buey, R.M.; de Pereda, J.M.; Balsera, M.
Deposited on : unknown
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 94% 5% </div> </div>
1	B	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 92% 5% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	410	-	-	-	X
3	SO4	B	403	-	-	-	X
3	SO4	B	405	-	-	-	X
4	EDO	A	408	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10299 atoms, of which 4797 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridine nucleotide-disulfide oxidoreductase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	315	Total	C	H	N	O	S	Se	0	0	0
			4742	1539	2333	409	447	4	10			
1	B	313	Total	C	H	N	O	S	Se	0	0	0
			4724	1529	2326	407	448	4	10			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP A0A0P0GNW3
A	2	SER	-	expression tag	UNP A0A0P0GNW3
A	3	HIS	-	expression tag	UNP A0A0P0GNW3
A	4	MSE	-	expression tag	UNP A0A0P0GNW3
A	5	VAL	-	expression tag	UNP A0A0P0GNW3
A	244	MSE	VAL	conflict	UNP A0A0P0GNW3
B	1	GLY	-	expression tag	UNP A0A0P0GNW3
B	2	SER	-	expression tag	UNP A0A0P0GNW3
B	3	HIS	-	expression tag	UNP A0A0P0GNW3
B	4	MSE	-	expression tag	UNP A0A0P0GNW3
B	5	VAL	-	expression tag	UNP A0A0P0GNW3
B	244	MSE	VAL	conflict	UNP A0A0P0GNW3

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		
2	A	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		
2	B	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		
2	B	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 10 2 6 2	0	0
4	A	1	Total C H O 10 2 6 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	223	Total	O	0	0
			223	223		
5	B	198	Total	O	0	0
			198	198		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyridine nucleotide-disulfide oxidoreductase

Chain A:  94% 5%



- Molecule 1: Pyridine nucleotide-disulfide oxidoreductase

Chain B:  92% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.73 Å 79.33 Å 127.05 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.08 – 1.95 49.59 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.08-1.95) 99.9 (49.59-1.95)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.95 Å)	Xtriage
Refinement program	PHENIX (dev_2341: ???)	Depositor
R, R_{free}	0.199 , 0.242 0.201 , 0.239	Depositor DCC
R_{free} test set	2651 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10299	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.22 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.8191e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2452	0.56	0/3318
1	B	0.35	0/2441	0.51	0/3304
All	All	0.39	0/4893	0.53	0/6622

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2409	2333	2343	5	0
1	B	2398	2326	2325	7	0
2	A	106	60	62	3	0
2	B	106	60	62	3	0
3	A	30	0	0	0	0
3	B	20	0	0	1	0
4	A	8	12	12	0	0
4	B	4	6	6	0	0
5	A	223	0	0	1	0
5	B	198	0	0	0	0
All	All	5502	4797	4810	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 13 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:267:GLY:O	1:B:284:SER:OG	2.00	0.80
1:A:16:LEU:N	5:A:505:HOH:O	2.32	0.62
1:A:156:VAL:HG21	2:A:401:FAD:HM83	1.84	0.59
1:B:22:ALA:O	1:B:45:CYS:HA	2.09	0.52
1:B:156:VAL:HG21	2:B:401:FAD:HM83	1.92	0.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/330 (95%)	305 (97%)	8 (3%)	0	100	100
1	B	311/330 (94%)	306 (98%)	5 (2%)	0	100	100
All	All	624/660 (94%)	611 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/258 (95%)	244 (100%)	0	100	100
1	B	244/258 (95%)	244 (100%)	0	100	100
All	All	488/516 (95%)	488 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	401	-	50,58,58	1.35	6 (12%)	53,89,89	2.08	10 (18%)
2	FAD	A	402	-	50,58,58	1.34	6 (12%)	53,89,89	2.04	9 (16%)
3	SO4	A	403	-	4,4,4	0.35	0	6,6,6	0.14	0
3	SO4	A	404	-	4,4,4	0.88	0	6,6,6	0.57	0
3	SO4	A	405	-	4,4,4	0.86	0	6,6,6	0.57	0
3	SO4	A	406	-	4,4,4	0.86	0	6,6,6	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	407	-	4,4,4	0.42	0	6,6,6	0.12	0
4	EDO	A	408	-	3,3,3	0.69	0	2,2,2	0.78	0
4	EDO	A	409	-	3,3,3	0.43	0	2,2,2	0.32	0
3	SO4	A	410	-	4,4,4	0.32	0	6,6,6	0.08	0
2	FAD	B	401	-	50,58,58	1.34	6 (12%)	53,89,89	2.08	10 (18%)
2	FAD	B	402	-	50,58,58	1.38	6 (12%)	53,89,89	2.07	9 (16%)
3	SO4	B	403	-	4,4,4	0.27	0	6,6,6	0.13	0
3	SO4	B	404	-	4,4,4	0.90	0	6,6,6	0.64	0
3	SO4	B	405	-	4,4,4	0.85	0	6,6,6	0.79	0
3	SO4	B	406	-	4,4,4	0.35	0	6,6,6	0.09	0
4	EDO	B	407	-	3,3,3	0.66	0	2,2,2	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	401	-	-	0/28/50/50	0/6/6/6
2	FAD	A	402	-	-	0/28/50/50	0/6/6/6
3	SO4	A	403	-	-	0/0/0/0	0/0/0/0
3	SO4	A	404	-	-	0/0/0/0	0/0/0/0
3	SO4	A	405	-	-	0/0/0/0	0/0/0/0
3	SO4	A	406	-	-	0/0/0/0	0/0/0/0
3	SO4	A	407	-	-	0/0/0/0	0/0/0/0
4	EDO	A	408	-	-	0/1/1/1	0/0/0/0
4	EDO	A	409	-	-	0/1/1/1	0/0/0/0
3	SO4	A	410	-	-	0/0/0/0	0/0/0/0
2	FAD	B	401	-	-	0/28/50/50	0/6/6/6
2	FAD	B	402	-	-	0/28/50/50	0/6/6/6
3	SO4	B	403	-	-	0/0/0/0	0/0/0/0
3	SO4	B	404	-	-	0/0/0/0	0/0/0/0
3	SO4	B	405	-	-	0/0/0/0	0/0/0/0
3	SO4	B	406	-	-	0/0/0/0	0/0/0/0
4	EDO	B	407	-	-	0/1/1/1	0/0/0/0

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	FAD	C9A-N10	2.54	1.42	1.38
2	A	402	FAD	C9A-N10	2.65	1.42	1.38
2	A	401	FAD	C9A-N10	2.68	1.42	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	FAD	C5A-C4A	2.79	1.46	1.40
2	B	402	FAD	C9A-N10	2.80	1.42	1.38

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FAD	N3A-C2A-N1A	-6.87	122.87	128.86
2	A	401	FAD	N3A-C2A-N1A	-6.79	122.94	128.86
2	A	402	FAD	N3A-C2A-N1A	-6.67	123.05	128.86
2	B	402	FAD	N3A-C2A-N1A	-6.53	123.17	128.86
2	B	401	FAD	C4A-C5A-N7A	-3.74	105.80	109.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FAD	2	0
2	A	402	FAD	1	0
2	B	401	FAD	1	0
2	B	402	FAD	2	0
3	B	406	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/330 (92%)	0.36	3 (0%) 82 88	21, 30, 55, 87	0
1	B	303/330 (91%)	0.32	1 (0%) 93 96	22, 31, 50, 68	0
All	All	608/660 (92%)	0.34	4 (0%) 87 92	21, 31, 52, 87	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	LEU	5.9
1	A	175	GLY	2.3
1	B	175	GLY	2.1
1	A	201	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	410	5/5	0.84	0.17	7.20	71,89,97,100	0
3	SO4	B	405	5/5	0.91	0.20	4.55	70,81,86,88	0
3	SO4	B	403	5/5	0.84	0.21	4.24	51,67,71,78	0
4	EDO	A	408	4/4	0.90	0.20	3.73	36,58,70,72	0
3	SO4	A	404	5/5	0.87	0.13	1.15	35,47,79,84	0
2	FAD	A	401	53/53	0.94	0.15	1.01	15,27,38,44	0
2	FAD	B	401	53/53	0.94	0.14	0.86	15,28,36,47	0
3	SO4	B	404	5/5	0.89	0.14	0.52	35,46,72,79	0
4	EDO	B	407	4/4	0.93	0.13	0.11	35,55,66,66	0
2	FAD	A	402	53/53	0.94	0.13	0.04	20,31,46,63	0
2	FAD	B	402	53/53	0.94	0.12	-0.53	19,33,46,61	0
4	EDO	A	409	4/4	0.85	0.33	-	20,20,20,20	0
3	SO4	A	403	5/5	0.91	0.13	-	35,44,62,66	0
3	SO4	B	406	5/5	0.80	0.23	-	129,130,131,133	0
3	SO4	A	406	5/5	0.65	0.18	-	101,109,117,124	0
3	SO4	A	405	5/5	0.80	0.20	-	88,103,106,106	0
3	SO4	A	407	5/5	0.70	0.40	-	20,20,20,20	0

6.5 Other polymers ⓘ

There are no such residues in this entry.