



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:12 pm GMT

PDB ID : 1JTU  
Title : E. coli Thymidylate Synthase in a Complex with dUMP and LY338913, A Polyglutamylated Pyrrolo(2,3-d)pyrimidine-based Antifolate  
Authors : Sayre, P.H.; Finer-Moore, J.S.; Fritz, T.A.; Biermann, D.; Gates, S.B.; MacKellar, W.C.; Patel, V.F.; Stroud, R.M.  
Deposited on : 2001-08-22  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	
1	B	264	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LYB	B	502	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2153	1375	371	395	12			
1	B	264	Total	C	N	O	S	0	0	0
			2153	1375	371	395	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
B	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $C_9H_{13}N_2O_8P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	20	9	2	8	1	0	0

- Molecule 3 is 2-{4-[4-(4-{4-[2-(2-AMINO-4-OXO-4,7-DIHYDRO-3H-PYRROLO[2,3-D]PYRIMIDIN-5-YL)-ETHYL]-BENZOYLAMINO}-4-CARBOXY-BUTYRYLAMINO)-4-CARBOXY-BUTYRYLAMINO}-PENTANEDIOIC ACID (three-letter code: LYB) (formula: C<sub>35</sub>H<sub>42</sub>N<sub>8</sub>O<sub>15</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	49	30	7	12	0	0
			49	30	7	12		
3	B	1	49	30	7	12	0	0
			49	30	7	12		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	112	Total	O	0	0
			112	112		
4	B	84	Total	O	0	0
			84	84		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: THYMIDYLATE SYNTHASE



#### • Molecule 1: THYMIDYLATE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.60Å 127.60Å 68.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.50 – 2.20 32.50 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.7 (32.50-2.20) 94.5 (32.50-2.20)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.20Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.198 , 0.241 0.198 , 0.240	Depositor DCC
$R_{free}$ test set	3060 reflections (10.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4640	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UMP, LYB, CXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2202	0.65	0/2990
1	B	0.33	0/2202	0.61	0/2990
All	All	0.34	0/4404	0.63	0/5980

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2079	48	0
1	B	2153	0	2079	57	0
2	A	20	0	10	0	0
2	B	20	0	10	0	0
3	A	49	0	31	1	0
3	B	49	0	31	2	0
4	A	112	0	0	5	0
4	B	84	0	0	6	0
All	All	4640	0	4240	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:GLN:NE2	1:A:215:GLN:H	1.60	0.98
1:A:215:GLN:HE21	1:A:215:GLN:H	1.13	0.95
1:B:215:GLN:HE21	1:B:215:GLN:H	1.15	0.89
1:A:222:ARG:NH1	1:A:256:PRO:HD2	1.95	0.81
1:A:215:GLN:N	1:A:215:GLN:HE21	1.82	0.77
1:B:228:PRO:HG3	1:B:249:ILE:HD11	1.70	0.73
1:B:215:GLN:NE2	1:B:215:GLN:H	1.87	0.72
1:B:243:ARG:HE	1:B:244:PHE:H	1.40	0.70
1:B:60:LEU:O	1:B:64:GLN:HG2	1.91	0.69
1:A:134:ASN:C	1:A:134:ASN:HD22	1.95	0.69
1:B:2:LYS:O	1:B:6:GLU:HG3	1.94	0.68
1:A:45:VAL:HG22	1:A:50:CYS:SG	2.34	0.67
1:A:44:LEU:HD21	1:A:52:LEU:HD21	1.78	0.66
1:B:237:GLU:HB2	4:B:672:HOH:O	1.96	0.64
1:B:134:ASN:HD22	1:B:134:ASN:C	2.03	0.61
1:B:61:TRP:CD1	1:B:66:ASP:HB3	2.36	0.60
1:A:245:GLU:CD	1:A:245:GLU:H	2.06	0.59
1:A:170:VAL:HB	1:A:208:LEU:HD13	1.84	0.59
1:A:15:GLY:HA3	1:A:27:LEU:HD22	1.85	0.59
1:A:222:ARG:HH12	1:A:256:PRO:HD2	1.68	0.58
1:B:228:PRO:CG	1:B:249:ILE:HD11	2.33	0.58
1:B:56:ILE:O	1:B:60:LEU:HG	2.04	0.58
1:A:170:VAL:HB	1:A:208:LEU:CD1	2.34	0.57
1:B:80:TRP:CH2	3:B:502:LYB:HC6	2.39	0.57
1:A:51:HIS:CE1	1:A:53:ARG:HB3	2.38	0.57
1:B:215:GLN:HE21	1:B:215:GLN:N	1.95	0.57
1:A:222:ARG:HH21	1:A:258:ILE:HD11	1.69	0.57
1:B:58:GLU:O	1:B:61:TRP:HB3	2.05	0.57
1:A:58:GLU:O	1:A:61:TRP:HB3	2.05	0.56
1:B:84:ALA:HA	1:B:90:LEU:HD23	1.88	0.56
1:A:110:ASP:OD2	1:A:113:THR:HG23	2.05	0.55
1:A:212:HIS:CD2	1:A:260:ALA:HB1	2.42	0.55
1:B:224:PRO:HD3	4:B:801:HOH:O	2.06	0.55
1:A:145:PRO:HG3	4:A:825:HOH:O	2.05	0.55
1:B:45:VAL:HG23	1:B:50:CYS:SG	2.47	0.55
1:B:186:HIS:CE1	1:B:196:VAL:HG11	2.42	0.55
1:B:243:ARG:NE	1:B:244:PHE:H	2.04	0.54
1:A:37:ASN:C	1:A:37:ASN:HD22	2.08	0.54
1:A:233:LYS:HE3	1:A:248:GLU:HB2	1.91	0.53
1:A:172:LEU:HD21	1:A:262:VAL:HG22	1.90	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:LEU:HD11	1:A:47:THR:HG21	1.90	0.53
1:A:256:PRO:HA	4:A:748:HOH:O	2.09	0.52
1:A:96:LYS:HE2	1:A:100:ALA:O	2.10	0.51
1:A:116:LEU:O	1:A:120:LYS:HG3	2.11	0.51
1:A:133:TRP:CZ2	1:A:138:LEU:HD21	2.46	0.50
1:A:73:HIS:HE1	1:A:81:ASP:OD1	1.94	0.50
1:B:192:CYS:O	1:B:194:LEU:HG	2.10	0.50
1:B:48:LYS:HE3	4:B:676:HOH:O	2.12	0.49
1:A:222:ARG:NH2	1:A:258:ILE:HD11	2.28	0.49
1:A:109:ILE:HD12	1:B:135:VAL:HG12	1.95	0.48
1:A:45:VAL:CG2	1:A:50:CYS:SG	3.01	0.48
1:B:255:HIS:HB3	1:B:256:PRO:HD2	1.94	0.48
1:A:37:ASN:ND2	1:A:39:GLN:H	2.11	0.48
1:B:211:ASN:ND2	1:B:261:PRO:HG2	2.29	0.48
1:B:163:LEU:HD22	1:B:164:TYR:N	2.29	0.47
1:B:174:LEU:O	1:B:178:ILE:HG13	2.14	0.47
1:B:111:GLN:OE1	1:B:130:VAL:HA	2.15	0.47
1:B:163:LEU:CD2	1:B:164:TYR:N	2.78	0.47
1:B:234:ARG:HG2	1:B:246:ASP:OD1	2.15	0.46
1:B:11:VAL:HG11	1:B:208:LEU:HB2	1.96	0.46
3:B:502:LYB:H12	4:B:741:HOH:O	2.16	0.46
1:A:134:ASN:ND2	1:A:134:ASN:C	2.67	0.46
1:A:1:CXM:HE2	1:A:45:VAL:N	2.31	0.45
1:A:1:CXM:HE1	1:A:45:VAL:HG12	1.99	0.45
1:B:56:ILE:HG12	1:B:183:LEU:HD11	1.98	0.45
1:A:174:LEU:HB3	1:A:175:PRO:HD3	1.99	0.45
1:A:136:GLY:HA3	4:B:749:HOH:O	2.17	0.45
1:B:225:ARG:O	1:B:226:PRO:C	2.55	0.45
1:B:227:LEU:H	1:B:227:LEU:HD22	1.82	0.45
1:B:116:LEU:HD12	1:B:120:LYS:HE3	2.00	0.44
1:B:128:ILE:CG2	1:B:152:PHE:HB2	2.48	0.44
1:B:45:VAL:CG2	1:B:50:CYS:SG	3.05	0.44
1:B:37:ASN:OD1	1:B:39:GLN:HB2	2.19	0.43
1:B:59:LEU:HD21	1:B:187:MET:HE3	2.01	0.43
1:A:27:LEU:HD12	1:A:213:MET:CE	2.49	0.43
1:B:57:HIS:NE2	1:B:75:ASN:ND2	2.67	0.43
1:B:122:ASP:OD2	1:B:125:SER:HB2	2.18	0.43
1:B:111:GLN:O	1:B:115:VAL:HG23	2.19	0.43
1:B:201:TRP:CH2	1:B:203:GLY:HA3	2.54	0.43
1:B:242:TYR:HB2	4:B:681:HOH:O	2.19	0.42
1:B:192:CYS:O	1:B:193:ASP:C	2.58	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ASP:OD2	1:A:107:ARG:HD3	2.20	0.42
1:B:170:VAL:HG23	1:B:207:HIS:O	2.20	0.42
1:A:79:ILE:HG13	1:A:80:TRP:CD1	2.54	0.42
1:A:27:LEU:HA	1:A:27:LEU:HD23	1.88	0.42
1:A:92:PRO:HG2	1:A:140:LYS:HE3	2.01	0.42
3:A:501:LYB:H12	4:A:699:HOH:O	2.19	0.41
1:B:230:LEU:C	1:B:231:ILE:HD12	2.40	0.41
1:B:78:THR:HB	1:B:81:ASP:OD2	2.20	0.41
4:A:764:HOH:O	1:B:104:PRO:HG2	2.21	0.41
1:B:10:LYS:HG3	1:B:14:GLU:OE2	2.21	0.41
1:B:69:ILE:HG21	1:B:80:TRP:O	2.20	0.41
1:A:9:GLN:HE21	1:A:13:ASP:CG	2.23	0.41
1:B:233:LYS:HE3	1:B:248:GLU:CD	2.40	0.41
1:A:10:LYS:HE3	1:A:32:HIS:CD2	2.56	0.41
1:B:230:LEU:HD13	1:B:231:ILE:N	2.36	0.41
1:B:27:LEU:HD12	1:B:27:LEU:HA	1.91	0.41
1:A:260:ALA:HA	1:A:261:PRO:HD3	1.90	0.41
1:B:223:GLU:HA	1:B:224:PRO:HD3	1.95	0.41
1:B:245:GLU:N	1:B:245:GLU:OE1	2.54	0.41
1:A:116:LEU:HD11	1:A:239:ILE:HB	2.02	0.40
1:A:37:ASN:HD22	1:A:39:GLN:H	1.68	0.40
1:A:82:GLU:HG3	4:A:828:HOH:O	2.21	0.40
1:B:26:THR:HG22	1:B:209:TYR:CD1	2.57	0.40
1:A:186:HIS:CD2	1:A:196:VAL:HG11	2.56	0.40
1:B:72:LEU:HB3	1:B:77:VAL:HB	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	262/264 (99%)	245 (94%)	16 (6%)	1 (0%)	38 41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	262/264 (99%)	241 (92%)	17 (6%)	4 (2%)	12	9
All	All	524/528 (99%)	486 (93%)	33 (6%)	5 (1%)	18	16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	226	PRO
1	A	94	TYR
1	B	94	TYR
1	B	82	GLU
1	B	93	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	216 (93%)	16 (7%)	18	19
1	B	232/232 (100%)	220 (95%)	12 (5%)	27	32
All	All	464/464 (100%)	436 (94%)	28 (6%)	22	25

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	10	LYS
1	A	27	LEU
1	A	37	ASN
1	A	39	GLN
1	A	60	LEU
1	A	134	ASN
1	A	138	LEU
1	A	166	ARG
1	A	208	LEU
1	A	215	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	223	GLU
1	A	225	ARG
1	A	230	LEU
1	A	241	ASP
1	A	254	PRO
1	B	38	LEU
1	B	52	LEU
1	B	53	ARG
1	B	116	LEU
1	B	130	VAL
1	B	134	ASN
1	B	163	LEU
1	B	183	LEU
1	B	215	GLN
1	B	222	ARG
1	B	226	PRO
1	B	243	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	HIS
1	A	33	GLN
1	A	37	ASN
1	A	73	HIS
1	A	75	ASN
1	A	87	ASN
1	A	97	GLN
1	A	117	ASN
1	A	118	GLN
1	A	134	ASN
1	A	151	GLN
1	A	162	GLN
1	A	215	GLN
1	B	32	HIS
1	B	51	HIS
1	B	64	GLN
1	B	75	ASN
1	B	76	ASN
1	B	117	ASN
1	B	118	GLN
1	B	134	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	151	GLN
1	B	215	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CXM	A	1	1	7,10,11	1.07	1 (14%)	5,11,13	1.39	0
1	CXM	B	1	1	7,10,11	1.35	2 (28%)	5,11,13	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
1	CXM	B	1	1	-	0/6/10/12	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	CXM	CA-C	2.17	1.53	1.50
1	A	1	CXM	CA-N	2.22	1.49	1.46
1	B	1	CXM	CA-N	2.46	1.49	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1	CXM	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	LYB	A	501	-	40,50,60	1.88	7 (17%)	42,68,82	2.33	7 (16%)
2	UMP	A	565	1	17,21,21	3.53	5 (29%)	23,31,31	3.34	6 (26%)
3	LYB	B	502	-	40,50,60	1.82	8 (20%)	42,68,82	2.20	4 (9%)
2	UMP	B	565	1	17,21,21	3.48	5 (29%)	23,31,31	3.10	6 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LYB	A	501	-	-	0/34/47/61	0/3/3/3
2	UMP	A	565	1	-	0/6/22/22	0/2/2/2
3	LYB	B	502	-	-	0/34/47/61	0/3/3/3
2	UMP	B	565	1	-	0/6/22/22	0/2/2/2

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	LYB	C22-N6	2.03	1.49	1.46
3	B	502	LYB	C1-N1	2.04	1.39	1.35
3	B	502	LYB	C7-C4	2.11	1.44	1.41
3	A	501	LYB	C7-C4	2.23	1.44	1.41
3	B	502	LYB	C3-N3	2.27	1.39	1.34
3	A	501	LYB	C6-N3	2.28	1.41	1.36
2	B	565	UMP	O4'-C4'	2.43	1.50	1.45
3	B	502	LYB	C6-N3	2.46	1.41	1.36
3	A	501	LYB	C3-N3	2.56	1.39	1.34
2	A	565	UMP	O4'-C4'	2.63	1.51	1.45
3	A	501	LYB	C1-N1	2.75	1.40	1.35
3	B	502	LYB	C3-N2	2.92	1.42	1.36
2	B	565	UMP	C4-N3	3.10	1.38	1.33
2	A	565	UMP	C4-N3	3.13	1.38	1.33
3	A	501	LYB	C3-N2	3.14	1.42	1.36
2	A	565	UMP	O4'-C1'	4.02	1.51	1.42
2	B	565	UMP	O4'-C1'	4.11	1.51	1.42
3	B	502	LYB	C5-N1	5.11	1.42	1.33
3	A	501	LYB	C5-N1	5.22	1.42	1.33
3	B	502	LYB	C1-N4	6.58	1.47	1.34
3	A	501	LYB	C1-N4	7.37	1.49	1.34
2	A	565	UMP	C6-C5	7.77	1.54	1.38
2	B	565	UMP	C6-C5	7.81	1.54	1.38
2	B	565	UMP	C6-N1	10.18	1.49	1.35
2	A	565	UMP	C6-N1	10.35	1.49	1.35

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	LYB	C4-C5-N1	-8.61	115.85	124.12
3	A	501	LYB	C4-C5-N1	-8.55	115.91	124.12
2	B	565	UMP	C5-C6-N1	-4.68	109.94	120.67
3	B	502	LYB	N2-C1-N1	-3.23	122.73	127.46
3	A	501	LYB	N2-C1-N1	-3.21	122.77	127.46
2	B	565	UMP	O4'-C4'-C3'	-2.61	99.41	105.68
2	A	565	UMP	C5-C6-N1	-2.58	114.76	120.67
3	A	501	LYB	O5-C16-N5	-2.50	117.90	122.46
3	A	501	LYB	C7-C4-C3	-2.24	106.46	110.22
2	B	565	UMP	O4'-C1'-C2'	-2.09	102.23	106.25
2	A	565	UMP	O4'-C4'-C3'	-2.00	100.88	105.68
3	A	501	LYB	O5-C16-C13	2.03	124.55	120.94
2	A	565	UMP	P-O5'-C5'	2.26	124.51	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	565	UMP	C6-N1-C2	2.35	125.09	121.28
2	B	565	UMP	C6-N1-C2	2.63	125.54	121.28
3	A	501	LYB	C5-N1-C1	3.76	121.47	116.06
2	A	565	UMP	O4'-C1'-N1	4.10	114.69	107.78
3	B	502	LYB	C5-N1-C1	4.12	121.98	116.06
2	B	565	UMP	O4'-C1'-N1	4.83	115.92	107.78
3	B	502	LYB	C5-C4-C3	8.66	119.89	115.02
3	A	501	LYB	C5-C4-C3	9.46	120.34	115.02
2	B	565	UMP	C4-N3-C2	11.68	124.17	114.13
2	A	565	UMP	C4-N3-C2	14.33	126.44	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	LYB	1	0
3	B	502	LYB	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/264 (99%)	-0.48	0 100 100	13, 30, 47, 55	0
1	B	263/264 (99%)	-0.19	5 (1%) 67 65	17, 38, 67, 74	0
All	All	526/528 (99%)	-0.34	5 (0%) 82 81	13, 34, 61, 74	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	86	GLU	3.4
1	B	88	GLY	2.4
1	B	243	ARG	2.4
1	B	233	LYS	2.4
1	B	264	ILE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CXM	B	1	11/12	0.94	0.14	-	39,40,48,48	0
1	CXM	A	1	11/12	0.95	0.11	-	30,33,34,36	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	LYB	B	502	49/58	0.59	0.26	3.20	68,81,97,99	0
3	LYB	A	501	49/58	0.78	0.19	1.16	37,60,100,103	0
2	UMP	A	565	20/20	0.98	0.10	-0.74	28,32,33,33	0
2	UMP	B	565	20/20	0.96	0.10	-0.82	33,43,45,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.