



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 10:25 pm GMT

PDB ID : 2JTY
Title : Self-complemented variant of FimA, the main subunit of type 1 pilus
Authors : Erilov, D.; Wider, G.; Glockshuber, R.; Puorger, C.; Vetsch, M.
Deposited on : 2007-08-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	184	 77% 8% 16%

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:19-A:89, A:95-A:160, A:167-A:184 (155)	0.31	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 6, 9, 14, 16, 17, 20
2	5, 8, 11, 13
3	4, 10, 15, 18
4	2, 3, 19
Single-model clusters	7; 12

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2489 atoms, of which 1222 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Type-1 fimbrial protein, A chain.


Mol	Chain	Residues	Atoms						Trace
1	A	184	Total	C	H	N	O	S	0
			2489	773	1222	226	266	2	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Type-1 fimbrial protein, A chain




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)


- Molecule 1: Type-1 fimbrial protein, A chain



4.2.2 Score per residue for model 2



- Molecule 1: Type-1 fimbrial protein, A chain





4.2.3 Score per residue for model 3

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.4 Score per residue for model 4

- Molecule 1: Type-1 fimbrial protein, A chain



4.2.5 Score per residue for model 5


- Molecule 1: Type-1 fimbrial protein, A chain



4.2.6 Score per residue for model 6


- Molecule 1: Type-1 fimbrial protein, A chain






4.2.7 Score per residue for model 7

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.8 Score per residue for model 8

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.9 Score per residue for model 9

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.10 Score per residue for model 10

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.11 Score per residue for model 11

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.12 Score per residue for model 12

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.13 Score per residue for model 13

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.14 Score per residue for model 14

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.15 Score per residue for model 15

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.16 Score per residue for model 16

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.17 Score per residue for model 17

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.18 Score per residue for model 18

- Molecule 1: Type-1 fimbrial protein, A chain




4.2.19 Score per residue for model 19

- Molecule 1: Type-1 fimbrial protein, A chain



4.2.20 Score per residue for model 20

- Molecule 1: Type-1 fimbrial protein, A chain



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
Opal	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 15423
Number of chemical shift lists	1
Total number of shifts	1932
Number of shifts mapped to atoms	1932
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.54±0.01	0±0/1102 (0.0±0.0%)	1.07±0.04	2±1/1506 (0.1±0.1%)
All	All	0.54	0/22040 (0.0%)	1.07	38/30120 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.0±0.8
All	All	0	20

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	156	VAL	CG1-CB-CG2	8.66	124.76	110.90	5	2
1	A	71	VAL	CG1-CB-CG2	8.57	124.62	110.90	2	10
1	A	137	TYR	CB-CG-CD2	-7.72	116.36	121.00	9	1
1	A	158	TYR	CB-CG-CD2	-7.63	116.42	121.00	11	1
1	A	124	LEU	CB-CG-CD1	6.79	122.54	111.00	15	2
1	A	104	LEU	CB-CG-CD1	6.26	121.64	111.00	17	1
1	A	136	ARG	NE-CZ-NH2	-5.96	117.32	120.30	5	4
1	A	21	CYS	CA-CB-SG	5.76	124.36	114.00	11	1
1	A	52	VAL	CA-CB-CG2	5.74	119.51	110.90	9	3
1	A	74	LEU	CB-CG-CD2	-5.64	101.42	111.00	18	1
1	A	28	VAL	CA-CB-CG1	5.54	119.22	110.90	2	1
1	A	104	LEU	CB-CG-CD2	5.53	120.40	111.00	6	1
1	A	19	ALA	C-N-CA	5.49	135.43	121.70	10	2
1	A	71	VAL	CA-CB-CG1	5.49	119.13	110.90	13	1
1	A	156	VAL	CA-CB-CG1	5.26	118.79	110.90	11	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	167	ALA	CB-CA-C	5.26	117.99	110.10	12	1
1	A	38	ARG	NE-CZ-NH1	5.22	122.91	120.30	2	1
1	A	119	SER	C-N-CA	5.20	134.71	121.70	6	1
1	A	65	VAL	CA-CB-CG2	5.18	118.67	110.90	4	1
1	A	106	ARG	NE-CZ-NH2	-5.18	117.71	120.30	6	1
1	A	38	ARG	NE-CZ-NH2	-5.18	117.71	120.30	17	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	69	ALA	Peptide	5
1	A	152	ALA	Peptide	3
1	A	38	ARG	Sidechain	3
1	A	106	ARG	Sidechain	2
1	A	95	SER	Peptide	2
1	A	119	SER	Peptide	2
1	A	159	GLN	Peptide	1
1	A	46	GLY	Peptide	1
1	A	154	PHE	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1090	1055	1053	5±3
All	All	21800	21100	21060	108

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:86:LEU:HD23	1:A:101:VAL:HG22	0.76	1.57	12	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:GLN:HE21	1:A:47:ALA:HB2	0.76	1.40	14	9
1:A:156:VAL:HG13	1:A:170:VAL:HG23	0.75	1.59	11	2
1:A:156:VAL:HG21	1:A:158:TYR:CE2	0.72	2.19	5	2
1:A:34:LEU:HD23	1:A:52:VAL:CG1	0.72	2.13	3	6
1:A:86:LEU:HD11	1:A:152:ALA:HB2	0.67	1.67	10	2
1:A:34:LEU:HD23	1:A:52:VAL:HG11	0.62	1.71	6	6
1:A:74:LEU:HD22	1:A:153:THR:OG1	0.61	1.95	18	1
1:A:74:LEU:HD23	1:A:74:LEU:C	0.58	2.19	18	1
1:A:124:LEU:HD22	1:A:124:LEU:H	0.57	1.59	18	1
1:A:25:ALA:HA	1:A:28:VAL:CG2	0.56	2.30	20	1
1:A:101:VAL:HG11	1:A:177:PHE:CD2	0.56	2.36	19	1
1:A:74:LEU:O	1:A:152:ALA:HB1	0.55	2.02	10	1
1:A:55:ASN:HB3	1:A:130:THR:HG22	0.55	1.78	13	1
1:A:54:PHE:CE1	1:A:135:ALA:HB2	0.54	2.38	14	6
1:A:156:VAL:CG2	1:A:158:TYR:CE2	0.53	2.91	5	2
1:A:156:VAL:CG1	1:A:170:VAL:HG23	0.52	2.33	5	1
1:A:86:LEU:HD13	1:A:150:ALA:HB1	0.51	1.81	14	3
1:A:73:PHE:CE1	1:A:103:ILE:HG21	0.51	2.40	16	2
1:A:69:ALA:HB2	1:A:124:LEU:HD11	0.50	1.83	19	1
1:A:42:LEU:O	1:A:143:ALA:HB3	0.50	2.06	6	6
1:A:44:GLN:NE2	1:A:47:ALA:HB2	0.50	2.22	2	5
1:A:104:LEU:HD23	1:A:110:ALA:HA	0.50	1.84	13	2
1:A:75:GLY:HA3	1:A:86:LEU:HD21	0.48	1.85	14	3
1:A:68:LYS:CA	1:A:124:LEU:HD21	0.48	2.39	18	2
1:A:31:THR:HG23	1:A:176:HIS:HB2	0.48	1.86	8	2
1:A:124:LEU:H	1:A:124:LEU:HD22	0.47	1.68	13	1
1:A:102:GLN:HE22	1:A:110:ALA:HB1	0.47	1.69	17	1
1:A:20:ALA:HA	1:A:65:VAL:HG11	0.47	1.85	14	1
1:A:156:VAL:HG11	1:A:158:TYR:CZ	0.47	2.45	3	2
1:A:19:ALA:HB1	1:A:158:TYR:CE2	0.47	2.45	2	1
1:A:55:ASN:HB3	1:A:130:THR:CG2	0.45	2.41	9	2
1:A:61:CYS:SG	1:A:124:LEU:HD12	0.45	2.52	9	1
1:A:88:LEU:HA	1:A:150:ALA:HB2	0.44	1.90	16	3
1:A:21:CYS:SG	1:A:58:LEU:HB3	0.44	2.52	10	1
1:A:157:GLN:HE21	1:A:159:GLN:NE2	0.44	2.10	9	2
1:A:56:ILE:O	1:A:130:THR:HG23	0.43	2.12	5	1
1:A:20:ALA:HB3	1:A:158:TYR:CE1	0.43	2.48	17	2
1:A:125:ASN:H	1:A:129:ASN:ND2	0.43	2.11	13	1
1:A:66:ALA:HB3	1:A:158:TYR:HB3	0.43	1.90	5	1
1:A:111:LEU:H	1:A:111:LEU:HD22	0.43	1.73	19	1
1:A:61:CYS:SG	1:A:124:LEU:HD13	0.43	2.54	17	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:101:VAL:HG11	1:A:177:PHE:CG	0.43	2.49	10	2
1:A:47:ALA:HB3	1:A:139:ALA:HB3	0.43	1.90	5	2
1:A:54:PHE:CD1	1:A:135:ALA:HB2	0.42	2.49	11	1
1:A:154:PHE:CD1	1:A:175:VAL:HG23	0.42	2.49	18	1
1:A:111:LEU:HD22	1:A:111:LEU:N	0.41	2.30	19	1
1:A:34:LEU:N	1:A:34:LEU:HD22	0.41	2.30	9	1
1:A:62:ASP:HB3	1:A:65:VAL:HG23	0.41	1.90	4	1
1:A:104:LEU:HD21	1:A:136:ARG:NH1	0.41	2.31	3	1
1:A:125:ASN:H	1:A:129:ASN:HD21	0.41	1.59	2	1
1:A:20:ALA:HB3	1:A:158:TYR:CD1	0.41	2.51	12	1
1:A:66:ALA:HB1	1:A:159:GLN:O	0.41	2.16	5	1
1:A:73:PHE:CD1	1:A:103:ILE:HG21	0.40	2.50	10	1
1:A:103:ILE:HG22	1:A:111:LEU:HD23	0.40	1.92	10	1
1:A:25:ALA:HA	1:A:28:VAL:HG12	0.40	1.92	7	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/184 (84%)	131±4 (85±2%)	19±3 (12±2%)	4±1 (3±1%)	10	46
All	All	3080/3680 (84%)	2620 (85%)	381 (12%)	79 (3%)	10	46

All 21 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	23	VAL	15
1	A	51	ALA	12
1	A	50	SER	8
1	A	117	THR	7
1	A	142	ALA	4
1	A	106	ARG	4
1	A	105	ASP	4
1	A	120	SER	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	20	ALA	3
1	A	126	ASN	2
1	A	119	SER	2
1	A	172	GLY	2
1	A	88	LEU	2
1	A	110	ALA	2
1	A	121	GLU	2
1	A	95	SER	2
1	A	73	PHE	1
1	A	71	VAL	1
1	A	55	ASN	1
1	A	28	VAL	1
1	A	60	ASP	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/125 (88%)	99±3 (90±3%)	11±3 (10±3%)	12	56
All	All	2200/2500 (88%)	1971 (90%)	229 (10%)	12	56

All 62 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	120	SER	13
1	A	86	LEU	12
1	A	183	ASN	12
1	A	170	VAL	10
1	A	114	ASP	10
1	A	71	VAL	10
1	A	125	ASN	8
1	A	99	VAL	8
1	A	129	ASN	8
1	A	52	VAL	8
1	A	30	GLN	7
1	A	38	ARG	6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	171	ASN	5
1	A	124	LEU	5
1	A	134	GLN	5
1	A	111	LEU	5
1	A	97	THR	4
1	A	55	ASN	4
1	A	180	GLU	4
1	A	79	ASP	4
1	A	156	VAL	4
1	A	57	GLN	4
1	A	104	LEU	3
1	A	126	ASN	3
1	A	50	SER	3
1	A	101	VAL	3
1	A	44	GLN	3
1	A	28	VAL	3
1	A	118	PHE	3
1	A	67	SER	3
1	A	21	CYS	3
1	A	151	ASP	3
1	A	169	THR	3
1	A	74	LEU	2
1	A	119	SER	2
1	A	95	SER	2
1	A	61	CYS	2
1	A	62	ASP	2
1	A	107	THR	2
1	A	24	ASP	2
1	A	128	THR	2
1	A	60	ASP	2
1	A	153	THR	2
1	A	31	THR	2
1	A	58	LEU	1
1	A	84	ASN	1
1	A	41	SER	1
1	A	105	ASP	1
1	A	29	ASP	1
1	A	76	THR	1
1	A	136	ARG	1
1	A	121	GLU	1
1	A	64	ASN	1
1	A	65	VAL	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	59	ASN	1
1	A	89	GLN	1
1	A	106	ARG	1
1	A	130	THR	1
1	A	27	SER	1
1	A	48	THR	1
1	A	102	GLN	1
1	A	113	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 15423

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1932
Number of shifts mapped to atoms	1932
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	6

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	182	0.01 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	157	-0.32 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}'$	176	0.16 ± 0.12	None needed (< 0.5 ppm)
^{15}N	180	-0.16 ± 0.32	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 1417 atoms were assigned a chemical shift out of a possible 1624. 25 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	766/771 (99%)	308/308 (100%)	305/310 (98%)	153/153 (100%)
Sidechain	623/758 (82%)	357/427 (84%)	242/298 (81%)	24/33 (73%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	28/95 (29%)	9/51 (18%)	19/40 (48%)	0/4 (0%)
Overall	1417/1624 (87%)	674/786 (86%)	566/648 (87%)	177/190 (93%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 86%, i.e. 1624 atoms were assigned a chemical shift out of a possible 1878. 27 out of 29 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	898/916 (98%)	360/366 (98%)	358/368 (97%)	180/182 (99%)
Sidechain	691/850 (81%)	395/478 (83%)	270/336 (80%)	26/36 (72%)
Aromatic	35/112 (31%)	12/60 (20%)	23/46 (50%)	0/6 (0%)
Overall	1624/1878 (86%)	767/904 (85%)	651/750 (87%)	206/224 (92%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	38	ARG	NE	114.70	92.63 – 76.73	18.9
1	A	106	ARG	NE	114.40	92.63 – 76.73	18.7
1	A	136	ARG	NE	113.30	92.63 – 76.73	18.0
1	A	72	ALA	HB3	-0.16	2.61 – 0.11	-6.1
1	A	72	ALA	HB1	-0.16	2.61 – 0.11	-6.1
1	A	72	ALA	HB2	-0.16	2.61 – 0.11	-6.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

