



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:32 am GMT

PDB ID : 3JUV
Title : Crystal structure of human lanosterol 14alpha-demethylase (CYP51)
Authors : Strushkevich, N.; MacKenzie, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Usanov, S.A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2009-09-15
Resolution : 3.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1000 (3.14-3.10)
Clashscore	112137	1099 (3.14-3.10)
Ramachandran outliers	110173	1060 (3.14-3.10)
Sidechain outliers	110143	1060 (3.14-3.10)
RSRZ outliers	101464	1005 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>21%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lanosterol 14-alpha demethylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	3605	2320	614	655	16	0	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	49	MET	-	EXPRESSION TAG	UNP Q16850
A	50	ALA	-	EXPRESSION TAG	UNP Q16850
A	51	LYS	-	EXPRESSION TAG	UNP Q16850
A	52	LYS	-	EXPRESSION TAG	UNP Q16850
A	53	THR	-	EXPRESSION TAG	UNP Q16850
A	503	THR	-	EXPRESSION TAG	UNP Q16850
A	504	HIS	-	EXPRESSION TAG	UNP Q16850
A	505	HIS	-	EXPRESSION TAG	UNP Q16850
A	506	HIS	-	EXPRESSION TAG	UNP Q16850
A	507	HIS	-	EXPRESSION TAG	UNP Q16850
A	508	HIS	-	EXPRESSION TAG	UNP Q16850
A	509	HIS	-	EXPRESSION TAG	UNP Q16850

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SUGAR (BETA-CYCLODEXTRIN) (three-letter code: BCD) (formula: $C_{42}H_{70}O_{35}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			77	42	35		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	O	0	0
			11	11		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lanosterol 14-alpha demethylase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	174.14Å 174.14Å 244.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.67 – 3.12 24.67 – 3.12	Depositor EDS
% Data completeness (in resolution range)	99.5 (24.67-3.12) 99.5 (24.67-3.12)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.05 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.198 , 0.227 0.198 , 0.227	Depositor DCC
R_{free} test set	1300 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	80.7	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3736	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCD, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3699	0.61	1/5011 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	252	ARG	NE-CZ-NH2	-5.37	117.62	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3605	0	3597	82	0
2	A	43	0	30	1	0
3	A	77	0	70	8	0
4	A	11	0	0	0	0
All	All	3736	0	3697	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:603:BCD:HO24	3:A:603:BCD:HO33	0.98	0.96
1:A:334:GLN:HE22	1:A:472:PHE:H	1.10	0.91
1:A:116:LEU:HD22	1:A:390:VAL:HG22	1.51	0.90
1:A:300:GLU:O	1:A:304:MET:HG2	1.76	0.85
1:A:353:THR:H	1:A:356:GLN:NE2	1.72	0.85
1:A:290:TYR:HB2	1:A:292:ASP:HB2	1.62	0.81
1:A:292:ASP:HB3	1:A:294:ARG:H	1.45	0.79
1:A:139:PHE:CD2	1:A:307:GLY:HA3	2.17	0.79
1:A:418:ARG:HG3	1:A:419:LEU:HG	1.67	0.77
1:A:289:THR:HG22	1:A:295:PRO:HA	1.69	0.73
1:A:369:GLU:OE1	1:A:372:ARG:NH1	2.22	0.72
3:A:603:BCD:H651	3:A:603:BCD:O66	1.89	0.72
1:A:471:GLU:OE1	1:A:499:LYS:HE2	1.94	0.68
1:A:328:ALA:HB1	1:A:473:ASP:HA	1.75	0.68
1:A:200:GLU:O	1:A:203:ILE:HG22	1.94	0.67
1:A:144:ALA:HA	1:A:152:PHE:CD2	2.30	0.67
1:A:203:ILE:O	1:A:207:SER:HB2	1.95	0.67
1:A:213:LYS:NZ	1:A:278:ILE:HD12	2.11	0.65
1:A:456:TYR:O	1:A:460:LYS:HB2	1.97	0.64
3:A:603:BCD:C65	3:A:603:BCD:O66	2.46	0.63
1:A:116:LEU:HD22	1:A:390:VAL:CG2	2.28	0.62
1:A:372:ARG:HG2	1:A:409:GLN:HB3	1.82	0.62
1:A:372:ARG:HD2	1:A:415:TRP:CZ3	2.35	0.62
1:A:68:ILE:HD12	1:A:68:ILE:N	2.16	0.61
1:A:484:TYR:HD2	1:A:489:HIS:CE1	2.18	0.61
1:A:352:LEU:HA	1:A:356:GLN:HE22	1.67	0.60
1:A:226:LEU:HD21	1:A:259:GLU:HG2	1.81	0.60
1:A:381:MET:HG3	1:A:402:CYS:SG	2.42	0.59
1:A:426:TYR:HA	1:A:430:ASN:OD1	2.02	0.59
1:A:242:PRO:HD2	1:A:245:LEU:HD11	1.83	0.58
1:A:338:TYR:O	1:A:341:GLN:HB2	2.02	0.58
1:A:334:GLN:HE21	1:A:501:ARG:HH12	1.53	0.57
1:A:208:HIS:CA	1:A:216:ARG:HG3	2.35	0.56
1:A:164:ASN:ND2	1:A:166:ALA:HB3	2.20	0.56
1:A:159:LEU:HD11	1:A:304:MET:HB2	1.87	0.56
1:A:422:ASN:HD21	1:A:424:ASP:HB2	1.73	0.54
1:A:204:LEU:HD23	1:A:219:LEU:HD23	1.90	0.54
1:A:213:LYS:HZ3	1:A:278:ILE:HD12	1.71	0.53
3:A:603:BCD:O33	3:A:603:BCD:O24	2.07	0.53
3:A:603:BCD:O64	3:A:603:BCD:H15	2.09	0.53
1:A:483:ASN:HD21	1:A:485:THR:HB	1.74	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:GLU:HG3	1:A:501:ARG:NH2	2.23	0.53
1:A:372:ARG:HA	1:A:409:GLN:NE2	2.24	0.52
1:A:121:LYS:HG3	1:A:446:ARG:HH22	1.74	0.52
1:A:484:TYR:CD2	1:A:489:HIS:CE1	2.98	0.52
1:A:422:ASN:O	1:A:425:ARG:HG2	2.11	0.51
1:A:430:ASN:ND2	1:A:432:ALA:HB3	2.25	0.51
1:A:369:GLU:CD	1:A:372:ARG:NH1	2.65	0.50
1:A:337:CYS:HB3	1:A:362:LEU:HD23	1.94	0.49
1:A:289:THR:HG22	1:A:295:PRO:CA	2.42	0.48
1:A:353:THR:H	1:A:356:GLN:HE22	1.58	0.48
1:A:315:THR:HB	2:A:601:HEM:HBB2	1.95	0.48
1:A:160:LYS:HG2	1:A:450:ILE:HB	1.96	0.48
1:A:369:GLU:CD	1:A:372:ARG:HH12	2.16	0.48
1:A:463:TRP:O	1:A:467:LEU:HB2	2.12	0.48
1:A:334:GLN:NE2	1:A:472:PHE:H	1.94	0.47
1:A:483:ASN:ND2	1:A:485:THR:H	2.13	0.47
1:A:191:GLU:CG	1:A:497:ARG:HG3	2.44	0.47
3:A:603:BCD:C13	3:A:603:BCD:H621	2.46	0.46
1:A:165:ILE:O	1:A:169:LYS:HG3	2.16	0.45
1:A:194:VAL:HG13	1:A:496:ILE:HG21	1.99	0.45
1:A:390:VAL:O	1:A:392:GLY:N	2.43	0.45
1:A:213:LYS:HZ1	1:A:278:ILE:HD12	1.80	0.45
1:A:369:GLU:HG3	1:A:421:PHE:CE1	2.52	0.45
1:A:121:LYS:HG3	1:A:446:ARG:NH2	2.32	0.45
3:A:603:BCD:O21	3:A:603:BCD:O37	2.29	0.45
1:A:230:LEU:HD12	1:A:230:LEU:HA	1.71	0.45
1:A:164:ASN:ND2	1:A:167:HIS:H	2.15	0.45
1:A:165:ILE:HG13	1:A:165:ILE:H	1.58	0.44
1:A:372:ARG:HG3	1:A:439:TYR:CD1	2.52	0.44
1:A:214:GLU:O	1:A:217:SER:OG	2.35	0.44
1:A:369:GLU:HG3	1:A:421:PHE:CD1	2.52	0.44
1:A:149:ASN:N	1:A:150:PRO:HD2	2.34	0.42
1:A:356:GLN:HB2	1:A:356:GLN:HE21	1.66	0.42
1:A:101:VAL:HG11	1:A:242:PRO:HD3	2.01	0.42
1:A:139:PHE:HB3	1:A:143:VAL:HG22	2.01	0.42
3:A:603:BCD:O64	3:A:603:BCD:C15	2.67	0.42
1:A:124:ASP:HB3	1:A:388:GLN:NE2	2.34	0.42
1:A:120:SER:HB3	1:A:125:LEU:HD12	2.01	0.42
1:A:381:MET:O	1:A:382:ARG:HD3	2.20	0.42
1:A:417:GLU:O	1:A:420:ASP:HB2	2.19	0.42
1:A:336:LYS:HE3	1:A:336:LYS:HB3	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:THR:N	1:A:356:GLN:NE2	2.53	0.41
1:A:66:SER:HB2	1:A:73:HIS:CE1	2.56	0.41
1:A:164:ASN:HD21	1:A:166:ALA:HB3	1.83	0.41
1:A:372:ARG:HD2	1:A:415:TRP:CH2	2.56	0.41
1:A:446:ARG:H	1:A:446:ARG:HG2	1.59	0.41
1:A:208:HIS:N	1:A:216:ARG:HG3	2.36	0.40
1:A:390:VAL:HB	1:A:391:ALA:H	1.75	0.40
1:A:194:VAL:HG13	1:A:496:ILE:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/461 (97%)	409 (92%)	32 (7%)	5 (1%)	17	53

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	144	ALA
1	A	273	GLN
1	A	347	GLU
1	A	428	GLN
1	A	390	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/403 (97%)	353 (90%)	38 (10%)	9	36

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	LYS
1	A	60	SER
1	A	123	GLU
1	A	143	VAL
1	A	160	LYS
1	A	164	ASN
1	A	180	LYS
1	A	194	VAL
1	A	207	SER
1	A	213	LYS
1	A	216	ARG
1	A	221	GLU
1	A	225	GLN
1	A	245	LEU
1	A	249	SER
1	A	271	ARG
1	A	274	SER
1	A	292	ASP
1	A	297	THR
1	A	313	GLN
1	A	317	SER
1	A	333	LEU
1	A	345	CYS
1	A	352	LEU
1	A	353	THR
1	A	381	MET
1	A	389	THR
1	A	390	VAL
1	A	418	ARG
1	A	420	ASP
1	A	422	ASN
1	A	430	ASN
1	A	446	ARG
1	A	473	ASP
1	A	483	ASN
1	A	485	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	487	MET
1	A	501	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	164	ASN
1	A	334	GLN
1	A	356	GLN
1	A	400	GLN
1	A	422	ASN
1	A	430	ASN
1	A	483	ASN
1	A	489	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	601	1	28,50,50	2.15	6 (21%)	17,82,82	1.92	3 (17%)
3	BCD	A	603	-	84,84,84	0.62	0	126,126,126	1.54	24 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	601	1	-	0/6/54/54	0/0/8/8
3	BCD	A	603	-	-	0/42/182/182	0/0/8/8

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	C3B-C2B	-5.10	1.33	1.40
2	A	601	HEM	C3C-C2C	-4.59	1.34	1.40
2	A	601	HEM	C1B-NB	2.04	1.39	1.36
2	A	601	HEM	C3B-CAB	3.41	1.54	1.47
2	A	601	HEM	C3C-CAC	3.79	1.55	1.47
2	A	601	HEM	C3D-C2D	4.77	1.51	1.37

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HEM	CBD-CAD-C3D	-5.17	102.60	112.47
2	A	601	HEM	C1D-C2D-C3D	-4.49	103.87	107.00
3	A	603	BCD	C13-O42-C42	-3.77	108.80	118.00
3	A	603	BCD	C16-O56-C56	-2.84	108.36	113.72
3	A	603	BCD	C15-O44-C44	-2.78	111.23	118.00
2	A	601	HEM	CAA-CBA-CGA	-2.70	108.04	112.66
3	A	603	BCD	C12-O41-C41	-2.61	111.63	118.00
3	A	603	BCD	O51-C11-C21	-2.27	105.92	110.30
3	A	603	BCD	C62-C52-C42	-2.14	107.40	113.24
3	A	603	BCD	O32-C32-C42	-2.07	105.16	109.87
3	A	603	BCD	C14-O43-C43	-2.06	112.97	118.00
3	A	603	BCD	C16-O45-C45	-2.01	113.09	118.00
3	A	603	BCD	O44-C44-C34	2.02	112.06	107.19
3	A	603	BCD	C37-C47-C57	2.05	115.23	110.88
3	A	603	BCD	O55-C55-C65	2.09	111.42	106.41
3	A	603	BCD	C21-C31-C41	2.57	114.94	109.61
3	A	603	BCD	C15-C25-C35	2.73	115.06	109.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	603	BCD	O52-C52-C62	2.82	113.16	106.41
3	A	603	BCD	C34-C44-C54	2.87	116.97	110.88
3	A	603	BCD	C17-O57-C57	2.89	119.16	113.72
3	A	603	BCD	O47-C11-C21	2.90	114.65	108.11
3	A	603	BCD	C24-C34-C44	3.03	115.89	109.61
3	A	603	BCD	C16-C26-C36	3.18	115.88	109.98
3	A	603	BCD	O54-C54-C44	3.34	116.59	109.75
3	A	603	BCD	C26-C36-C46	3.68	117.24	109.61
3	A	603	BCD	C17-C27-C37	3.99	117.39	109.98
3	A	603	BCD	C27-C37-C47	4.12	118.14	109.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	1	0
3	A	603	BCD	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	448/461 (97%)	-0.25	8 (1%) 69 48	53, 75, 97, 119	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	PRO	3.7
1	A	277	LYS	3.3
1	A	502	SER	2.9
1	A	275	GLN	2.3
1	A	273	GLN	2.2
1	A	56	ALA	2.1
1	A	293	GLY	2.1
1	A	274	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	BCD	A	603	77/77	0.84	0.25	0.85	64,70,72,75	77
2	HEM	A	601	43/43	0.99	0.15	-0.61	53,59,67,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.