



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:02 am GMT

PDB ID : 4JV1
Title : Ternary complex of gamma-OHPDG adduct modified dna with dna (-1 primer)
polymerase iv and incoming dgtp
Authors : Banerjee, S.; Shanmugam, G.; Stone, M.P.
Deposited on : 2013-03-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	<div> <div></div> <div>81%</div> <div>14%</div> <div>..</div> </div>
2	B	15	<div> <div>60%</div> <div>40%</div> </div>
3	C	13	<div> <div>77%</div> <div>23%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CA	A	402	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	0	0
			2744	1760	472	505	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP Q97W02
A	-4	HIS	-	EXPRESSION TAG	UNP Q97W02
A	-3	HIS	-	EXPRESSION TAG	UNP Q97W02
A	-2	HIS	-	EXPRESSION TAG	UNP Q97W02
A	-1	HIS	-	EXPRESSION TAG	UNP Q97W02
A	0	HIS	-	EXPRESSION TAG	UNP Q97W02


- Molecule 2 is a DNA chain called DNA (5'-D(P*CP*(HN1)P*GP*AP*AP*TP*CP*CP*TP*TP*CP*CP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	P	0	0	0
			302	145	50	92	15			

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*TP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	P	0	0	0
			273	129	57	75	12			

- Molecule 4 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: DGT) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		



- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Ca	0	0
			3	3		
5	C	1	Total	Ca	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	59	Total	O	0	0
			59	59		
6	B	12	Total	O	0	0
			12	12		
6	C	13	Total	O	0	0
			13	13		

- Molecule 1: DNA polymerase IV



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	93.27Å 102.18Å 52.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.87 – 2.30 41.87 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.8 (41.87-2.30) 95.2 (41.87-2.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.29Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7.2_869)	Depositor
R, R_{free}	0.213 , 0.269 0.204 , 0.261	Depositor DCC
R_{free} test set	1985 reflections (8.68%)	DCC
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3438	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HN1, CA, DGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2783	0.55	0/3736
2	B	0.75	0/305	1.49	6/463 (1.3%)
3	C	0.78	0/308	1.30	3/476 (0.6%)
All	All	0.52	0/3396	0.80	9/4675 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	617	DC	O4'-C1'-N1	9.57	114.70	108.00
2	B	612	DT	N3-C4-O4	6.54	123.83	119.90
2	B	607	DA	C1'-O4'-C4'	-6.12	103.97	110.10
3	C	507	DA	O4'-C1'-N9	-5.85	103.91	108.00
2	B	612	DT	C5-C4-O4	-5.71	120.91	124.90
2	B	615	DC	C1'-O4'-C4'	-5.47	104.63	110.10
2	B	607	DA	O4'-C1'-N9	5.32	111.72	108.00
3	C	504	DG	O4'-C1'-N9	5.22	111.66	108.00
3	C	513	DC	O4'-C1'-N1	5.06	111.54	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2889	24	0
2	B	302	0	171	5	0
3	C	273	0	147	0	0
4	A	31	0	12	2	0
5	A	3	0	0	0	0
5	C	1	0	0	0	0
6	A	59	0	0	1	1
6	B	12	0	0	1	0
6	C	13	0	0	0	0
All	All	3438	0	3219	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:605:HN1:N9	2:B:605:HN1:C1'	1.73	1.47
2:B:605:HN1:C4	2:B:605:HN1:C1'	2.54	0.84
1:A:100:GLU:OE1	1:A:240:ARG:NH2	2.17	0.73
1:A:242:ARG:HH11	1:A:242:ARG:HA	1.53	0.71
1:A:256:ARG:NH2	1:A:323:LEU:O	2.30	0.65
2:B:605:HN1:C2'	2:B:605:HN1:N9	2.57	0.64
1:A:270:GLU:OE2	1:A:312:TYR:OH	2.17	0.62
1:A:51:ARG:NH1	4:A:401:DGT:O1G	2.32	0.62
1:A:230:ARG:HB3	1:A:232:GLU:HG2	1.82	0.61
1:A:180:ILE:HD13	1:A:194:LEU:HD13	1.87	0.57
1:A:115:VAL:HG22	1:A:120:GLU:HB3	1.86	0.57
1:A:63:GLU:HA	1:A:66:LYS:HD2	1.86	0.57
1:A:36:ARG:NH1	1:A:254:ASN:OD1	2.38	0.56
1:A:285:HIS:HD2	6:A:514:HOH:O	1.89	0.56
1:A:79:GLU:H	1:A:79:GLU:CD	2.10	0.55
1:A:214:LYS:HE2	1:A:219:GLU:HB2	1.88	0.55
1:A:292:ASP:N	1:A:292:ASP:OD1	2.38	0.55
1:A:12:TYR:CD2	4:A:401:DGT:H2'	2.44	0.53
1:A:166:ILE:HG23	1:A:170:GLU:HB3	1.94	0.49
1:A:66:LYS:HB3	1:A:66:LYS:HE3	1.61	0.47
1:A:255:SER:OG	1:A:256:ARG:N	2.49	0.45
1:A:292:ASP:OD2	1:A:328:ARG:NH1	2.45	0.45
1:A:251:MET:HG2	1:A:264:TYR:CD2	2.52	0.44
1:A:41:GLY:HA2	2:B:604:DC:H5'	1.99	0.44
1:A:95:TYR:OH	1:A:131:LYS:HE3	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:GLN:OE1	1:A:138:ILE:HA	2.19	0.41
1:A:95:TYR:CD1	1:A:124:LEU:HD11	2.56	0.41
2:B:604:DC:OP1	6:B:705:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:513:HOH:O	6:A:542:HOH:O[1_556]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/347 (98%)	330 (97%)	7 (2%)	2 (1%)	28	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	ASP
1	A	277	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	300/306 (98%)	270 (90%)	30 (10%)	9 10

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	24	LYS
1	A	38	GLU
1	A	51	ARG
1	A	63	GLU
1	A	65	LYS
1	A	66	LYS
1	A	94	GLU
1	A	115	VAL
1	A	116	ARG
1	A	119	ARG
1	A	138	ILE
1	A	169	GLU
1	A	176	ARG
1	A	177	GLU
1	A	178	LEU
1	A	189	ILE
1	A	196	LYS
1	A	201	LYS
1	A	238	ARG
1	A	241	VAL
1	A	242	ARG
1	A	256	ARG
1	A	260	GLU
1	A	270	GLU
1	A	292	ASP
1	A	309	GLU
1	A	325	GLU
1	A	326	ASP
1	A	327	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HN1	B	605	2	19,29,37	5.68	6 (31%)	19,43,55	3.15	8 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HN1	B	605	2	-	0/3/31/43	0/3/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	605	HN1	O5'-C5'	-22.48	1.13	1.44
2	B	605	HN1	C3-N2	-2.01	1.42	1.47
2	B	605	HN1	C2-N2	2.48	1.40	1.32
2	B	605	HN1	C6-N1	4.18	1.44	1.38
2	B	605	HN1	C4-N3	4.98	1.40	1.33
2	B	605	HN1	O6-C6	6.72	1.41	1.24

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	605	HN1	C2'-C1'-N9	-4.45	103.71	114.23
2	B	605	HN1	C4-C5-N7	-2.65	106.84	109.41
2	B	605	HN1	C3-N2-C2	-2.46	112.34	117.41
2	B	605	HN1	C4'-O4'-C1'	-2.40	103.54	109.42
2	B	605	HN1	O4'-C1'-N9	-2.30	103.91	107.78
2	B	605	HN1	C-C3-N2	2.77	121.69	113.49
2	B	605	HN1	O11-C1-N1	6.33	124.70	109.66
2	B	605	HN1	O5'-C5'-C4'	9.32	141.81	109.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	605	HN1	3	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	DGT	A	401	5	26,33,33	1.91	5 (19%)	28,52,52	2.17	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DGT	A	401	5	-	0/18/34/34	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	401	DGT	C5'-C4'	-2.74	1.42	1.51
4	A	401	DGT	O3'-C3'	-2.71	1.37	1.43
4	A	401	DGT	C2'-C3'	-2.29	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	401	DGT	C2-N2	3.90	1.42	1.34
4	A	401	DGT	O6-C6	6.92	1.42	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	DGT	N3-C2-N1	-4.25	121.26	127.46
4	A	401	DGT	C4-C5-N7	-3.39	106.14	109.41
4	A	401	DGT	C2'-C1'-N9	-3.38	106.24	114.23
4	A	401	DGT	C5-C6-N1	-3.21	118.91	123.48
4	A	401	DGT	C6-C5-C4	-3.07	117.80	120.84
4	A	401	DGT	O5'-C5'-C4'	3.44	121.20	109.00
4	A	401	DGT	C2-N3-C4	3.61	119.37	115.16
4	A	401	DGT	C6-N1-C2	4.13	122.00	116.06
4	A	401	DGT	O4'-C1'-N9	4.53	115.41	107.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	401	DGT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/347 (98%)	-0.02	3 (0%) 84 87	36, 50, 69, 82	8 (2%)
2	B	14/15 (93%)	-0.58	0 100 100	49, 60, 66, 68	0
3	C	13/13 (100%)	-0.53	0 100 100	44, 53, 69, 72	0
All	All	368/375 (98%)	-0.06	3 (0%) 86 89	36, 51, 69, 82	8 (2%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	116	ARG	3.5
1	A	134	GLU	2.5
1	A	34	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	HN1	B	605	26/34	0.93	0.12	-	54,57,62,69	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CA	A	402	1/1	0.97	0.19	2.85	50,50,50,50	0
5	CA	A	403	1/1	0.91	0.19	1.58	66,66,66,66	0
4	DGT	A	401	31/31	0.97	0.12	-0.23	42,48,56,64	5
5	CA	A	404	1/1	0.50	0.29	-	80,80,80,80	0
5	CA	C	601	1/1	0.85	0.11	-	97,97,97,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.