



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 07:55 am GMT

PDB ID : 4K5M  
Title : Phosphonic Arginine Mimetics as Inhibitors of the M1 Aminopeptidases from Plasmodium falciparum  
Authors : McGowan, S.  
Deposited on : 2013-04-14  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1762 (1.76-1.76)
Clashscore	112137	1889 (1.76-1.76)
Ramachandran outliers	110173	1868 (1.76-1.76)
Sidechain outliers	110143	1868 (1.76-1.76)
RSRZ outliers	101464	1770 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	898	 89% 10% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	1102	-	-	-	X
4	GOL	A	1104	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1105	-	-	-	X
4	GOL	A	1107	-	-	-	X
5	1OV	A	1106[A]	-	-	-	X
5	1OV	A	1106[B]	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M1 family aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	889	Total	C	N	O	S	7	12	0
			7272	4683	1178	1383	28			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	194	MET	-	EXPRESSION TAG	UNP O96935
A	213	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	223	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	378	PRO	HIS	ENGINEERED MUTATION	UNP O96935
A	501	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	745	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	795	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	1069	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	1086	HIS	-	EXPRESSION TAG	UNP O96935
A	1087	HIS	-	EXPRESSION TAG	UNP O96935
A	1088	HIS	-	EXPRESSION TAG	UNP O96935
A	1089	HIS	-	EXPRESSION TAG	UNP O96935
A	1090	HIS	-	EXPRESSION TAG	UNP O96935
A	1091	HIS	-	EXPRESSION TAG	UNP O96935


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is [(1R)-1-AMINO-4-CARBAMIMIDAMIDOBUTYL]PHOSPHONIC ACID (three-letter code: 1OV) (formula:  $C_5H_{15}N_4O_3P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	1
			18	7	7	3	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1147	Total	O	0	0
			1147	1147		



- Molecule 1: M1 family aminopeptidase

HIS	P798	T492	MET
HIS	V799	G496	GLU
HIS	K808	H496	P196
HIS	L811	E497	H199
HIS	R831	Y502	K252
	I834	E526	I256
	N835	K535	S257
	L840	H543	Y264
	T848	L546	E265
	M867	L553	G266
	E872	S867	E267
	D890	S570	E268
	R898	F574	Y269
	E919	K808	S286
	S934	D634	S291
	K951	L643	E348
	R969	Y658	P351
	D989	K665	H370
	R997	N672	D391
	P1002	Q673	L392
	R1008	Y674	K393
	R1009	E679	T398
	L1045	F688	Y403
	W1046	F721	K407
	T1051	V726	E416
	K1052	F736	S432
	R1053	F736	D436
	Q1054	D748	L442
	N1059	K774	E443
	N1073	M778	L449
	E1076	N779	A453
	R1080	Y780	V459
	N1083	N781	L475
	K1084	K785	K479
	F1091	T792	K480
			F485

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.00Å 109.03Å 118.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.26 – 1.75 80.26 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.8 (80.26-1.75) 98.8 (80.26-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.163 , 0.207 0.163 , 0.209	Depositor DCC
$R_{free}$ test set	4938 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.4	Xtriage
Anisotropy	0.464	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8458	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1OV, ZN, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.17	6/7471 (0.1%)	0.96	14/10116 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1076	GLU	CB-CG	7.77	1.67	1.52
1	A	934	SER	CB-OG	-5.74	1.34	1.42
1	A	658	TYR	CD2-CE2	5.55	1.47	1.39
1	A	736	PHE	CE2-CZ	5.32	1.47	1.37
1	A	799	VAL	CB-CG1	5.16	1.63	1.52
1	A	721	PHE	CE1-CZ	5.03	1.47	1.37

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	778[A]	MET	CG-SD-CE	7.51	112.22	100.20
1	A	778[B]	MET	CG-SD-CE	7.51	112.22	100.20
1	A	546	LEU	CB-CG-CD2	6.98	122.86	111.00
1	A	898	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	A	748	ASP	CB-CG-OD1	5.94	123.64	118.30
1	A	449	LEU	CB-CG-CD2	5.86	120.95	111.00
1	A	898	ARG	NE-CZ-NH1	5.79	123.19	120.30
1	A	1008	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	997	ARG	NE-CZ-NH2	5.49	123.04	120.30
1	A	748	ASP	CB-CG-OD2	-5.40	113.44	118.30
1	A	643	LEU	CB-CG-CD2	-5.24	102.09	111.00
1	A	890	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	1009	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	A	774	LYS	CD-CE-NZ	-5.12	99.92	111.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7272	0	7141	66	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	18	0	24	1	0
5	A	18	0	14	1	0
6	A	1147	0	0	40	1
All	All	8458	0	7179	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459:VAL:N	6:A:2337:HOH:O	1.83	1.04
1:A:665:LYS:CD	6:A:1914:HOH:O	2.05	1.02
1:A:834:ILE:HB	6:A:1667:HOH:O	1.60	1.00
1:A:781[A]:ASN:ND2	6:A:2297:HOH:O	1.95	0.98
1:A:1080:ARG:NH1	6:A:1365:HOH:O	2.01	0.93
1:A:286:SER:HB2	6:A:1642:HOH:O	1.70	0.91
1:A:459:VAL:CA	6:A:2337:HOH:O	2.20	0.87
1:A:543[B]:HIS:CD2	6:A:1859:HOH:O	2.29	0.85
1:A:543[B]:HIS:HD2	6:A:1859:HOH:O	1.58	0.84
1:A:679:GLU:OE2	6:A:2343:HOH:O	1.99	0.81
1:A:1009:ARG:HD3	6:A:1811:HOH:O	1.81	0.81
1:A:492:THR:HG23	1:A:526[A]:GLU:OE2	1.85	0.76
1:A:264:VAL:HG22	1:A:268:GLU:HG3	1.69	0.74
1:A:608:LYS:CB	6:A:2219:HOH:O	2.35	0.72
1:A:1051:THR:HB	6:A:2338:HOH:O	1.92	0.68
1:A:1083:ASN:OD1	6:A:2021:HOH:O	2.12	0.68
1:A:256:ILE:HD11	1:A:269:TYR:CD1	2.29	0.67

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:872[B]:GLU:OE1	6:A:1744:HOH:O	2.13	0.67
1:A:1051:THR:CB	6:A:2338:HOH:O	2.41	0.66
1:A:674:TYR:HB3	6:A:1676:HOH:O	1.95	0.64
1:A:459:VAL:O	5:A:1106[B]:1OV:N2	2.29	0.64
1:A:252:LYS:HE3	6:A:2242:HOH:O	1.99	0.63
1:A:264:VAL:HG23	1:A:267:GLU:HB3	1.81	0.61
1:A:570:SER:OG	1:A:1073[A]:ASN:ND2	2.34	0.61
1:A:785:LYS:HD2	6:A:1743:HOH:O	2.00	0.61
1:A:989:ASP:CB	6:A:2169:HOH:O	2.51	0.58
1:A:403:TYR:N	1:A:443:GLU:OE2	2.33	0.57
1:A:672:ASN:ND2	6:A:1643:HOH:O	2.37	0.57
1:A:951:LYS:HE2	6:A:2215:HOH:O	2.05	0.57
1:A:969[A]:ARG:HD3	1:A:1002:PRO:HB3	1.87	0.56
1:A:1059[B]:ASN:ND2	6:A:1920:HOH:O	2.24	0.56
1:A:808:LYS:HE3	6:A:2237:HOH:O	2.06	0.55
1:A:872[A]:GLU:OE1	6:A:1800:HOH:O	2.18	0.54
1:A:264:VAL:CG2	1:A:267:GLU:HB3	2.38	0.54
1:A:432[A]:SER:OG	1:A:495:GLY:HA2	2.07	0.54
1:A:778[A]:MET:SD	1:A:797:THR:HG21	2.47	0.54
1:A:480:LYS:CD	6:A:2135:HOH:O	2.56	0.53
1:A:834:ILE:HD13	1:A:1045:LEU:HD22	1.90	0.53
1:A:252:LYS:HD2	6:A:1868:HOH:O	2.09	0.52
1:A:726:VAL:HG22	6:A:1691:HOH:O	2.08	0.52
1:A:835:ASN:ND2	6:A:2282:HOH:O	2.12	0.52
1:A:674:TYR:CB	6:A:1676:HOH:O	2.55	0.50
1:A:1046:TRP:O	1:A:1054:GLN:HG2	2.11	0.50
1:A:1084:LYS:HA	6:A:2096:HOH:O	2.12	0.49
1:A:781[B]:ASN:ND2	6:A:1743:HOH:O	2.45	0.49
1:A:831:ARG:NH2	6:A:1873:HOH:O	2.37	0.49
1:A:553:LEU:HD23	1:A:553:LEU:C	2.35	0.47
1:A:257:SER:HB2	1:A:291:SER:OG	2.15	0.47
1:A:199:HIS:O	1:A:567:SER:HA	2.15	0.47
1:A:196:PRO:HD2	6:A:1851:HOH:O	2.16	0.45
1:A:442:LEU:HB3	1:A:502:TYR:CE2	2.52	0.45
1:A:1076:GLU:HG2	1:A:1080:ARG:HE	1.81	0.45
1:A:535:LYS:NZ	6:A:1762:HOH:O	2.26	0.44
1:A:348:GLU:CB	6:A:1978:HOH:O	2.66	0.44
1:A:811:LEU:HB3	1:A:867:MET:SD	2.58	0.44
1:A:351:PRO:HG2	1:A:391:ASP:HA	1.99	0.44
1:A:407:LYS:HE3	6:A:1933:HOH:O	2.16	0.43
1:A:919:GLU:OE2	6:A:2345:HOH:O	2.21	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459:VAL:C	6:A:2337:HOH:O	2.55	0.42
1:A:969[B]:ARG:NE	6:A:2199:HOH:O	2.27	0.42
1:A:496:HIS:CE1	4:A:1107:GOL:H32	2.55	0.41
1:A:393:LYS:HG3	1:A:416:GLU:HG3	2.03	0.41
1:A:779:ASN:HB2	1:A:848:THR:HG23	2.03	0.41
1:A:453:ALA:HB2	1:A:475:LEU:HD23	2.02	0.40
1:A:570:SER:CB	1:A:1073[A]:ASN:HD22	2.34	0.40
1:A:398:THR:HG21	1:A:407:LYS:HD3	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2128:HOH:O	6:A:2289:HOH:O[2_554]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	899/898 (100%)	883 (98%)	15 (2%)	1 (0%)	55 35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	459	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	805/830 (97%)	791 (98%)	14 (2%)	66	47

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	GLU
1	A	370	HIS
1	A	432[A]	SER
1	A	432[B]	SER
1	A	436	ASP
1	A	449	LEU
1	A	479	LYS
1	A	497	GLU
1	A	546	LEU
1	A	634	ASP
1	A	688	PHE
1	A	840	LEU
1	A	1045	LEU
1	A	1052	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	ASN
1	A	653	HIS
1	A	882	ASN
1	A	915	ASN
1	A	1083	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	1104	-	5,5,5	0.65	0	5,5,5	1.06	0
4	GOL	A	1105	-	5,5,5	1.04	0	5,5,5	1.02	0
5	1OV	A	1106[A]	-	11,12,12	2.08	4 (36%)	9,16,16	2.31	4 (44%)
5	1OV	A	1106[B]	-	11,12,12	2.10	4 (36%)	9,16,16	2.31	4 (44%)
4	GOL	A	1107	-	5,5,5	0.75	0	5,5,5	1.80	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1104	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1105	-	-	0/4/4/4	0/0/0/0
5	1OV	A	1106[A]	-	-	1/12/13/13	0/0/0/0
5	1OV	A	1106[B]	-	-	0/12/13/13	0/0/0/0
4	GOL	A	1107	-	-	0/4/4/4	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1106[B]	1OV	P-C5	-3.19	1.82	1.84
5	A	1106[A]	1OV	P-C5	-3.19	1.82	1.84
5	A	1106[B]	1OV	P-O3	2.35	1.58	1.54
5	A	1106[A]	1OV	P-O3	2.35	1.58	1.54
5	A	1106[B]	1OV	P-O1	3.53	1.56	1.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1106[A]	1OV	P-O1	3.53	1.56	1.49
5	A	1106[B]	1OV	P-O2	3.61	1.60	1.54
5	A	1106[A]	1OV	P-O2	3.61	1.60	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1106[B]	1OV	O2-P-O1	-4.53	101.89	113.41
5	A	1106[A]	1OV	O2-P-O1	-4.53	101.89	113.41
5	A	1106[A]	1OV	N3-C1-N1	-2.43	116.22	120.58
5	A	1106[A]	1OV	C4-C3-C2	-2.10	105.84	112.11
5	A	1106[B]	1OV	C4-C3-C2	2.25	118.83	112.11
5	A	1106[B]	1OV	C3-C2-N3	2.59	119.64	112.18
4	A	1107	GOL	O2-C2-C1	3.10	123.47	108.84
5	A	1106[B]	1OV	O3-P-O2	3.53	117.68	107.69
5	A	1106[A]	1OV	O3-P-O2	3.53	117.68	107.69

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1106[A]	1OV	N2-C1-N3-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1106[B]	1OV	1	0
4	A	1107	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	889/898 (98%)	-0.27	3 (0%) 93 95	7, 17, 31, 50	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	674	TYR	2.5
1	A	485	PHE	2.3
1	A	574	PHE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	1104	6/6	0.91	0.16	8.77	25,26,35,36	0
4	GOL	A	1105	6/6	0.94	0.12	5.62	14,16,20,20	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	1102	1/1	0.98	0.21	5.31	33,33,33,33	0
4	GOL	A	1107	6/6	0.91	0.22	4.91	25,32,35,37	0
5	1OV	A	1106[A]	13/13	0.92	0.15	3.03	11,16,25,27	5
5	1OV	A	1106[B]	13/13	0.92	0.15	1.71	9,15,18,21	5
2	ZN	A	1101	1/1	1.00	0.07	-3.06	12,12,12,12	0
3	MG	A	1103	1/1	0.96	0.16	-	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.