



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 25, 2017 – 06:25 PM EDT

PDB ID : 3K6C  
Title : Crystal structure of protein ne0167 from nitrosomonas europaea  
Authors : Chang, C.; Evdokimova, E.; Savchenko, A.; Edwards, A.M.; Joachimiak, A.;  
Midwest Center for Structural Genomics (MCSG)  
Deposited on : unknown  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	95	<div> <div>3%</div> <div>66%</div> <div>27%</div> <div>• •</div> </div>
1	B	95	<div> <div>5%</div> <div>71%</div> <div>21%</div> <div>• 5%</div> </div>
1	C	95	<div> <div>7%</div> <div>65%</div> <div>26%</div> <div>• • •</div> </div>
1	D	95	<div> <div>4%</div> <div>71%</div> <div>26%</div> <div>•</div> </div>
1	E	95	<div> <div>5%</div> <div>64%</div> <div>29%</div> <div>• •</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	95	<div><div></div><div>5%</div><div>65%</div><div>29%</div><div></div><div></div></div>
1	G	95	<div><div></div><div>11%</div><div>65%</div><div>31%</div><div></div><div></div></div>
1	H	95	<div><div></div><div>5%</div><div>64%</div><div>26%</div><div></div><div></div></div>
1	I	95	<div><div></div><div>4%</div><div>65%</div><div>27%</div><div></div><div></div></div>
1	J	95	<div><div></div><div>5%</div><div>67%</div><div>25%</div><div></div><div>5%</div></div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7911 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein NE0167.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	B	90	Total	C	N	O	S	Se	0	0	0
			748	467	131	146	2	2			
1	C	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	D	92	Total	C	N	O	S	Se	0	0	0
			767	478	135	150	2	2			
1	E	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	F	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	G	91	Total	C	N	O	S	Se	0	0	0
			756	471	133	148	2	2			
1	H	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	I	91	Total	C	N	O	S	Se	0	0	0
			758	473	134	147	2	2			
1	J	90	Total	C	N	O	S	Se	0	0	0
			748	467	131	146	2	2			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	39	Total	O	0	0
			39	39		
2	B	45	Total	O	0	0
			45	45		
2	C	46	Total	O	0	0
			46	46		
2	D	37	Total	O	0	0
			37	37		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	29	Total 29	O 29	0	0
2	F	32	Total 32	O 32	0	0
2	G	30	Total 30	O 30	0	0
2	H	33	Total 33	O 33	0	0
2	I	23	Total 23	O 23	0	0
2	J	30	Total 30	O 30	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized protein NE0167




- Molecule 1: Uncharacterized protein NE0167




- Molecule 1: Uncharacterized protein NE0167



- Molecule 1: Uncharacterized protein NE0167




- Molecule 1: Uncharacterized protein NE0167




GLU


- Molecule 1: Uncharacterized protein NE0167




- Molecule 1: Uncharacterized protein NE0167




- Molecule 1: Uncharacterized protein NE0167



- Molecule 1: Uncharacterized protein NE0167



- Molecule 1: Uncharacterized protein NE0167



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.42Å 134.55Å 78.46Å 90.00° 96.24° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 45.32 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.20) 99.9 (45.32-2.20)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.86 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.221 , 0.261 0.260 , 0.289	Depositor DCC
$R_{free}$ test set	2567 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 29.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7911	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	6.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.79	0/770	0.80	1/1033 (0.1%)
1	B	0.85	1/759 (0.1%)	0.81	1/1018 (0.1%)
1	C	0.91	1/770 (0.1%)	0.87	3/1033 (0.3%)
1	D	0.88	0/779	0.90	2/1045 (0.2%)
1	E	0.83	0/770	0.84	4/1033 (0.4%)
1	F	0.86	1/770 (0.1%)	0.84	1/1033 (0.1%)
1	G	0.75	0/767	0.79	1/1029 (0.1%)
1	H	0.76	0/770	0.89	3/1033 (0.3%)
1	I	0.72	0/770	0.81	0/1033
1	J	0.91	1/759 (0.1%)	0.91	2/1018 (0.2%)
All	All	0.83	4/7684 (0.1%)	0.85	18/10308 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	H	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	89	ASN	C-O	6.27	1.35	1.23
1	J	74	CYS	CB-SG	-6.08	1.72	1.82
1	B	60	GLU	CD-OE2	5.42	1.31	1.25
1	C	74	CYS	CB-SG	-5.12	1.73	1.81

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	66	MSE	N-CA-C	-8.05	89.26	111.00
1	D	22	ARG	NE-CZ-NH2	-7.95	116.33	120.30
1	F	22	ARG	NE-CZ-NH2	-7.50	116.55	120.30
1	B	22	ARG	NE-CZ-NH2	-7.41	116.60	120.30
1	C	86	LEU	CB-CG-CD2	-7.31	98.57	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	93	ALA	Peptide
1	H	65	ALA	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	758	0	732	26	0
1	B	748	0	725	23	0
1	C	758	0	732	34	0
1	D	767	0	738	27	0
1	E	758	0	732	40	0
1	F	758	0	732	32	0
1	G	756	0	731	32	0
1	H	758	0	732	42	0
1	I	758	0	732	33	0
1	J	748	0	725	27	0
2	A	39	0	0	2	0
2	B	45	0	0	1	0
2	C	46	0	0	0	0
2	D	37	0	0	4	0
2	E	29	0	0	1	0
2	F	32	0	0	3	0
2	G	30	0	0	2	0
2	H	33	0	0	2	0
2	I	23	0	0	0	0
2	J	30	0	0	1	0
All	All	7911	0	7311	223	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:27:LEU:HD21	1:C:68:LEU:HD12	1.38	1.04
1:H:4:ASP:N	1:H:21:HIS:HE2	1.60	0.98
1:E:17:THR:OG1	1:F:42:ASN:ND2	2.01	0.93
1:H:19:ASP:OD1	1:H:22:ARG:NH2	2.00	0.93
1:F:69:GLU:OE1	1:F:72:ARG:HD3	1.69	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	89/95 (94%)	88 (99%)	1 (1%)	0	100	100
1	B	88/95 (93%)	88 (100%)	0	0	100	100
1	C	89/95 (94%)	85 (96%)	3 (3%)	1 (1%)	17	13
1	D	90/95 (95%)	86 (96%)	4 (4%)	0	100	100
1	E	89/95 (94%)	84 (94%)	5 (6%)	0	100	100
1	F	89/95 (94%)	89 (100%)	0	0	100	100
1	G	89/95 (94%)	85 (96%)	4 (4%)	0	100	100
1	H	89/95 (94%)	86 (97%)	2 (2%)	1 (1%)	17	13
1	I	89/95 (94%)	85 (96%)	3 (3%)	1 (1%)	17	13
1	J	88/95 (93%)	88 (100%)	0	0	100	100
All	All	889/950 (94%)	864 (97%)	22 (2%)	3 (0%)	44	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	66	MSE
1	I	10	THR
1	C	80	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/81 (100%)	78 (96%)	3 (4%)	39	49
1	B	80/81 (99%)	78 (98%)	2 (2%)	53	65
1	C	81/81 (100%)	77 (95%)	4 (5%)	29	35
1	D	82/81 (101%)	80 (98%)	2 (2%)	54	67
1	E	81/81 (100%)	80 (99%)	1 (1%)	75	86
1	F	81/81 (100%)	78 (96%)	3 (4%)	39	49
1	G	81/81 (100%)	79 (98%)	2 (2%)	53	65
1	H	81/81 (100%)	76 (94%)	5 (6%)	21	24
1	I	81/81 (100%)	78 (96%)	3 (4%)	39	49
1	J	80/81 (99%)	77 (96%)	3 (4%)	38	47
All	All	809/810 (100%)	781 (96%)	28 (4%)	41	51

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	4	ASP
1	G	38	ASN
1	J	38	ASN
1	F	46	ASP
1	F	94	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	38	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	42	ASN
1	H	42	ASN
1	E	42	ASN
1	G	42	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	89/95 (93%)	0.48	3 (3%) 46 43	2, 2, 13, 50	1 (1%)
1	B	88/95 (92%)	0.54	5 (5%) 24 24	2, 2, 10, 18	0
1	C	89/95 (93%)	0.62	7 (7%) 13 12	2, 2, 16, 56	0
1	D	90/95 (94%)	0.49	4 (4%) 35 33	2, 3, 11, 54	1 (1%)
1	E	89/95 (93%)	0.60	5 (5%) 25 24	2, 3, 15, 53	0
1	F	89/95 (93%)	0.66	5 (5%) 25 24	2, 6, 17, 25	0
1	G	89/95 (93%)	0.64	10 (11%) 6 5	2, 9, 20, 58	0
1	H	89/95 (93%)	0.48	5 (5%) 25 24	2, 7, 18, 22	0
1	I	89/95 (93%)	0.62	4 (4%) 34 32	2, 3, 18, 57	0
1	J	88/95 (92%)	0.48	5 (5%) 24 24	2, 4, 15, 22	0
All	All	889/950 (93%)	0.56	53 (5%) 23 22	2, 3, 18, 58	2 (0%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	94	HIS	8.3
1	A	94	HIS	7.4
1	E	78	PHE	5.2
1	G	93	ALA	4.8
1	H	65	ALA	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.