



wwPDB X-ray Structure Validation Summary Report i

Jul 18, 2017 – 09:41 AM EDT

PDB ID : 5KAH
Title : Crystal structure of a dioxygenase in the Crotonase superfamily in P21, V425T mutant
Authors : Li, K.; Fielding, E.N.; Condurso, H.L.; Bruner, S.D.
Deposited on : unknown
Resolution : 2.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3276 (2.80-2.76)
Clashscore	112137	3771 (2.80-2.76)
Ramachandran outliers	110173	3707 (2.80-2.76)
Sidechain outliers	110143	3709 (2.80-2.76)
RSRZ outliers	101464	3307 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	F	438	1%	81%	15%	..
1	G	438		83%	12%	..
1	H	438		83%	11%	• 5%
1	I	438		81%	13%	..
1	J	438	3%	84%	11%	..
1	K	438		85%	10%	..
1	L	438	2%	82%	12%	• 5%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 40568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	421	3272	2055	611	596	10	0	0	0
1	B	418	3251	2042	607	592	10	0	0	0
1	C	417	3244	2038	606	590	10	0	0	0
1	D	421	3273	2055	611	597	10	0	0	0
1	E	421	3272	2055	611	596	10	0	0	0
1	F	421	3272	2055	611	596	10	0	0	0
1	G	419	3256	2045	608	593	10	0	0	0
1	H	418	3251	2042	607	592	10	0	0	0
1	I	421	3272	2055	611	596	10	0	0	0
1	J	419	3253	2043	608	592	10	0	0	0
1	K	419	3260	2047	608	595	10	0	0	0
1	L	418	3246	2040	607	589	10	0	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	425	THR	VAL	engineered mutation	UNP Q8KLK7
B	425	THR	VAL	engineered mutation	UNP Q8KLK7
C	425	THR	VAL	engineered mutation	UNP Q8KLK7
D	425	THR	VAL	engineered mutation	UNP Q8KLK7
E	425	THR	VAL	engineered mutation	UNP Q8KLK7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	425	THR	VAL	engineered mutation	UNP Q8KLK7
G	425	THR	VAL	engineered mutation	UNP Q8KLK7
H	425	THR	VAL	engineered mutation	UNP Q8KLK7
I	425	THR	VAL	engineered mutation	UNP Q8KLK7
J	425	THR	VAL	engineered mutation	UNP Q8KLK7
K	425	THR	VAL	engineered mutation	UNP Q8KLK7
L	425	THR	VAL	engineered mutation	UNP Q8KLK7

- Molecule 2 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXY-3-(PHOSPHO NOOXY)TETRAHYDROFURAN-2-YL]METHYL (3R)-4-({3-[{2-[(3,5-DIHYDROXYPHENYL)ACETYL]AMINO}ETHYL]AMINO}-3-OXOPROPYL)AMINO)-3-HYDROXY-2,2-DIMETHYL-4-OXOBUTYL DIHYDROGEN DIPHOSPHATE (three-letter code: YE1) (formula: C₂₉H₄₃N₈O₁₉P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	B	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	C	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	D	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	E	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	F	1	Total	C	N	O	P	0	0
			59	29	8	19	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	G	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	H	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	I	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	J	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	K	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	L	1	Total	C	N	O	P	0	0
			59	29	8	19	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		
3	B	56	Total	O	0	0
			56	56		
3	C	62	Total	O	0	0
			62	62		
3	D	61	Total	O	0	0
			61	61		
3	E	74	Total	O	0	0
			74	74		
3	F	60	Total	O	0	0
			60	60		
3	G	65	Total	O	0	0
			65	65		
3	H	66	Total	O	0	0
			66	66		
3	I	53	Total	O	0	0
			53	53		
3	J	53	Total	O	0	0
			53	53		
3	K	67	Total	O	0	0
			67	67		
3	L	63	Total	O	0	0
			63	63		


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase






- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain E:



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain F:



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain G



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain H-





- D271 R272 R280 I281 F292 G296 Q299 L300 V303 F304 P318 ALA ALA LYS GLU G323 P326 G327 A328 A329 N330 R335 E348 R349 W353 E356 E377 D386 P402 Y408 D427 K428 V429 R430 I431 F432 GLY GLY ARG PRO PRO ALA

Chain I:  81% 13% · ·



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

A horizontal bar chart titled "Chain J:" at the top left. The x-axis represents percentages from 0% to 100%, with major tick marks at 0%, 3%, 84%, 11%, and 100%. A single green bar spans from 0% to approximately 84%. Above the bar, the percentage values are labeled: 3% at the start, 84% in the middle, 11% at the end, and 100% at the far right. Ellipses (...) are placed after the 100% label.



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase


Chain K: 85% 10% ...




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

A horizontal bar chart illustrating the distribution of Chain L. The x-axis represents the percentage of Chain L, ranging from 0% to 100%. The y-axis lists categories: 'Chain L', '82%', '12%', and '5%'. Each category is represented by a colored bar: red for 'Chain L' (2%), yellow for '82%', orange for '12%', and dark blue for '5%'.

Category	Percentage
Chain L	2%
82%	82%
12%	12%
5%	5%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	139.77Å 171.96Å 156.11Å 90.00° 90.02° 90.00°	Depositor
Resolution (Å)	39.03 – 2.78 39.03 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.03-2.78) 99.7 (39.03-2.78)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle^1$	2.57 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ????)	Depositor
R , R_{free}	0.198 , 0.231 0.193 , 0.226	Depositor DCC
R_{free} test set	9405 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.885	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 9.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.447 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	40568	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.46 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.6739e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
YE1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/3333	0.53	0/4515
1	B	0.28	0/3311	0.52	0/4485
1	C	0.26	0/3304	0.51	0/4475
1	D	0.29	0/3333	0.51	0/4513
1	E	0.29	1/3333 (0.0%)	0.52	0/4515
1	F	0.31	1/3333 (0.0%)	0.52	0/4515
1	G	0.28	0/3316	0.52	0/4492
1	H	0.28	0/3311	0.51	0/4485
1	I	0.27	0/3333	0.51	0/4515
1	J	0.28	0/3313	0.51	0/4487
1	K	0.27	0/3320	0.51	0/4497
1	L	0.27	0/3306	0.51	0/4478
All	All	0.28	2/39846 (0.0%)	0.51	0/53972

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	317	LEU	C-N	5.70	1.45	1.34
1	E	324	ILE	C-N	5.20	1.46	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3272	0	3293	26	0
1	B	3251	0	3270	33	0
1	C	3244	0	3263	37	0
1	D	3273	0	3292	23	0
1	E	3272	0	3293	37	0
1	F	3272	0	3293	42	0
1	G	3256	0	3275	37	0
1	H	3251	0	3270	37	0
1	I	3272	0	3293	57	0
1	J	3253	0	3271	28	0
1	K	3260	0	3276	28	0
1	L	3246	0	3269	41	0
2	A	59	0	37	3	0
2	B	59	0	39	3	0
2	C	59	0	39	6	0
2	D	59	0	38	2	0
2	E	59	0	39	4	0
2	F	59	0	38	4	0
2	G	59	0	37	7	0
2	H	59	0	39	3	0
2	I	59	0	38	7	0
2	J	59	0	38	2	0
2	K	59	0	39	4	0
2	L	59	0	38	2	0
3	A	58	0	0	0	0
3	B	56	0	0	2	0
3	C	62	0	0	1	0
3	D	61	0	0	3	0
3	E	74	0	0	5	0
3	F	60	0	0	4	0
3	G	65	0	0	3	0
3	H	66	0	0	4	0
3	I	53	0	0	1	0
3	J	53	0	0	1	0
3	K	67	0	0	4	0
3	L	63	0	0	2	0
All	All	40568	0	39817	417	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 417 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:319:ALA:HB1	1:E:324:ILE:O	1.54	1.08
1:I:322:GLU:CG	1:I:428:LYS:HG3	1.87	1.04
1:E:322:GLU:HG3	1:E:428:LYS:HE2	1.07	1.02
1:E:355:LYS:HB2	3:E:633:HOH:O	1.65	0.96
1:E:322:GLU:HG3	1:E:428:LYS:CE	1.99	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	419/438 (96%)	401 (96%)	12 (3%)	6 (1%)	13 37
1	B	414/438 (94%)	404 (98%)	7 (2%)	3 (1%)	25 57
1	C	413/438 (94%)	405 (98%)	8 (2%)	0	100 100
1	D	417/438 (95%)	411 (99%)	6 (1%)	0	100 100
1	E	419/438 (96%)	399 (95%)	16 (4%)	4 (1%)	18 47
1	F	419/438 (96%)	407 (97%)	8 (2%)	4 (1%)	18 47
1	G	415/438 (95%)	405 (98%)	9 (2%)	1 (0%)	51 82
1	H	414/438 (94%)	405 (98%)	9 (2%)	0	100 100
1	I	419/438 (96%)	407 (97%)	11 (3%)	1 (0%)	51 82
1	J	415/438 (95%)	405 (98%)	9 (2%)	1 (0%)	51 82
1	K	415/438 (95%)	405 (98%)	10 (2%)	0	100 100
1	L	414/438 (94%)	402 (97%)	11 (3%)	1 (0%)	51 82
All	All	4993/5256 (95%)	4856 (97%)	116 (2%)	21 (0%)	38 70

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	HIS
1	A	322	GLU
1	B	270	ASP
1	F	40	SER
1	F	322	GLU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	331/345 (96%)	316 (96%)	15 (4%)	32 64
1	B	330/345 (96%)	321 (97%)	9 (3%)	50 81
1	C	329/345 (95%)	321 (98%)	8 (2%)	54 84
1	D	332/345 (96%)	319 (96%)	13 (4%)	37 70
1	E	331/345 (96%)	318 (96%)	13 (4%)	37 70
1	F	331/345 (96%)	319 (96%)	12 (4%)	40 73
1	G	330/345 (96%)	319 (97%)	11 (3%)	43 75
1	H	330/345 (96%)	323 (98%)	7 (2%)	59 86
1	I	331/345 (96%)	319 (96%)	12 (4%)	40 73
1	J	329/345 (95%)	316 (96%)	13 (4%)	36 69
1	K	331/345 (96%)	320 (97%)	11 (3%)	43 75
1	L	328/345 (95%)	317 (97%)	11 (3%)	42 74
All	All	3963/4140 (96%)	3828 (97%)	135 (3%)	42 74

5 of 135 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	163	GLU
1	G	303	VAL
1	L	56	LEU
1	F	224	ARG
1	F	386	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	100	GLN
1	H	79	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	YE1	A	501	-	55,62,62	2.22	7 (12%)	66,92,92	1.29	9 (13%)
2	YE1	B	501	-	55,62,62	1.94	6 (10%)	66,92,92	1.29	5 (7%)
2	YE1	C	501	-	55,62,62	2.12	8 (14%)	66,92,92	1.44	7 (10%)
2	YE1	D	501	-	55,62,62	1.95	7 (12%)	66,92,92	1.23	6 (9%)
2	YE1	E	501	-	55,62,62	1.93	6 (10%)	66,92,92	1.35	9 (13%)
2	YE1	F	501	-	55,62,62	2.05	10 (18%)	66,92,92	1.40	9 (13%)
2	YE1	G	501	-	55,62,62	2.22	7 (12%)	66,92,92	1.29	9 (13%)
2	YE1	H	501	-	55,62,62	1.96	9 (16%)	66,92,92	1.48	10 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	YE1	I	501	-	55,62,62	2.05	11 (20%)	66,92,92	1.44	9 (13%)
2	YE1	J	501	-	55,62,62	2.00	8 (14%)	66,92,92	1.39	11 (16%)
2	YE1	K	501	-	55,62,62	2.02	8 (14%)	66,92,92	1.39	9 (13%)
2	YE1	L	501	-	55,62,62	1.96	7 (12%)	66,92,92	1.36	7 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YE1	A	501	-	-	0/51/71/71	0/4/4/4
2	YE1	B	501	-	-	0/51/71/71	0/4/4/4
2	YE1	C	501	-	-	0/51/71/71	0/4/4/4
2	YE1	D	501	-	-	0/51/71/71	0/4/4/4
2	YE1	E	501	-	-	0/51/71/71	0/4/4/4
2	YE1	F	501	-	-	0/51/71/71	0/4/4/4
2	YE1	G	501	-	-	0/51/71/71	0/4/4/4
2	YE1	H	501	-	-	0/51/71/71	0/4/4/4
2	YE1	I	501	-	-	0/51/71/71	0/4/4/4
2	YE1	J	501	-	-	0/51/71/71	0/4/4/4
2	YE1	K	501	-	-	0/51/71/71	0/4/4/4
2	YE1	L	501	-	-	0/51/71/71	0/4/4/4

The worst 5 of 94 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	501	YE1	O10-C10	-3.00	1.36	1.42
2	J	501	YE1	O4'-C1'	-2.67	1.37	1.41
2	F	501	YE1	O3'-C3'	-2.55	1.34	1.44
2	H	501	YE1	O3'-C3'	-2.55	1.34	1.44
2	I	501	YE1	O3'-C3'	-2.49	1.34	1.44

The worst 5 of 100 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	501	YE1	C4'-O4'-C1'	-5.92	103.47	109.77
2	H	501	YE1	C4'-O4'-C1'	-4.49	104.99	109.77
2	C	501	YE1	CAC-CAB-NAA	-4.21	110.28	116.25
2	H	501	YE1	C6P-C5P-N4P	-4.02	109.56	116.49
2	K	501	YE1	C6P-C5P-N4P	-3.98	109.62	116.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 47 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	YE1	3	0
2	B	501	YE1	3	0
2	C	501	YE1	6	0
2	D	501	YE1	2	0
2	E	501	YE1	4	0
2	F	501	YE1	4	0
2	G	501	YE1	7	0
2	H	501	YE1	3	0
2	I	501	YE1	7	0
2	J	501	YE1	2	0
2	K	501	YE1	4	0
2	L	501	YE1	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	421/438 (96%)	0.31	14 (3%)	47	40	24, 40, 88, 104	0
1	B	418/438 (95%)	0.13	1 (0%)	94	95	23, 38, 59, 72	0
1	C	417/438 (95%)	0.25	4 (0%)	82	79	23, 38, 66, 78	0
1	D	421/438 (96%)	0.04	3 (0%)	87	85	20, 30, 48, 73	0
1	E	421/438 (96%)	0.13	1 (0%)	94	95	21, 36, 55, 72	0
1	F	421/438 (96%)	0.12	5 (1%)	79	75	21, 34, 54, 77	0
1	G	419/438 (95%)	0.08	1 (0%)	94	95	20, 32, 54, 72	0
1	H	418/438 (95%)	0.19	2 (0%)	90	89	23, 39, 57, 76	0
1	I	421/438 (96%)	0.15	2 (0%)	90	89	21, 36, 54, 67	0
1	J	419/438 (95%)	0.27	13 (3%)	49	42	26, 42, 75, 91	0
1	K	419/438 (95%)	0.12	1 (0%)	94	95	20, 34, 52, 64	0
1	L	418/438 (95%)	0.29	10 (2%)	59	53	22, 37, 77, 96	0
All	All	5033/5256 (95%)	0.17	57 (1%)	80	77	20, 36, 63, 104	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	430	GLY	5.6
1	A	37	ALA	5.4
1	A	430	GLY	4.3
1	A	35	HIS	4.2
1	J	37	ALA	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	YE1	G	501	59/59	0.94	0.21	0.63	25,39,63,71	0
2	YE1	B	501	59/59	0.95	0.18	-0.14	30,48,66,72	0
2	YE1	K	501	59/59	0.96	0.18	-0.21	26,35,46,54	0
2	YE1	A	501	59/59	0.93	0.19	-0.25	32,46,70,79	0
2	YE1	H	501	59/59	0.95	0.19	-0.27	32,38,45,46	0
2	YE1	D	501	59/59	0.96	0.18	-0.32	24,35,44,48	0
2	YE1	L	501	59/59	0.95	0.19	-0.41	35,40,50,55	0
2	YE1	J	501	59/59	0.95	0.18	-0.44	37,48,62,64	0
2	YE1	E	501	59/59	0.95	0.17	-0.57	30,41,51,55	0
2	YE1	F	501	59/59	0.96	0.17	-0.60	27,37,50,57	0
2	YE1	I	501	59/59	0.96	0.18	-0.61	28,36,42,48	0
2	YE1	C	501	59/59	0.94	0.18	-0.62	31,43,58,66	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.