



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2017 – 01:10 PM EDT

PDB ID : 5KFJ
Title : Human DNA polymerase eta-DNA ternary complex: reaction with 10 mM Mn²⁺ for 180s
Authors : Gao, Y.; Yang, W.
Deposited on : unknown
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>8%</div> <div> <div></div> <div>91%</div> <div>7%</div> </div> </div>
2	T	12	<div> <div>8%</div> <div> <div></div> <div>83%</div> <div>17%</div> </div> </div>
3	P	9	<div> <div></div> <div> <div></div> <div>67%</div> <div>22%</div> <div>11%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	504	-	-	-	X
6	GOL	A	505	-	-	-	X
6	GOL	A	506	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	20	0
			3450	2155	627	639	29			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y253
A	-1	PRO	-	expression tag	UNP Q9Y253
A	0	HIS	-	expression tag	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*TP*TP*AP*TP*GP*AP*CP*GP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	12	Total	C	N	O	P	0	1	1
			246	118	41	75	12			

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*GP*CP*GP*TP*CP*AP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	P	0	4	0
			208	99	37	62	10			


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	1	Total	Mn	0	1
			1	1		
4	A	2	Total	Mn	0	1
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	1
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: $C_{10}H_{16}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	
			30	10	5	12	3	
								0
								1

- Molecule 8 is DIPHOSPHATE (three-letter code: DPO) (formula: O_7P_2).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	O P		
			9	7 2	0	1

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	404	Total 405	O 405	0	5
9	T	34	Total 34	O 34	0	0
9	P	26	Total 26	O 26	0	0

- Molecule 1: DNA polymerase eta



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	98.12Å 98.12Å 82.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.99 – 1.70 19.99 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.99-1.70) 100.0 (19.99-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.168 , 0.210 0.168 , 0.210	Depositor DCC
R_{free} test set	3827 reflections (7.74%)	DCC
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4430	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DPO, GOL, CA, MN, DTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3586	0.52	0/4833
2	T	0.74	0/274	1.05	0/422
3	P	4.05	2/272 (0.7%)	1.82	2/419 (0.5%)
All	All	1.12	2/4132 (0.0%)	0.75	2/5674 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	8[A]	DT	O3'-P	46.46	2.16	1.61
3	P	8[B]	DT	O3'-P	46.46	2.16	1.61

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	8[A]	DT	P-O3'-C3'	-21.88	93.45	119.70
3	P	8[B]	DT	P-O3'-C3'	-21.88	93.45	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3450	0	3519	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	T	246	0	138	1	0
3	P	208	0	99	3	0
4	A	2	0	0	0	0
4	P	1	0	0	0	0
5	A	1	0	0	0	0
6	A	18	0	24	0	0
7	A	30	0	11	0	0
8	A	9	0	0	0	0
9	A	405	0	0	4	2
9	P	26	0	0	1	0
9	T	34	0	0	0	0
All	All	4430	0	3791	21	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:8[B]:DT:H2''	9:P:203:HOH:O	1.11	1.26
3:P:8[B]:DT:H2'	3:P:9[B]:DA:C8	2.24	0.72
1:A:256:ARG:NH1	9:A:604:HOH:O	2.24	0.66
1:A:372:VAL:HG11	1:A:417:PRO:HG2	1.81	0.61
2:T:3[A]:DT:H2'	2:T:4:DT:H71	1.88	0.55
1:A:231[B]:LYS:HE3	9:A:956:HOH:O	2.09	0.52
1:A:319:ILE:HG23	1:A:351[B]:ARG:CZ	2.43	0.49
1:A:367:VAL:HG22	1:A:384[B]:CYS:SG	2.53	0.49
1:A:369:SER:HB3	1:A:423:PHE:HB3	1.97	0.47
1:A:162:GLN:HG3	1:A:165:GLY:H	1.80	0.47
1:A:372:VAL:CG1	1:A:417:PRO:HG2	2.44	0.47
1:A:180:ILE:HA	1:A:188:LEU:HD22	1.98	0.45
1:A:356:ARG:NH2	9:A:618:HOH:O	2.46	0.44
1:A:106:PHE:CG	1:A:200[B]:MET:HG2	2.52	0.44
1:A:290:PHE:O	1:A:294:ASN:HB2	2.18	0.43
1:A:368:VAL:HG21	1:A:399:ALA:HA	2.02	0.42
1:A:398:ASP:O	1:A:402:VAL:HG13	2.19	0.42
3:P:7:DA:H2''	3:P:8[A]:DT:H5'	2.02	0.41
1:A:-1:PRO:HG2	1:A:0:HIS:CD2	2.56	0.41
1:A:318:THR:HG22	1:A:427:THR:HG22	2.02	0.41
1:A:351[B]:ARG:NH2	9:A:627:HOH:O	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:732:HOH:O	9:A:791:HOH:O[5_664]	2.10	0.10
9:A:802:HOH:O	9:A:862:HOH:O[5_554]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/435 (102%)	438 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/372 (104%)	382 (98%)	6 (2%)	70	55

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	ASP
1	A	119	VAL
1	A	321[A]	CYS
1	A	321[B]	CYS
1	A	381	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	408	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	504	-	5,5,5	0.54	0	5,5,5	0.29	0
6	GOL	A	505	-	5,5,5	0.35	0	5,5,5	0.25	0
6	GOL	A	506	-	5,5,5	0.30	0	5,5,5	0.19	0
7	DTP	A	507[A]	5,4	26,32,32	1.25	4 (15%)	26,50,50	2.24	3 (11%)
8	DPO	A	508[B]	4	8,8,8	1.26	1 (12%)	8,13,13	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	504	-	-	0/4/4/4	0/0/0/0
6	GOL	A	505	-	-	0/4/4/4	0/0/0/0
6	GOL	A	506	-	-	0/4/4/4	0/0/0/0
7	DTP	A	507[A]	5,4	-	0/18/34/34	0/3/3/3
8	DPO	A	508[B]	4	-	0/6/6/6	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	507[A]	DTP	C3'-C4'	-2.78	1.45	1.53
8	A	508[B]	DPO	P1-O4	-2.48	1.56	1.60
7	A	507[A]	DTP	C2'-C3'	-2.38	1.46	1.52
7	A	507[A]	DTP	C6-N6	2.39	1.43	1.34
7	A	507[A]	DTP	C2-N3	2.97	1.37	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	507[A]	DTP	N3-C2-N1	-10.36	119.84	128.86
7	A	507[A]	DTP	C4-C5-N7	-2.18	107.30	109.41
7	A	507[A]	DTP	C2'-C1'-N9	-2.13	109.21	114.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/435 (98%)	0.15	35 (8%) 13 15	9, 19, 55, 81	0
2	T	12/12 (100%)	-0.04	1 (8%) 12 14	17, 23, 51, 53	0
3	P	9/9 (100%)	-0.53	0 100 100	14, 20, 29, 33	1 (11%)
All	All	451/456 (98%)	0.13	36 (7%) 13 16	9, 19, 55, 81	1 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	7.7
1	A	378	LEU	6.9
1	A	376	LYS	6.6
1	A	411	ILE	6.1
1	A	412	GLN	5.6
1	A	-1	PRO	5.0
1	A	160	THR	4.7
1	A	0	HIS	4.5
1	A	2	ALA	4.3
1	A	410	GLY	4.3
1	A	154	PRO	4.3
1	A	161	VAL	4.2
1	A	153	GLY	4.1
1	A	183	LEU	3.8
1	A	133	GLN	3.8
1	A	374	GLY	3.8
1	A	181	ASP	3.7
1	A	3	THR	3.6
1	A	180	ILE	3.5
1	A	408	THR	3.4
1	A	134	GLY	3.4
1	A	432	SER	3.4
1	A	179	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	T	3[A]	DT	3.3
1	A	327	GLY	2.8
1	A	162	GLN	2.7
1	A	182	ASN	2.4
1	A	184	THR	2.3
1	A	132	LEU	2.3
1	A	328	LYS	2.3
1	A	131	LYS	2.3
1	A	130	GLN	2.2
1	A	380	SER	2.1
1	A	127	GLU	2.1
1	A	135	GLN	2.1
1	A	373	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	GOL	A	506	6/6	0.75	0.30	11.14	29,38,45,47	0
6	GOL	A	504	6/6	0.94	0.15	6.48	16,18,24,28	0
6	GOL	A	505	6/6	0.90	0.16	3.64	18,30,35,45	0
7	DTP	A	507[A]	30/30	0.98	0.05	-0.81	10,12,15,16	30
8	DPO	A	508[B]	9/9	0.99	0.04	-1.65	10,11,12,13	9
5	CA	A	502[A]	1/1	0.99	0.04	-1.78	12,12,12,12	1
4	MN	A	503[B]	1/1	0.99	0.04	-1.78	12,12,12,12	0
4	MN	A	501	1/1	1.00	0.03	-2.31	12,12,12,12	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MN	P	101[B]	1/1	0.99	0.08	-	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.