



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 07:42 PM EST

PDB ID : 4KH0  
Title : The R state structure of E. coli ATCase with ATP and Magnesium bound  
Authors : Cockrell, G.M.; Zheng, Y.; Guo, W.; Peterson, A.W.; Kantrowitz, E.R.  
Deposited on : unknown  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	<div> <div style="width: 93%;"></div> <div style="width: 7%;"></div> </div>
1	C	310	<div> <div style="width: 89%;"></div> <div style="width: 11%;"></div> </div>
2	B	153	<div> <div style="width: 4%;"></div> <div style="width: 86%;"></div> <div style="width: 9%;"></div> <div style="width: 1%;"></div> </div>
2	D	153	<div> <div style="width: 7%;"></div> <div style="width: 77%;"></div> <div style="width: 20%;"></div> <div style="width: 1%;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	B	201	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aspartate carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			
1	C	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			

- Molecule 2 is a protein called Aspartate carbamoyltransferase regulatory chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	150	Total	C	N	O	S	0	0	0
			1176	737	208	226	5			
2	D	152	Total	C	N	O	S	0	0	0
			1193	747	212	229	5			

- Molecule 3 is N-(PHOSPHONACETYL)-L-ASPARTIC ACID (three-letter code: PAL) (formula: C<sub>6</sub>H<sub>10</sub>NO<sub>8</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			16	6	1	8	1		
3	C	1	Total	C	N	O	P	0	0
			16	6	1	8	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Mg 1	0	0
6	D	1	Total 1	Mg 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	184	Total 184	O 184	0	0
7	B	69	Total 69	O 69	0	0
7	C	165	Total 165	O 165	0	0
7	D	49	Total 49	O 49	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate carbamoyltransferase

Chain A: 




- Molecule 1: Aspartate carbamoyltransferase

Chain C: 




- Molecule 2: Aspartate carbamoyltransferase regulatory chain

Chain B: 



- Molecule 2: Aspartate carbamoyltransferase regulatory chain

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.13Å 121.13Å 155.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.69 – 2.25 49.69 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.69-2.25) 99.8 (49.69-2.25)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.33 (at 2.25Å)	Xtriage
Refinement program	PHENIX 1.8.1 _1168	Depositor
R, $R_{free}$	0.168 , 0.208 0.166 , 0.207	Depositor DCC
$R_{free}$ test set	3183 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7826	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PAL, ZN, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2461	0.55	0/3339
1	C	0.37	0/2461	0.52	0/3339
2	B	0.34	0/1193	0.51	0/1612
2	D	0.31	0/1211	0.51	0/1637
All	All	0.36	0/7326	0.53	0/9927

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	0	2422	12	0
1	C	2415	0	2422	17	1
2	B	1176	0	1193	15	0
2	D	1193	0	1207	20	1
3	A	16	0	6	0	0
3	C	16	0	6	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	B	62	0	24	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	62	0	24	2	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	184	0	0	5	0
7	B	69	0	0	2	0
7	C	165	0	0	2	0
7	D	49	0	0	4	0
All	All	7826	0	7304	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:GLU:O	1:A:258:LYS:NZ	2.14	0.81
2:B:56:LYS:NZ	5:B:203:ATP:O2A	2.15	0.78
2:D:144:GLU:OE2	7:D:325:HOH:O	2.06	0.73
1:C:271:ASP:OD1	7:C:665:HOH:O	2.09	0.70
1:A:253:ASP:O	7:A:684:HOH:O	2.10	0.69
1:A:271:ASP:OD1	7:A:670:HOH:O	2.11	0.69
1:A:31:LYS:NZ	7:A:581:HOH:O	2.25	0.68
2:B:43:THR:HG21	5:B:203:ATP:HN62	1.60	0.67
2:B:5:ASN:HB3	2:B:7:LEU:HG	1.77	0.67
2:D:130:ARG:HE	2:D:135:ALA:HB2	1.58	0.66
2:D:22:PRO:HG2	2:D:25:ILE:HG13	1.79	0.64
1:C:35:GLN:HG3	1:C:310:LEU:HD11	1.80	0.64
5:D:202:ATP:O2'	7:D:336:HOH:O	2.15	0.62
2:D:99:LEU:HD12	2:D:100:PRO:HD2	1.82	0.62
2:D:13:LYS:HD2	2:D:88:ASN:HA	1.85	0.58
1:C:130:GLY:O	1:C:167:ARG:HD3	2.03	0.58
2:D:128:ARG:NH1	2:D:130:ARG:HD3	2.20	0.56
1:A:275:THR:O	7:A:597:HOH:O	2.17	0.56
2:D:132:ASN:OD1	7:D:331:HOH:O	2.18	0.56
2:D:2:THR:N	2:D:3:HIS:HD1	2.03	0.55
2:B:41:ARG:NH2	2:B:62:GLU:OE1	2.39	0.55
1:C:279:LYS:HG2	7:C:664:HOH:O	2.07	0.55
2:B:14:ARG:HH11	2:B:65:PHE:HE2	1.55	0.53
2:D:9:VAL:HG21	2:D:43:THR:HG21	1.90	0.53
2:D:76:LEU:HG	2:D:151:LEU:HD11	1.91	0.53
1:A:130:GLY:O	1:A:167:ARG:HD3	2.11	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:66:LEU:HD21	1:C:297:GLN:HE21	1.76	0.51
2:D:72:ASP:HB3	2:D:100:PRO:HG3	1.94	0.50
1:C:104:MET:HE1	1:C:115:ALA:CB	2.42	0.50
1:C:104:MET:HE1	1:C:115:ALA:HB2	1.94	0.49
2:D:130:ARG:NH1	2:D:133:ASP:OD1	2.44	0.49
2:D:130:ARG:NE	2:D:135:ALA:HB2	2.25	0.49
1:C:75:ASP:OD2	1:C:77:ALA:HB3	2.14	0.48
5:D:203:ATP:N7	7:D:336:HOH:O	2.35	0.48
1:A:221:GLU:C	1:A:258:LYS:NZ	2.67	0.47
1:C:232:LYS:HA	1:C:235:LEU:HG	1.96	0.47
1:A:279:LYS:HD3	7:A:597:HOH:O	2.15	0.47
2:B:96:ARG:NH1	7:B:341:HOH:O	2.44	0.46
1:C:33:ASN:HA	1:C:34:PRO:HD3	1.61	0.46
1:C:36:PRO:HA	1:C:65:ARG:O	2.16	0.46
1:A:36:PRO:HA	1:A:65:ARG:O	2.16	0.45
2:B:133:ASP:OD2	2:B:147:HIS:CE1	2.69	0.45
2:B:10:GLU:OE1	2:B:10:GLU:N	2.39	0.45
1:C:237:PRO:HA	1:C:240:TYR:CZ	2.52	0.45
2:B:80:GLN:HA	2:B:96:ARG:NH2	2.31	0.45
2:D:33:PHE:HB2	2:D:35:LEU:HG	1.99	0.45
2:D:13:LYS:HD3	2:D:89:TYR:CE2	2.53	0.44
2:B:80:GLN:CD	2:B:80:GLN:H	2.21	0.44
2:B:94:LYS:NZ	5:B:202:ATP:O3G	2.51	0.44
2:D:134:ILE:HG23	2:D:147:HIS:CG	2.52	0.44
1:C:122:VAL:HA	1:C:123:PRO:HD3	1.88	0.43
2:B:4:ASP:HB3	2:B:5:ASN:H	1.58	0.43
1:C:50:GLU:HG3	1:C:105:ARG:HG2	2.00	0.43
1:C:214:SER:OG	1:C:217:GLU:HB2	2.19	0.43
2:D:42:ILE:HG12	2:D:61:ILE:HG23	2.01	0.42
1:A:194:MET:HA	1:A:195:PRO:HD3	1.90	0.42
1:A:271:ASP:OD1	1:A:271:ASP:N	2.53	0.42
2:B:10:GLU:CD	2:B:10:GLU:H	2.16	0.42
2:B:70:GLN:NE2	7:B:355:HOH:O	2.53	0.42
2:B:44:ILE:HB	2:D:44:ILE:HB	2.01	0.41
1:A:218:VAL:O	1:A:222:VAL:HG13	2.21	0.41
1:C:106:HIS:CG	1:C:107:PRO:HD2	2.56	0.40
1:C:269:ARG:HA	1:C:272:GLU:OE2	2.22	0.40
2:D:110:PRO:HG2	2:D:145:PHE:CE1	2.57	0.40
2:D:45:GLY:HA3	2:D:48:LEU:HD11	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1:ALA:N	2:D:68:GLU:OE2[4_556]	2.05	0.15

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/310 (99%)	299 (97%)	8 (3%)	1 (0%)	44	50
1	C	308/310 (99%)	299 (97%)	9 (3%)	0	100	100
2	B	148/153 (97%)	140 (95%)	8 (5%)	0	100	100
2	D	150/153 (98%)	141 (94%)	7 (5%)	2 (1%)	14	9
All	All	914/926 (99%)	879 (96%)	32 (4%)	3 (0%)	44	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	3	HIS
2	D	131	ALA
1	A	270	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/261 (100%)	254 (97%)	7 (3%)	50	60
1	C	261/261 (100%)	257 (98%)	4 (2%)	70	79

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	134/137 (98%)	127 (95%)	7 (5%)	27	29
2	D	136/137 (99%)	132 (97%)	4 (3%)	48	57
All	All	792/796 (100%)	770 (97%)	22 (3%)	49	58

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	PHE
1	A	75	ASP
1	A	104	MET
1	A	134	HIS
1	A	204	GLU
1	A	214	SER
1	A	285	TYR
2	B	4	ASP
2	B	5	ASN
2	B	10	GLU
2	B	74	LEU
2	B	80	GLN
2	B	95	SER
2	B	147	HIS
1	C	59	PHE
1	C	104	MET
1	C	146	GLN
1	C	285	TYR
2	D	2	THR
2	D	5	ASN
2	D	130	ARG
2	D	134	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	297	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PAL	A	401	-	9,15,15	2.09	3 (33%)	8,21,21	0.73	0
5	ATP	B	202	6	27,33,33	1.04	1 (3%)	25,52,52	1.69	3 (12%)
5	ATP	B	203	6	27,33,33	0.99	1 (3%)	25,52,52	1.59	2 (8%)
3	PAL	C	401	-	9,15,15	2.01	3 (33%)	8,21,21	0.78	0
5	ATP	D	202	6	27,33,33	1.00	1 (3%)	25,52,52	1.63	2 (8%)
5	ATP	D	203	6	27,33,33	1.02	1 (3%)	25,52,52	1.68	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PAL	A	401	-	-	0/11/17/17	0/0/0/0
5	ATP	B	202	6	-	0/18/38/38	0/3/3/3
5	ATP	B	203	6	-	0/18/38/38	0/3/3/3
3	PAL	C	401	-	-	0/11/17/17	0/0/0/0
5	ATP	D	202	6	-	0/18/38/38	0/3/3/3
5	ATP	D	203	6	-	0/18/38/38	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	PAL	C2-N2	-2.77	1.42	1.46
3	C	401	PAL	C3-C2	-2.59	1.50	1.53
3	A	401	PAL	C2-N2	-2.55	1.43	1.46
3	A	401	PAL	C3-C2	-2.42	1.50	1.53
5	D	202	ATP	C5-C4	3.10	1.47	1.40
5	B	203	ATP	C5-C4	3.17	1.47	1.40
5	D	203	ATP	C5-C4	3.28	1.47	1.40
5	B	202	ATP	C5-C4	3.32	1.48	1.40
3	C	401	PAL	C1-N2	4.04	1.42	1.34
3	A	401	PAL	C1-N2	4.37	1.43	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	203	ATP	N3-C2-N1	-6.01	123.62	128.86
5	B	202	ATP	N3-C2-N1	-5.99	123.64	128.86
5	D	202	ATP	N3-C2-N1	-5.97	123.65	128.86
5	B	203	ATP	N3-C2-N1	-5.97	123.65	128.86
5	D	203	ATP	C4-C5-N7	-3.20	106.32	109.41
5	D	202	ATP	C4-C5-N7	-3.13	106.38	109.41
5	B	202	ATP	C4-C5-N7	-3.01	106.50	109.41
5	B	203	ATP	C4-C5-N7	-2.64	106.86	109.41
5	B	202	ATP	C2-N1-C6	2.06	122.37	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	202	ATP	1	0
5	B	203	ATP	2	0
5	D	202	ATP	1	0
5	D	203	ATP	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	310/310 (100%)	-0.52	0	100	100	20, 37, 62, 107	0
1	C	310/310 (100%)	-0.70	0	100	100	24, 38, 62, 96	0
2	B	150/153 (98%)	-0.25	6 (4%)	39	42	28, 53, 93, 148	0
2	D	152/153 (99%)	-0.07	11 (7%)	16	18	35, 60, 113, 135	0
All	All	922/926 (99%)	-0.46	17 (1%)	69	72	20, 43, 79, 148	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	7	LEU	7.7
2	B	4	ASP	6.9
2	B	8	GLN	6.0
2	B	9	VAL	5.0
2	D	153	ASN	4.3
2	B	5	ASN	4.1
2	D	131	ALA	3.9
2	D	9	VAL	3.8
2	D	132	ASN	3.1
2	D	8	GLN	3.1
2	D	129	LYS	3.0
2	D	5	ASN	2.9
2	D	7	LEU	2.9
2	D	152	ALA	2.8
2	B	6	LYS	2.5
2	D	151	LEU	2.4
2	D	3	HIS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ZN	B	201	1/1	1.00	0.11	3.43	31,31,31,31	0
4	ZN	D	201	1/1	1.00	0.08	1.04	38,38,38,38	0
3	PAL	A	401	16/16	0.99	0.12	0.39	13,27,32,33	0
3	PAL	C	401	16/16	0.98	0.11	0.39	22,31,36,39	0
5	ATP	B	203	31/31	0.92	0.18	-0.07	86,121,132,253	0
5	ATP	D	203	31/31	0.96	0.14	-0.38	66,120,129,131	0
5	ATP	D	202	31/31	0.96	0.09	-0.64	48,64,77,87	0
5	ATP	B	202	31/31	0.95	0.10	-0.85	46,63,84,90	0
6	MG	B	204	1/1	0.80	0.04	-	84,84,84,84	0
6	MG	D	204	1/1	0.96	0.07	-	67,67,67,67	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.