



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 12:42 am GMT

PDB ID : 4KIN
Title : Crystal structure of mitogen-activated protein kinase 14 (P38-H5) complex with 5-(2-CHLOROPHENYL)-N-(5-(CYCLOPROPYLCARBAMOYL)-2-METHYLPHENYL)-2-THIOPHENECARBOXAMIDE
Authors : Sack, J.S.; Tokarski, J.S.
Deposited on : 2013-05-02
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	9293 (2.00-1.96)
Clashscore	112137	10621 (2.00-1.96)
Ramachandran outliers	110173	10502 (2.00-1.96)
Sidechain outliers	110143	10501 (2.00-1.96)
RSRZ outliers	101464	9395 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>• • 8%</div> </div> </div>
1	B	366	<div> <div>7%</div> <div> <div></div> <div>86%</div> <div>6% • 7%</div> </div> </div>
1	C	366	<div> <div>10%</div> <div> <div></div> <div>86%</div> <div>5% • 8%</div> </div> </div>
1	D	366	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>5% 7%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2721	1748	463	498	12			
1	B	341	Total	C	N	O	S	0	0	0
			2748	1764	470	502	12			
1	C	338	Total	C	N	O	S	0	0	0
			2729	1751	469	497	12			
1	D	339	Total	C	N	O	S	0	0	0
			2741	1759	470	500	12			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	EXPRESSION TAG	UNP Q16539
A	-4	ALA	-	EXPRESSION TAG	UNP Q16539
A	-3	HIS	-	EXPRESSION TAG	UNP Q16539
A	-2	HIS	-	EXPRESSION TAG	UNP Q16539
A	-1	HIS	-	EXPRESSION TAG	UNP Q16539
A	0	HIS	-	EXPRESSION TAG	UNP Q16539
A	1	HIS	-	EXPRESSION TAG	UNP Q16539
B	-5	MET	-	EXPRESSION TAG	UNP Q16539
B	-4	ALA	-	EXPRESSION TAG	UNP Q16539
B	-3	HIS	-	EXPRESSION TAG	UNP Q16539
B	-2	HIS	-	EXPRESSION TAG	UNP Q16539
B	-1	HIS	-	EXPRESSION TAG	UNP Q16539
B	0	HIS	-	EXPRESSION TAG	UNP Q16539
B	1	HIS	-	EXPRESSION TAG	UNP Q16539
C	-5	MET	-	EXPRESSION TAG	UNP Q16539
C	-4	ALA	-	EXPRESSION TAG	UNP Q16539
C	-3	HIS	-	EXPRESSION TAG	UNP Q16539
C	-2	HIS	-	EXPRESSION TAG	UNP Q16539
C	-1	HIS	-	EXPRESSION TAG	UNP Q16539
C	0	HIS	-	EXPRESSION TAG	UNP Q16539
C	1	HIS	-	EXPRESSION TAG	UNP Q16539

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	MET	-	EXPRESSION TAG	UNP Q16539
D	-4	ALA	-	EXPRESSION TAG	UNP Q16539
D	-3	HIS	-	EXPRESSION TAG	UNP Q16539
D	-2	HIS	-	EXPRESSION TAG	UNP Q16539
D	-1	HIS	-	EXPRESSION TAG	UNP Q16539
D	0	HIS	-	EXPRESSION TAG	UNP Q16539
D	1	HIS	-	EXPRESSION TAG	UNP Q16539

- Molecule 2 is 5-(2-CHLOROPHENYL)-N-[5-(CYCLOPROPYLCARBAMOYL)-2-METHYLPHENYL]THIOPHENE-2-CARBOXAMIDE (three-letter code: 1M8) (formula: C₂₂H₁₉ClN₂O₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 28	C 22	Cl 1	N 2	O 2	S 1	0	0
2	B	1	Total 28	C 22	Cl 1	N 2	O 2	S 1	0	0
2	C	1	Total 28	C 22	Cl 1	N 2	O 2	S 1	0	0
2	D	1	Total 28	C 22	Cl 1	N 2	O 2	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	240	Total	O	0	0
			240	240		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	196	Total 196	O 196	0	0
3	C	129	Total 129	O 129	0	0
3	D	289	Total 289	O 289	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Mitogen-activated protein kinase 14




• Molecule 1: Mitogen-activated protein kinase 14




• Molecule 1: Mitogen-activated protein kinase 14



• Molecule 1: Mitogen-activated protein kinase 14





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.88Å 73.55Å 100.95Å 90.00° 96.38° 90.00°	Depositor
Resolution (Å)	28.33 – 1.97 28.33 – 1.97	Depositor EDS
% Data completeness (in resolution range)	98.5 (28.33-1.97) 98.3 (28.33-1.97)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 1.96Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.184 , 0.217 0.192 , 0.227	Depositor DCC
R_{free} test set	5134 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.114 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11905	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.74 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.6267e-04.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1M8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2784	0.60	0/3780
1	B	0.49	0/2812	0.64	0/3819
1	C	0.48	0/2792	0.64	1/3790 (0.0%)
1	D	0.54	0/2804	0.61	0/3807
All	All	0.50	0/11192	0.62	1/15196 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	198	MET	C-N-CA	5.67	135.88	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2721	0	2712	7	0
1	B	2748	0	2743	15	0
1	C	2729	0	2723	10	0
1	D	2741	0	2735	9	0
2	A	28	0	19	3	0
2	B	28	0	19	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	28	0	19	0	0
2	D	28	0	19	4	0
3	A	240	0	0	0	0
3	B	196	0	0	0	0
3	C	129	0	0	0	0
3	D	289	0	0	1	0
All	All	11905	0	10989	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:70:ARG:CZ	1:D:174:HIS:HB3	2.14	0.78
1:B:182:TYR:N	1:B:182:TYR:HD1	1.95	0.64
1:B:182:TYR:N	1:B:182:TYR:CD1	2.67	0.63
1:B:30:VAL:HB	2:B:401:1M8:H17	1.81	0.63
1:D:108:LEU:O	2:D:401:1M8:H15	2.03	0.58
1:C:197:TRP:HD1	1:C:198:MET:HG3	1.69	0.57
1:C:197:TRP:CD1	1:C:198:MET:HG3	2.40	0.57
1:B:196:ASN:HA	1:B:197:TRP:HB3	1.89	0.54
1:C:116:ILE:HG13	1:C:158:VAL:HG11	1.91	0.53
1:D:75:LEU:HD11	2:D:401:1M8:H2	1.90	0.52
1:B:116:ILE:HG13	1:B:158:VAL:HG11	1.92	0.51
1:D:255:ALA:O	1:D:259:ILE:HD12	2.10	0.51
1:A:118:LYS:HE2	1:A:120:GLN:HA	1.93	0.51
1:A:30:VAL:HB	2:A:401:1M8:H17	1.92	0.51
1:C:195:LEU:HD11	1:C:232:LEU:HD22	1.92	0.51
1:B:192:GLU:HA	1:B:197:TRP:CE3	2.46	0.50
1:A:75:LEU:HD11	2:A:401:1M8:H2	1.93	0.50
1:C:198:MET:HA	1:C:199:HIS:HB3	1.93	0.49
1:B:195:LEU:HD21	1:B:232:LEU:HD22	1.95	0.47
1:B:108:LEU:O	2:B:401:1M8:H15	2.14	0.47
1:C:275:ILE:H	1:C:275:ILE:HG12	1.58	0.46
1:A:189:ARG:HG2	1:A:194:MET:HG3	1.98	0.45
1:B:197:TRP:CE2	1:B:250:ILE:HG12	2.52	0.44
1:C:158:VAL:HG22	1:C:162:CYS:HA	1.99	0.44
1:A:246:LEU:HA	1:A:249:LYS:HZ2	1.82	0.44
1:D:263:THR:HG22	1:D:265:MET:HE1	1.99	0.44
1:D:53:LYS:HB2	2:D:401:1M8:H6	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:VAL:HG21	1:A:291:LEU:HD13	2.01	0.43
1:D:112:ASP:HB3	2:D:401:1M8:CL1	2.56	0.43
1:B:75:LEU:HD11	2:B:401:1M8:H2	2.01	0.42
1:C:284:LEU:HG	1:C:288:MET:HE2	2.02	0.42
1:D:74:LEU:HG	3:D:525:HOH:O	2.19	0.42
1:D:147:ILE:HD11	1:D:149:ARG:NE	2.34	0.41
1:A:108:LEU:O	2:A:401:1M8:H15	2.19	0.41
1:B:239:VAL:HG21	1:B:291:LEU:HD13	2.02	0.41
1:B:120:GLN:HG3	1:B:120:GLN:H	1.75	0.41
1:C:239:VAL:HG21	1:C:291:LEU:HD13	2.02	0.41
1:B:229:ILE:HG23	1:B:258:TYR:OH	2.21	0.40
1:C:121:LYS:HG3	1:C:217:LEU:O	2.22	0.40
1:B:182:TYR:HA	1:B:183:VAL:HA	1.66	0.40
1:B:190:ALA:HB3	1:B:193:ILE:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/366 (91%)	322 (97%)	9 (3%)	1 (0%)	44	38
1	B	337/366 (92%)	328 (97%)	8 (2%)	1 (0%)	44	38
1	C	332/366 (91%)	319 (96%)	9 (3%)	4 (1%)	15	7
1	D	333/366 (91%)	326 (98%)	5 (2%)	2 (1%)	28	20
All	All	1334/1464 (91%)	1295 (97%)	31 (2%)	8 (1%)	28	20

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	196	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	264	GLN
1	D	14	ASN
1	D	173	ARG
1	B	196	ASN
1	C	173	ARG
1	C	199	HIS
1	A	183	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/325 (91%)	290 (98%)	7 (2%)	54	53
1	B	300/325 (92%)	289 (96%)	11 (4%)	39	33
1	C	298/325 (92%)	290 (97%)	8 (3%)	50	48
1	D	300/325 (92%)	297 (99%)	3 (1%)	80	83
All	All	1195/1300 (92%)	1166 (98%)	29 (2%)	54	53

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	67	ARG
1	A	113	LEU
1	A	120	GLN
1	A	173	ARG
1	A	183	VAL
1	A	275	ILE
1	B	27	LEU
1	B	32	SER
1	B	113	LEU
1	B	120	GLN
1	B	182	TYR
1	B	183	VAL
1	B	186	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	193	ILE
1	B	198	MET
1	B	233	LYS
1	B	275	ILE
1	C	27	LEU
1	C	113	LEU
1	C	186	ARG
1	C	197	TRP
1	C	199	HIS
1	C	220	ARG
1	C	233	LYS
1	C	275	ILE
1	D	26	ASN
1	D	27	LEU
1	D	186	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1M8	A	401	-	28,31,31	0.27	0	35,44,44	0.62	0
2	1M8	B	401	-	28,31,31	0.25	0	35,44,44	0.59	0
2	1M8	C	401	-	28,31,31	0.32	0	35,44,44	0.56	0
2	1M8	D	401	-	28,31,31	0.25	0	35,44,44	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1M8	A	401	-	-	0/16/22/22	0/3/4/4
2	1M8	B	401	-	-	0/16/22/22	0/3/4/4
2	1M8	C	401	-	-	0/16/22/22	0/3/4/4
2	1M8	D	401	-	-	0/16/22/22	0/3/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	1M8	3	0
2	B	401	1M8	3	0
2	D	401	1M8	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/366 (92%)	0.08	14 (4%) 38 41	15, 27, 58, 75	0
1	B	341/366 (93%)	0.19	27 (7%) 13 15	18, 31, 65, 95	0
1	C	338/366 (92%)	0.50	36 (10%) 7 8	21, 39, 71, 94	0
1	D	339/366 (92%)	-0.17	9 (2%) 55 58	16, 24, 45, 87	0
All	All	1356/1464 (92%)	0.15	86 (6%) 21 23	15, 30, 63, 95	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	MET	11.6
1	B	183	VAL	10.2
1	B	198	MET	9.5
1	A	183	VAL	9.4
1	C	119	CYS	9.2
1	B	197	TRP	8.2
1	A	117	VAL	8.2
1	B	116	ILE	7.8
1	B	119	CYS	7.6
1	C	33	GLY	7.1
1	D	174	HIS	6.9
1	A	119	CYS	6.8
1	B	199	HIS	6.5
1	C	121	LYS	6.4
1	C	197	TRP	5.9
1	C	116	ILE	5.7
1	C	185	THR	5.6
1	C	120	GLN	5.5
1	C	195	LEU	5.4
1	B	115	ASN	5.3
1	B	258	TYR	5.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	258	TYR	5.2
1	D	36	GLY	5.0
1	A	197	TRP	4.6
1	C	117	VAL	4.5
1	A	199	HIS	4.4
1	C	174	HIS	4.4
1	C	160	GLU	4.4
1	A	184	ALA	4.3
1	C	36	GLY	4.0
1	B	118	LYS	4.0
1	B	32	SER	3.9
1	B	117	VAL	3.9
1	D	198	MET	3.9
1	B	193	ILE	3.7
1	D	354	ASP	3.5
1	C	14	ASN	3.3
1	C	173	ARG	3.3
1	C	199	HIS	3.3
1	C	352	PRO	3.3
1	C	196	ASN	3.2
1	B	184	ALA	3.1
1	B	260	GLN	3.1
1	C	184	ALA	3.0
1	C	48	LEU	3.0
1	C	35	TYR	2.9
1	B	253	GLU	2.8
1	C	186	ARG	2.8
1	C	258	TYR	2.8
1	D	35	TYR	2.6
1	C	49	ARG	2.6
1	B	232	LEU	2.6
1	A	195	LEU	2.6
1	B	196	ASN	2.6
1	C	30	VAL	2.6
1	B	195	LEU	2.6
1	C	346	ILE	2.6
1	C	226	THR	2.6
1	B	262	LEU	2.5
1	C	27	LEU	2.5
1	C	122	LEU	2.5
1	D	14	ASN	2.5
1	D	199	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	162	CYS	2.4
1	C	22	GLU	2.4
1	A	275	ILE	2.4
1	C	13	LEU	2.3
1	B	31	GLY	2.3
1	C	115	ASN	2.3
1	B	352	PRO	2.2
1	A	182	TYR	2.2
1	B	228	HIS	2.2
1	B	14	ASN	2.2
1	A	352	PRO	2.2
1	C	260	GLN	2.2
1	C	265	MET	2.1
1	B	35	TYR	2.1
1	D	200	TYR	2.1
1	B	186	ARG	2.1
1	B	42	PHE	2.1
1	C	42	PHE	2.0
1	A	120	GLN	2.0
1	A	264	GLN	2.0
1	B	120	GLN	2.0
1	C	60	GLN	2.0
1	D	310	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	1M8	A	401	28/28	0.96	0.09	0.00	12,19,38,40	0
2	1M8	B	401	28/28	0.95	0.09	-0.21	17,22,46,48	0
2	1M8	C	401	28/28	0.93	0.11	-0.28	25,29,51,57	0
2	1M8	D	401	28/28	0.97	0.08	-0.63	13,19,30,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.