



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:37 am GMT

PDB ID : 1KLF
Title : FIMH ADHESIN-FIMC CHAPERONE COMPLEX WITH D-MANNOSE
Authors : Hung, C.S.; Bouckaert, J.
Deposited on : 2001-12-11
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	<div><div>5%</div><div><div>45%</div><div>47%</div><div>8%</div></div></div>
1	C	205	<div><div>6%</div><div><div>45%</div><div>46%</div><div>8%</div></div></div>
1	E	205	<div><div>%</div><div><div>46%</div><div>46%</div><div>8%</div></div></div>
1	G	205	<div><div>8%</div><div><div>43%</div><div>49%</div><div>8%</div></div></div>
1	I	205	<div><div>71%</div><div><div>38%</div><div>55%</div><div>6%</div></div></div>
1	K	205	<div><div>83%</div><div><div>39%</div><div>54%</div><div>6%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	M	205	<div><div></div><div>80%</div><div>38%</div><div>56%</div><div>6%</div></div>
1	O	205	<div><div></div><div>72%</div><div>38%</div><div>55%</div><div>6%</div></div>
2	B	279	<div><div></div><div>2%</div><div>57%</div><div>35%</div><div>8%</div></div>
2	D	279	<div><div></div><div>3%</div><div>58%</div><div>34%</div><div>8%</div></div>
2	F	279	<div><div></div><div>%</div><div>58%</div><div>34%</div><div>8%</div></div>
2	H	279	<div><div></div><div>2%</div><div>56%</div><div>36%</div><div>8%</div></div>
2	J	279	<div><div></div><div>38%</div><div>34%</div><div>58%</div><div>8%</div></div>
2	L	279	<div><div></div><div>44%</div><div>33%</div><div>58%</div><div>8%</div></div>
2	N	279	<div><div></div><div>51%</div><div>34%</div><div>57%</div><div>8%</div></div>
2	P	279	<div><div></div><div>39%</div><div>33%</div><div>59%</div><div>8%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 29775 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHAPERONE PROTEIN FIMC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	C	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	E	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	G	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	I	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	K	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	M	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			
1	O	205	Total	C	N	O	S	0	0	0
			1596	1010	276	304	6			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	VAL	GLU	CONFLICT	UNP P31697
C	18	VAL	GLU	CONFLICT	UNP P31697
E	18	VAL	GLU	CONFLICT	UNP P31697
G	18	VAL	GLU	CONFLICT	UNP P31697
I	18	VAL	GLU	CONFLICT	UNP P31697
K	18	VAL	GLU	CONFLICT	UNP P31697
M	18	VAL	GLU	CONFLICT	UNP P31697
O	18	VAL	GLU	CONFLICT	UNP P31697

- Molecule 2 is a protein called FIMH PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	D	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	F	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	H	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	J	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	L	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	N	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			
2	P	279	Total	C	N	O	S	0	0	0
			2052	1297	342	409	4			

- Molecule 3 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			12	6	6		
3	D	1	Total	C	O	0	0
			12	6	6		
3	F	1	Total	C	O	0	0
			12	6	6		
3	H	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	C	O	0	0
			12	6	6		
3	J	1	Total	C	O	0	0
			12	6	6		
3	N	1	Total	C	O	0	0
			12	6	6		
3	P	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	34	Total	O	0	0
			34	34		
4	B	82	Total	O	0	0
			82	82		
4	C	37	Total	O	0	0
			37	37		
4	D	69	Total	O	0	0
			69	69		
4	E	39	Total	O	0	0
			39	39		
4	F	78	Total	O	0	0
			78	78		
4	G	39	Total	O	0	0
			39	39		
4	H	73	Total	O	0	0
			73	73		
4	I	2	Total	O	0	0
			2	2		
4	J	9	Total	O	0	0
			9	9		
4	K	4	Total	O	0	0
			4	4		
4	L	8	Total	O	0	0
			8	8		
4	M	2	Total	O	0	0
			2	2		
4	N	8	Total	O	0	0
			8	8		
4	O	4	Total	O	0	0
			4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CHAPERONE PROTEIN FIMC




• Molecule 1: CHAPERONE PROTEIN FIMC




• Molecule 1: CHAPERONE PROTEIN FIMC








● Molecule 1: CHAPERONE PROTEIN FIMC




● Molecule 1: CHAPERONE PROTEIN FIMC




● Molecule 1: CHAPERONE PROTEIN FIMC




- Molecule 1: CHAPERONE PROTEIN FIMC




- Molecule 1: CHAPERONE PROTEIN FIMC




- Molecule 2: FIMH PROTEIN




- Molecule 2: FIMH PROTEIN




• Molecule 2: FIMH PROTEIN




• Molecule 2: FIMH PROTEIN




• Molecule 2: FIMH PROTEIN







• Molecule 2: FIMH PROTEIN




• Molecule 2: FIMH PROTEIN





• Molecule 2: FIMH PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.08Å 138.13Å 215.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.68 – 2.79 43.68 – 2.79	Depositor EDS
% Data completeness (in resolution range)	98.9 (43.68-2.79) 99.0 (43.68-2.79)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.81Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.239 , 0.280 0.242 , 0.281	Depositor DCC
R_{free} test set	9936 reflections (11.14%)	DCC
Wilson B-factor (Å ²)	59.1	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 68.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.477 for k,h,-l 0.476 for -k,-h,-l 0.477 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	29775	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1625	0.74	0/2209
1	C	0.42	0/1625	0.74	0/2209
1	E	0.42	0/1625	0.73	0/2209
1	G	0.42	0/1625	0.74	0/2209
1	I	0.29	0/1625	0.58	0/2209
1	K	0.29	0/1625	0.59	0/2209
1	M	0.29	0/1625	0.59	0/2209
1	O	0.29	0/1625	0.59	0/2209
2	B	0.48	0/2097	0.76	0/2881
2	D	0.48	0/2097	0.76	0/2881
2	F	0.48	0/2097	0.76	0/2881
2	H	0.48	0/2097	0.76	0/2881
2	J	0.31	0/2097	0.59	0/2881
2	L	0.31	0/2097	0.59	0/2881
2	N	0.31	0/2097	0.59	0/2881
2	P	0.31	0/2097	0.59	0/2881
All	All	0.38	0/29776	0.67	0/40720

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1596	0	1639	133	0
1	C	1596	0	1639	142	0
1	E	1596	0	1639	125	0
1	G	1596	0	1639	135	0
1	I	1596	0	1639	149	0
1	K	1596	0	1639	149	0
1	M	1596	0	1639	152	0
1	O	1596	0	1639	152	0
2	B	2052	0	2007	141	0
2	D	2052	0	2007	142	0
2	F	2052	0	2007	135	0
2	H	2052	0	2007	141	0
2	J	2052	0	2007	200	0
2	L	2052	0	2007	205	0
2	N	2052	0	2007	203	0
2	P	2052	0	2007	208	0
3	B	12	0	12	0	0
3	D	12	0	12	0	0
3	F	12	0	12	0	0
3	H	12	0	12	0	0
3	J	12	0	12	4	0
3	L	12	0	12	2	0
3	N	12	0	12	1	0
3	P	12	0	12	2	0
4	A	34	0	0	3	0
4	B	82	0	0	3	0
4	C	37	0	0	1	0
4	D	69	0	0	4	0
4	E	39	0	0	0	0
4	F	78	0	0	4	0
4	G	39	0	0	3	0
4	H	73	0	0	4	0
4	I	2	0	0	0	0
4	J	9	0	0	1	0
4	K	4	0	0	0	0
4	L	8	0	0	1	0
4	M	2	0	0	0	0
4	N	8	0	0	0	0
4	O	4	0	0	0	0
4	P	7	0	0	0	0
All	All	29775	0	29264	2407	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

The worst 5 of 2407 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:ARG:HB3	1:C:141:THR:HG22	1.26	1.14
2:F:126:ILE:HD12	2:F:150:ALA:HB2	1.33	1.10
2:P:224:GLN:HE21	2:P:231:ILE:HG21	1.16	1.10
2:H:126:ILE:HD12	2:H:150:ALA:HB2	1.34	1.09
2:D:126:ILE:HD12	2:D:150:ALA:HB2	1.33	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/205 (99%)	163 (80%)	29 (14%)	11 (5%)	2	6
1	C	203/205 (99%)	164 (81%)	28 (14%)	11 (5%)	2	6
1	E	203/205 (99%)	163 (80%)	29 (14%)	11 (5%)	2	6
1	G	203/205 (99%)	164 (81%)	28 (14%)	11 (5%)	2	6
1	I	203/205 (99%)	162 (80%)	29 (14%)	12 (6%)	2	5
1	K	203/205 (99%)	162 (80%)	29 (14%)	12 (6%)	2	5
1	M	203/205 (99%)	162 (80%)	29 (14%)	12 (6%)	2	5
1	O	203/205 (99%)	163 (80%)	28 (14%)	12 (6%)	2	5
2	B	277/279 (99%)	244 (88%)	24 (9%)	9 (3%)	5	16
2	D	277/279 (99%)	243 (88%)	24 (9%)	10 (4%)	4	13
2	F	277/279 (99%)	244 (88%)	23 (8%)	10 (4%)	4	13
2	H	277/279 (99%)	243 (88%)	24 (9%)	10 (4%)	4	13
2	J	277/279 (99%)	195 (70%)	60 (22%)	22 (8%)	1	2
2	L	277/279 (99%)	195 (70%)	60 (22%)	22 (8%)	1	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	N	277/279 (99%)	195 (70%)	60 (22%)	22 (8%)	1	2
2	P	277/279 (99%)	196 (71%)	59 (21%)	22 (8%)	1	2
All	All	3840/3872 (99%)	3058 (80%)	563 (15%)	219 (6%)	2	5

5 of 219 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	SER
1	A	119	LYS
1	A	181	ALA
1	A	183	SER
2	B	175	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/176 (100%)	160 (91%)	16 (9%)	11	31
1	C	176/176 (100%)	159 (90%)	17 (10%)	9	27
1	E	176/176 (100%)	160 (91%)	16 (9%)	11	31
1	G	176/176 (100%)	160 (91%)	16 (9%)	11	31
1	I	176/176 (100%)	165 (94%)	11 (6%)	21	51
1	K	176/176 (100%)	164 (93%)	12 (7%)	18	47
1	M	176/176 (100%)	165 (94%)	11 (6%)	21	51
1	O	176/176 (100%)	165 (94%)	11 (6%)	21	51
2	B	226/226 (100%)	208 (92%)	18 (8%)	14	38
2	D	226/226 (100%)	208 (92%)	18 (8%)	14	38
2	F	226/226 (100%)	207 (92%)	19 (8%)	13	35
2	H	226/226 (100%)	207 (92%)	19 (8%)	13	35
2	J	226/226 (100%)	214 (95%)	12 (5%)	26	59
2	L	226/226 (100%)	214 (95%)	12 (5%)	26	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	N	226/226 (100%)	214 (95%)	12 (5%)	26	59
2	P	226/226 (100%)	213 (94%)	13 (6%)	23	55
All	All	3216/3216 (100%)	2983 (93%)	233 (7%)	17	43

5 of 233 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	23	THR
2	H	126	ILE
1	O	85	MET
1	G	47	ARG
1	G	200	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 181 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	255	ASN
2	J	211	ASN
2	P	70	ASN
1	I	28	ASN
1	I	126	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MAN	B	1500	-	12,12,12	0.40	0	17,17,17	0.39	0
3	MAN	D	1601	-	12,12,12	0.36	0	17,17,17	0.52	0
3	MAN	F	1502	-	12,12,12	0.43	0	17,17,17	0.59	0
3	MAN	H	1603	-	12,12,12	0.44	0	17,17,17	0.51	0
3	MAN	J	1605	-	12,12,12	0.28	0	17,17,17	0.35	0
3	MAN	L	1504	-	12,12,12	0.29	0	17,17,17	0.44	0
3	MAN	N	1506	-	12,12,12	0.24	0	17,17,17	0.44	0
3	MAN	P	1607	-	12,12,12	0.28	0	17,17,17	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	B	1500	-	-	0/2/22/22	0/1/1/1
3	MAN	D	1601	-	-	0/2/22/22	0/1/1/1
3	MAN	F	1502	-	-	0/2/22/22	0/1/1/1
3	MAN	H	1603	-	-	0/2/22/22	0/1/1/1
3	MAN	J	1605	-	-	0/2/22/22	0/1/1/1
3	MAN	L	1504	-	-	0/2/22/22	0/1/1/1
3	MAN	N	1506	-	-	0/2/22/22	0/1/1/1
3	MAN	P	1607	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	1605	MAN	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	1504	MAN	2	0
3	N	1506	MAN	1	0
3	P	1607	MAN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	205/205 (100%)	0.51	10 (4%)	30	20	20, 55, 108, 132	0
1	C	205/205 (100%)	0.48	12 (5%)	23	15	20, 57, 108, 130	0
1	E	205/205 (100%)	0.41	3 (1%)	74	67	20, 59, 108, 129	0
1	G	205/205 (100%)	0.47	16 (7%)	14	7	19, 58, 110, 132	0
1	I	205/205 (100%)	3.93	145 (70%)	0	0	20, 168, 190, 196	0
1	K	205/205 (100%)	4.77	171 (83%)	0	0	20, 171, 195, 198	0
1	M	205/205 (100%)	4.55	165 (80%)	0	0	20, 170, 193, 198	0
1	O	205/205 (100%)	3.98	148 (72%)	0	0	20, 168, 193, 198	0
2	B	279/279 (100%)	0.30	5 (1%)	69	60	17, 40, 79, 114	0
2	D	279/279 (100%)	0.30	7 (2%)	58	47	18, 41, 80, 116	0
2	F	279/279 (100%)	0.23	4 (1%)	75	69	18, 41, 83, 112	0
2	H	279/279 (100%)	0.28	5 (1%)	69	60	17, 42, 83, 114	0
2	J	279/279 (100%)	2.04	105 (37%)	0	0	51, 112, 192, 200	0
2	L	279/279 (100%)	2.36	124 (44%)	0	0	56, 113, 190, 200	0
2	N	279/279 (100%)	2.60	143 (51%)	0	0	53, 115, 193, 200	0
2	P	279/279 (100%)	2.14	109 (39%)	0	0	50, 112, 191, 200	0
All	All	3872/3872 (100%)	1.75	1172 (30%)	0	0	17, 85, 186, 200	0

The worst 5 of 1172 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	48	PHE	17.8
1	I	102	THR	16.6
1	M	102	THR	15.5
2	L	270	SER	15.4
1	K	95	LYS	15.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MAN	D	1601	12/12	0.98	0.19	0.07	28,36,41,41	0
3	MAN	P	1607	12/12	0.90	0.26	0.02	110,112,116,117	0
3	MAN	H	1603	12/12	0.96	0.17	-0.03	28,34,39,39	0
3	MAN	F	1502	12/12	0.95	0.19	-0.04	29,35,41,41	0
3	MAN	J	1605	12/12	0.88	0.25	-0.28	96,100,105,106	0
3	MAN	B	1500	12/12	0.97	0.15	-0.80	30,35,39,39	0
3	MAN	L	1504	12/12	0.81	0.23	-0.80	96,99,103,106	0
3	MAN	N	1506	12/12	0.91	0.18	-1.22	90,94,100,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.